

*-ZERO DIVISORS AND *-PRIME IDEALS

Usama A. Aburawash and Khadija B. Sola

Department of Mathematics, Faculty of Science
Alexandria University, Alexandria, Egypt
aburawash@sci.alex.edu.eg

Abstract

Throughout this note we introduce the concept of *-zero divisors in rings with involution and its correlation with the concept of zero divisors in rings without involution. Moreover, some related definitions; such as *-completely prime ideals and rings and *-cancellation laws are introduced. Nevertheless, we characterize *-prime and *-completely prime ideals using *-zero divisors.

By a ring we mean an associative ring. A ring A is said to be an *involution ring* if on A there is defined a unary operation (called *involution*) $*$ subject to the identities $a^{**} = a$, $(a + b)^* = a^* + b^*$ and $(ab)^* = b^*a^*$, for all $a, b \in A$. In other words, the involution is an anti-isomorphism of order 2 on A . For a commutative ring A , it is evident that the identity mapping of A onto A is an involution on A (see [1]-[4]). Considering the category of involution rings, all morphisms (and also embeddings) must preserve involution. So we are looking here for a particular concept for zero divisors that works in the category of involution rings.

If the ideal I of A is closed under involution; that is $I^{(*)} = \{a^* \in A \mid a \in I\} \subseteq I$, then it is called a **-ideal* of A and will be denoted by $I \triangleleft^* A$.

We start by defining *-zero divisors for an involution ring A .

Definition 1 A nonzero element $a \in A$ is said to be a *-zero divisor if there exists a nonzero element $b \in A$ such that $ab = 0$ and $a^*b = 0$.

Remark 2 If we start by defining left *-zero divisor as in definition 1, we get $b^*a^* = 0$ and $b^*a = 0$ which mean that a is a right *-zero divisor, too. By reversing the roles, a right *-zero divisor is also a left *-zero divisor. Thus we have only the concept of *-zero divisor, as one expects in the category of involution rings.

Key words: zero divisors, involution, *-cancellation, *-completely prime ideal
2000 AMS Mathematics Subject Classification: 16W10, 16U10, 16U30.

Clearly, a *-zero divisor is a zero divisor, but the converse is not always true as it is obvious from the following example.

Example 3 Consider the direct sum $R = A \oplus A^{op}$, where A is an integral domain and A^{op} is its opposite domain. R is an involution ring under the exchange involution given by $(a, b)^* = (b, a)$ for all $(a, b) \in R$. For any $0 \neq a \in A$, the element $(a, 0)$ of R is a zero divisor since $(a, 0)(0, b) = 0 = (0, b)(a, 0)$ for every $0 \neq b \in A$. Because neither a nor b are zero divisors, from $(0, a)(0, b) \neq (0, 0)$, we conclude that $(a, 0)$ is not a *-zero divisor.

In particular, if a is a symmetric ($a^* = a$) or a skew symmetric ($a^* = -a$) element, then a is a zero divisor if and only if it is a *-zero divisor. Moreover, we can construct symmetric or skew symmetric *-zero divisors from given *-zero divisors as in the following result.

Proposition 4 Let A be an involution ring and $a \in A$. If a is a *-zero divisor, then there exists a (nonzero) symmetric or skew symmetric *-zero divisor in A .

Proof If a is a symmetric or skew symmetric element, then we are done. If a is not symmetric, then $a - a^* \neq 0$ is a skew symmetric element in A such that $(a - a^*)b = ab - a^*b = 0$ and $(a - a^*)^*b = (a^* - a)b = a^*b - ab = 0$, with an appropriately chosen $b \in A$. \square

Nevertheless, the next example shows that there are zero divisors which are *-zero divisors.

Example 5 In the involution ring of all 3×3 matrices over the integers Z with the transpose as involution, the element $a = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ is both zero and *-zero divisor. In fact the matrix $b = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$ satisfies $ab = ba = 0$ and $ab = a^*b = 0$.

Definition 6 A commutative involution ring without *-zero divisors is said to be a *-integral domain.

Since a commutative involution ring is an integral domain if it has no zero divisors, so it has also no *-zero divisors and consequently it is a *-integral domain. Moreover, each involution division ring is a *-integral domain.

Next, we define the *-cancellation law to work with *-zero divisors as follows.

Definition 7 We say that The *-cancellation law holds in an involution ring A if $ab = ac$ and $a^*b = a^*c$ imply $b = c$, for any $0 \neq a \in A$.

Again, if one defines *left *-cancellation law* to be hold in A as in Definition 7, we can easily show that *the right *-cancellation law* holds also in A . Therefore, we have only the *-cancellation law as one expects.

It is obvious that if the left (right) cancellation law holds in an involution ring A , then the *-cancellation law holds in A , too.

Remind that an ideal P of a ring A is called a *completely prime ideal* if $ab \in P$ implies $a \in P$ or $b \in P$ for all $a, b \in A$ (see for instance [5]).

Now, we give the involutive version of this definition.

Definition 8 An ideal P of an involution ring A is called a **-completely prime ideal* if $ab \in P$ and $a^*b \in P$ imply $a \in P$ or $b \in P$ for all $a, b \in A$. The ring A is said to be a **-completely prime ring* if the zero ideal is a *-completely prime ideal.

It is evident that in an involution ring A , a completely prime ideal is *-completely prime, too.

From the definition, it follows that the ring A is *-completely prime if and only if it has no *-zero divisors. We remind also that a ring A is completely prime if and only if it has no zero divisors. By this remark, a completely prime involution ring A is also *-completely prime, since A has no zero divisors implies that A has no *-zero divisors.

Following [3], an ideal P of an involution ring A is called a **-prime ideal* if $JK \subseteq P$ implies $J \subseteq P$ or $K \subseteq P$, for any $J, K \triangleleft^* A$. An involution ring A is a **-prime ring* if the zero ideal is a *-prime ideal. By the way, Birkenmeier and Groenewald gave in [3] the following equivalents for *-primeness of ideals.

Proposition 9 ([3], Proposition 5.4) *Let A be an involution ring and $P \triangleleft^* A$. Then the following conditions are equivalent:*

- (i) P is a *-prime *-ideal of A .
- (ii) If $a, b \in A$ such that $aAb \subseteq P$ and $a^*Ab \subseteq P$, then $a \in P$ or $b \in P$.
- (iii) If $I \triangleleft A$ and $K \triangleleft^* A$ such that $IK \subseteq P$, then $I \subseteq P$ or $K \subseteq P$.

We start our results by a classical one which gives the relation between the *-cancellation law and *-zero divisors.

Proposition 10 *Let A be an involution ring. Then the *-cancellation law holds in A if and only if A has no *-zero divisors.*

Proof Suppose that the *-cancellation law hold in A . If $0 \neq a \in A$ is such that $ab = 0, a^*b = 0$, then $b = 0$ follows and consequently A has no *-zero divisors. Conversely, let A have no *-zero divisors. For $0 \neq a \in A$, if $ab = ac$ and $a^*b = a^*c$, then $a(b - c) = 0$ and $a^*(b - c) = 0$ which forces $b - c = 0$. Thus $b = c$ and the *-cancellation law holds in A . \square

For *-prime rings without nonzero nilpotent elements, we claim that they have no *-zero divisors.

Proposition 11 *If A is a *-prime ring having no nonzero nilpotent elements, then A has no *-zero divisors.*

Proof Let $0 \neq a, b \in A$ be such that $ab = 0, a^*b = 0$. Then $(ba)^2 = b(ab)a = 0$. Since A has no nonzero nilpotent elements, it follows that $ba = 0$. Thus for all $x \in A$, we have $(axb)^2 = ax(ba)xb = 0$, whence $axb = 0$ and consequently $aAb = 0$. Similarly, we get $a^*Ab = 0$. Because A is *-prime, we deduce from Proposition 9 that $b = 0$, from which A has no *-zero divisors. \square

From the definitions, it is easy to check that a *-completely prime *-ideal of A is also a *-prime *-ideal. The converse is true only in particular cases; for instance if A possesses identity. For commutative involution rings, we have the following equivalences.

Theorem 12 *Let A be a commutative ring with involution and $P \triangleleft^* A$. Then the following conditions are equivalent:*

- (i) P is a *-prime *-ideal.
- (ii) P is a *-completely prime *-ideal.
- (iii) The factor ring A/P is a *-integral domain.

Proof (i) \rightarrow (ii). Let $a, b \in A$ such that $ab \in P$ and $a^*b \in P$. Then $aAb \subseteq P$ and $a^*Ab \subseteq P$. Hence, by Proposition 9, $a \in P$ or $b \in P$ and consequently P is a *-completely prime *-ideal.

(ii) \rightarrow (iii). A/P is commutative because A is commutative. Since P is a *-completely prime *-ideal, then $ab \in P$ and $a^*b \in P$ imply $a \in P$ or $b \in P$ for all $a, b \in A$. In other words, $(a + P)(b + P) = P$ and $(a + P)^*(b + P) = P$ imply $a + P = P$ or $b + P = P$, whence A/P is a *-integral domain.

(iii) \rightarrow (i). Suppose that $aAb \subseteq P$ and $a^*Ab \subseteq P$. By the commutativity of A , we get $(ab)b \in P$, $(ab)^*b \in P$ and $(a^*b)b \in P$, $(a^*b)^*b \in P$. Since A/P has no *-zero divisors, it follows that $ab \in P$ or $b \in P$ and $a^*b \in P$ or $b \in P$. If $b \notin P$, then $ab \in P$ and $a^*b \in P$, from which $a \in P$. Thus P is a *-prime *-ideal, by Proposition 9. \square

Proposition 13 *For a commutative ring A with involution, the following are true:*

- (i) The set $K = \{\text{all *-zero divisors of } A\} \cup \{0\}$ is a *-ideal of A .
- (ii) The factor ring A/K is a *-integral domain.

Proof (i) Let $a, b \in K$ and $r \in A$, then there exist $c, d \in A$ such that $ac = a^*c = 0$ and $bd = b^*d = 0$. Hence $(a-b)cd = 0$, $(a-b)^*cd = (a^*-b^*)cd = 0$ and $rac = 0$, $(ra)^*c = 0$. Thus $a - b, ra \in K$. Moreover $a^* \in K$, since $a^*c = a^{**}c = ac = 0$.

(ii) Since A/K is commutative and has no *-zero divisors, it is a *-integral domain. \square

The following proposition gives a necessary condition for an element in the center of a *-ideal to be in the center of the ring.

Proposition 14 *Let N be a $*$ -ideal of an involution ring A and $c \in C(N)$; the center of N . If c is not a $*$ -zero divisor, then $c \in C(A)$.*

Proof $C(N) = \{n \in N \mid nx = xn, \text{ for all } x \in N\}$ is a $*$ -subring of A , since for $n \in C(N)$, $x \in N$, we have $nx^* = x^*n$. Hence $n^*x = xn^*$ and $n^* \in C(N)$. Now for any $y \in A$, we have $cy, yc, c^*, c^*y, yc^* \in N$. Hence

$$c(cy - yc) = c(cy) - c(yc) = c^*yc - c^*yc = 0$$

and

$$c^*(cy - yc) = c^*(cy) - c^*(yc) = (c^*c)y - c^*yc = c(c^*y) - c^*yc = c^*yc - c^*yc = 0.$$

But c is not a $*$ -zero divisor, whence $cy - yc = 0$ and $c \in C(A)$ follows. \square

Finally, since an involution ring without zero divisors having no $*$ -zero divisors, we have the following immediate result from Proposition 3 in [1].

Proposition 15 *Every involution ring A without zero divisors is embeddable as a $*$ -ideal (up to isomorphism) into one and only one involution ring \bar{A}^1 with identity and without $*$ -zero divisors such that \bar{A}^1 is a minimal $*$ -extension of A possessing identity.*

References

- [1] U. A. Aburawash, *On embedding of involution rings*, Math. Pannonica, **82**, (1997), 245-250.
- [2] U. A. Aburawash, *On involution rings*, East-West J.Math., **2**(2)(2000), 109-126.
- [3] G. F. Birkenmeier and N. J. Groenewald, *Prime ideals in rings with involution*, Quaestiones in Mathematica, 20 (1997), 591-603.
- [4] I. N. Herstein, "Rings with Involution", Univ. Chicago Press, 1976.
- [5] A. Kertész, "Lectures on Artinian Rings", Akad. Kiadó, Budapest, 1987.