

ON REVERSIBILITY OF RINGS WITH INVOLUTION

Usama A. Aburawash and Muna E. Abdulhafed⁰

*Department of Mathematics and Computer Science
Faculty of Science, Alexandria University, Alexandria, Egypt
e-mail: aburawash@alexu.edu.eg*

*Faculty of Arts and Science, Azzaytuna University, Tarhunah, Libya
muna.am2016@gmail.com*

Abstract

Let R be a ring with involution $*$. We give the notion of central $*$ -reversible $*$ -rings which generalizes weakly $*$ -reversible $*$ -rings. Moreover, we introduce the class of weakly $*$ -rings which is a generalization of central $*$ -reversible $*$ -rings and investigate their properties. Further, a generalization of the class of quasi- $*$ -IFP $*$ -rings is given; namely weakly quasi- $*$ -IFP $*$ -rings. Since every $*$ -reversible $*$ -ring is central $*$ -reversible, we give sufficient conditions for central $*$ -reversible, weakly $*$ -reversible and weakly quasi- $*$ -IFP $*$ -rings to be $*$ -reversible and some examples are given to illustrate these situations. Finally, we show that the properties of $*$ -reversible, central $*$ -reversible, weakly $*$ -reversible and weakly quasi- $*$ -IFP can be transfer to some extensions of the $*$ -ring.

1 Introduction

Throughout this paper, a ring will always mean an associative ring with unity unless otherwise stated. A ring R is said to be $*$ -ring if on R there is defined an involution $*$; that is an anti-isomorphism of order two. The right annihilator of the nonempty set A of R is denoted by $r_R(A)$ and the right $*$ -annihilator of A is denoted by $r_{*R}(A) = \{x \in R \mid Ax = Ax^* = 0\}$. If there is no ambiguity, we omit the subsuffix R . A $*$ -ideal (self-adjoint) I of R is an ideal closed under involution. A self adjoint idempotent; $e^2 = e = e^*$, is

Key words: Involution; $*$ -Reversible; Central $*$ -reversible; Weakly $*$ -reversible; Quasi- $*$ -IFP; Weakly $*$ -IFP; Weakly quasi- $*$ -IFP $*$ -rings.

2010 AMS Mathematics classification:

called projection. A nonzero element a of a $*$ -ring R is called $*$ -zero divisor if $ab = 0 = a^*b$, for some nonzero element $b \in R$ and R is $*$ -domain if it has no nonzero $*$ -zero divisors, from [6]. A $*$ -ring R is said to be *Abelian* ($*$ -Abelian) if every idempotent (projection) of R is center. A $*$ -ring R is reduced if it has no nonzero nilpotent elements. A ring R is called *semicommutative* or has (IFP) if for all $a, b \in R$, $ab = 0$ implies $aRb = 0$ (equivalently $r(a)$ is an ideal of R for all $a \in R$) (see [10]). A $*$ -ring R is said to have $*$ -IFP if for all $a, b \in R$, $ab = 0$ implies $aRb^* = 0$ (equivalently $r(a)$ is a $*$ -ideal of R for all $a \in R$) (see [4]). From [13], recall a ring R is *weakly semicommutative* if for all $a, b \in R$, $ab = 0$ implies arb is a nilpotent element for each $r \in R$. By [7], a ring R is called *reversible* if for all $a, b \in R$, $ab = 0$ implies $ba = 0$. According to [3], a $*$ -ring R is called $*$ -reversible if for all $a, b \in R$, $ab = 0 = ab^*$ implies $ba = 0$, and R has *quasi- $*$ -IFP* if for all $a, b \in R$, $ab = ab^* = 0$ implies $aRb = 0$. From [5], an element a of a $*$ -ring R is called $*$ -nilpotent if $a^m = (aa^*)^n = 0$, for some positive integers m and n . R is $*$ -reduced if it has no nonzero $*$ -nilpotent elements. Following [9], a $*$ -ring R is called *Baer $*$ -ring* if the right annihilator of every nonempty subset of R is generated, as a right ideal, by a projection. By [5], a $*$ -ring R is called $*$ -Baer $*$ -ring if the $*$ -right annihilator of every nonempty subset of R is generated, as a biideal, by a projection. From [8] a ring R is *central reversible* rings if for all $a, b \in R$, $ab = 0$ implies ba belongs to the center of R and a ring R is called *weakly reversible* if $ab = 0$ implies $Rbra$ is nil left ideal of R , for all $a, b, r \in R$, from [11]. The natural numbers and the integers will be denoted by $\mathbb{N}$ and $\mathbb{Z}$, respectively. $M_n(R)$ will denote the full matrix ring of all $n \times n$ matrices over the ring R , while $T_n(R)$ ($T_{nE}(R)$) will denote the $n \times n$ upper triangular matrix ring (with equal diagonal elements) over R .

In this paper, we introduce central and weakly $*$ -reversible $*$ -rings, both are proper generalizations of $*$ -reversible $*$ -rings. Moreover, the class of weakly $*$ -reversible $*$ -rings contains strictly central $*$ -reversible $*$ -rings. We also prove that central $*$ -reversible $*$ -rings are $*$ -Abelian and there exists a $*$ -Abelian $*$ -ring which is not central $*$ -reversible. Clearly $*$ -reversible $*$ -rings are quasi- $*$ -IFP and example is given to show that the converse is not true and another example shows that commutative weakly $*$ -reversible $*$ -rings do not necessarily have quasi- $*$ -IFP. It is also shown that if R is a commutative $*$ -ring, then $T_{nE}(R)$ is weakly $*$ -reversible (weakly quasi- $*$ -IFP) $*$ -ring. Moreover, weakly quasi- $*$ -IFP condition is given for $*$ -rings which generalizes quasi- $*$ -IFP. We show also that commutative weakly quasi- $*$ -IFP $*$ -rings may not be quasi- $*$ -IFP. Moreover, for a $*$ -Armendariz $*$ -ring R , we prove that R is $*$ -reversible (central $*$ -reversible) if and only if the polynomial $*$ -rings $R[x]$ is $*$ -reversible (central $*$ -reversible) if and only if the Laurent polynomial $*$ -ring $R[x; x^{-1}]$ is $*$ -reversible (central $*$ -reversible). Furthermore, it is proved that R is $*$ -reversible (central $*$ -reversible) if and only if the Dorroh extension $D(R, \mathbb{Z})$ of

R is $*$ -reversible (central $*$ -reversible). Finally, the Ore $*$ -ring R is shown to be $*$ -reversible if and only if its classical quotient Q is $*$ -reversible.

2 Central $*$ -Reversible $*$ -Rings

In this section, we introduce and study the class of central $*$ -reversible $*$ -rings, which is a generalization of $*$ -reversible $*$ -rings. We start by giving the main definition.

Definition. A $*$ -ring R is called *central $*$ -reversible* if for all $a, b \in R$, $ab = 0 = ab^*$ implies ba is central in R . Consequently, b^*a is central in R .

Clearly, a central reversible $*$ -ring is central $*$ -reversible and a $*$ -reversible $*$ -ring is central $*$ -reversible. However, the next result shows that $T_{3E}(R)$, in general, is central $*$ -reversible but not $*$ -reversible.

Proposition 1. *Let R be a commutative $*$ -ring, then the ring*

$$T_{3E}(R) = \left\{ \begin{pmatrix} a & b & c \\ 0 & a & d \\ 0 & 0 & a \end{pmatrix} \mid a, b, c, d \in R \right\}$$

with involution defined as $\begin{pmatrix} a & b & c \\ 0 & a & d \\ 0 & 0 & a \end{pmatrix}^ = \begin{pmatrix} a & d & c \\ 0 & a & b \\ 0 & 0 & a \end{pmatrix}$ is central $*$ -reversible $*$ -ring.*

Proof. Let $x = \begin{pmatrix} a_1 & b_1 & c_1 \\ 0 & a_1 & d_1 \\ 0 & 0 & a_1 \end{pmatrix}$ and $y = \begin{pmatrix} a_2 & b_2 & c_2 \\ 0 & a_2 & d_2 \\ 0 & 0 & a_2 \end{pmatrix} \in T_{3E}(R)$. If $xy = 0 = xy^*$, then we have the following equations :

$$a_1a_2 = 0 \tag{1}$$

$$a_1b_2 + b_1a_2 = 0, \quad a_1d_2 + b_1a_2 = 0 \tag{2}$$

$$a_1c_2 + b_1d_2 + c_1a_2 = 0, \quad a_1c_2 + b_1b_2 + c_1a_2 = 0 \tag{3}$$

$$a_1d_2 + d_1a_2 = 0, \quad a_1b_2 + d_1a_2 = 0. \tag{4}$$

Hence $yx = \begin{pmatrix} 0 & 0 & b_2d_1 - b_1d_2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \neq 0$, is central and consequently $T_{3E}(R)$

is central $*$ -reversible. On the other hand, $T_{3E}(R)$ is not $*$ -reversible, since $yx \neq 0$, while the converse is clear from [3, Example 3.8]. $\square$

In general, **Proposition 1** is not true for $n \geq 4$ which is clear from the following example.

Example 1. Consider the $*$ -ring $T_{4E}(\mathbb{Z})$ with the involution $*$ defined as:

$$\begin{pmatrix} a & a_{12} & a_{13} & a_{14} \\ 0 & a & a_{23} & a_{24} \\ 0 & 0 & a & a_{34} \\ 0 & 0 & 0 & a \end{pmatrix}^* = \begin{pmatrix} a & a_{34} & a_{24} & a_{14} \\ 0 & a & a_{23} & a_{13} \\ 0 & 0 & a & a_{12} \\ 0 & 0 & 0 & a \end{pmatrix}.$$

The matrices $A = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$ satisfies $AB =$

$0 = AB^*$, but $BA = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$ is not central and so $T_{4E}(\mathbb{Z})$ is not

central $*$ -reversible.

It is clear that each central reversible is central $*$ -reversible. However, the converse is true when the ring has $*$ -IFP as shown in the next result.

Proposition 2. *Let R be a $*$ -ring. If R is central $*$ -reversible and has $*$ -IFP, then R is central reversible.*

Proof. Obvious, since $ab = 0$, implies $aRb^* = 0$, by $*$ -IFP property, and R is central reversible. $\square$

Recall that a $*$ -ring R is $*$ -semiprime if and only if it is semiprime (see ([1])). Next, we give some particular conditions for a central $*$ -reversible $*$ -ring to be $*$ -reversible.

Proposition 3. *A semiprime central $*$ -reversible $*$ -ring is $*$ -reversible.*

Proof. Assume that R is a semiprime central $*$ -reversible $*$ -ring. If $ab = ab^* = 0$, then ba is central and consequently $baRba = 0$. Form semiprimeness, we get $ba = 0$ and so R is $*$ -reversible. $\square$

Proposition 4. *If R is a $*$ -Baer and central $*$ -reversible $*$ -ring, then R is $*$ -reversible.*

Proof. Let R be a $*$ -Baer $*$ -ring and $ab = 0 = ab^*$, then there exists a projection $e \in R$ such that $r_*(a) = eRe$. We have $ae = 0$ and $b = ebe = eb$, since $b \in r_*(a) = eRe$. Hence $ba = eba = bae = 0$, since ba is central, and so R is $*$ -reversible. $\square$

Since each Bear $*$ -ring is $*$ -Bear, we have the following corollary.

Corollary 1. *If R is a Baer and central $*$ -reversible $*$ -ring, then R is $*$ -reversible.*

Furthermore, the class of central $*$ -reversible $*$ -rings is clearly closed under direct sums (with changeless involution) and under taking $*$ -subrings by [3], since every $*$ -reversible $*$ -ring is central $*$ -reversible.

Proposition 5. *The class of central $*$ -reversible $*$ -ring is closed under direct sums and under taking $*$ -subrings.*

Proposition 6. *Let R be a $*$ -ring and e be a central projection of R . Then eR and $(1 - e)R$ are $*$ -reversible if and only if R is $*$ -reversible.*

Proof. It suffices to show the necessity by [3, Proposition 3.15]. Let $ab = ab^* = 0$ with $a, b \in R$, then $eab = eab^* = 0$ and $(1 - e)ab = (1 - e)ab^* = 0$. By assumption, we have $bea = 0$ and $b(1 - e)a = 0$. Hence $ba = bea + [b(1 - e)a] = 0$ and so R is $*$ -reversible. $\square$

By a similar proof as **Proposition 6**, and using **Proposition 5**, the following corollary is immediate.

Corollary 2. *Let R be a $*$ -ring and e be a central projection of R . Then eR and $(1 - e)R$ are central $*$ -reversible if and only if R is central $*$ -reversible.*

Recall that a $*$ -ideal I of a $*$ -ring R is $*$ -nil if each element of I is $*$ -nilpotent.

Obviously, each $*$ -nil ideal is nil. The following example shows that the converse is not always true.

Example 2. For the $*$ -ring $R = M_2(\mathbb{Z})$ of all 2×2 matrices over the integers $\mathbb{Z}$ with transpose of matrices as involution, the nonzero elements of the form $\begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix}$ are all nilpotent but not $*$ -nilpotent, since $\begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix}^2 = 0$ but $\begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ x & 0 \end{pmatrix} = \begin{pmatrix} x^2 & 0 \\ 0 & 0 \end{pmatrix} \neq 0$

We note that the homomorphic image of a central $*$ -reversible $*$ -ring need not be central $*$ -reversible as seen from the following example.

Example 3. Let D be a $*$ -division ring, $R = D[x, y]$ and $I = \langle xy \rangle$, where $xy \neq yx$. Since R is $*$ -domain, R is central $*$ -reversible. On the other hand, $(x+I)(y+I)$ and $(x+I)^*(y+I) = (x+I)(y+I)$ are both zero. But $(y+I)(x+I)$ is not central in R/I , hence R/I is not central $*$ -reversible.

Moreover, the next example shows that if the homomorphic image of a $*$ -ring R is central $*$ -reversible, then R need not be central $*$ -reversible.

Example 4. Let $R = \begin{pmatrix} \mathbb{F} & \mathbb{F} \\ 0 & \mathbb{F} \end{pmatrix}$, where $\mathbb{F}$ is a field, with the adjoint involution $*$ definition by $\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}^* = \begin{pmatrix} c & -b \\ 0 & a \end{pmatrix}$ for all $a, b, c \in \mathbb{F}$. Consider the $*$ -ideal $I = \begin{pmatrix} 0 & \mathbb{F} \\ 0 & 0 \end{pmatrix}$ of R . Then R/I is central $*$ -reversible, because of the commutativity property of R/I . For $A = \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \in R$ where $B^* = \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix} \in R$, we have $AB = 0 = AB^*$. Consider $C = \begin{pmatrix} c_1 & c_2 \\ 0 & c_3 \end{pmatrix} \in R$ with $c_1 \neq c_3$. It is clear that $CBA \neq BAC$ and therefore R is not central $*$ -reversible.

Our next endeavour is to give a condition on the homomorphic image of a $*$ -ring to be central $*$ -reversible. Recall that a $*$ -ring R is called *unit-central*, if all unit elements of R are central in R . Moreover, we show that every unit central $*$ -ring is $*$ -Abelian.

Proposition 7. *Let R be a unit-central $*$ -ring. If I is a $*$ -nil ideal of R , then R/I is central $*$ -reversible.*

Proof. Let $a, b \in R$ with $(a + I)(b + I) = (a + I)(b + I)^* = I$. Then $ab \in I$, $ab^* \in I$ and so there exists a positive integers m, n, p and q such that $(ab)^m = 0$, $((ab)(ab)^*)^n = 0$, $(ab^*)^p = 0$ and $((ab^*)(ab^*)^*)^q = 0$. It follows that $(ba)^{m+1} = 0$, whence $1 - ba$ is unit and so central by hypothesis. Thus $rba = bar$ for any $r \in R$ and therefore $(b + I)(a + I)$ is central in R/I . $\square$

Since each $*$ -reversible $*$ -ring is central $*$ -reversible and each $*$ -domain is $*$ -reversible, by [3, Example 3.2], we have immediately the following corollary.

Corollary 3. *Every $*$ -domain is a central $*$ -reversible $*$ -ring.*

The converse of **Corollary 3** is not true by **Example 4**. However, the converse is true for $*$ -prime $*$ -rings as follows.

Proposition 8. *Let R be a $*$ -ring. Then R is $*$ -prime and central $*$ -reversible if and only if it is $*$ -domain.*

Proof. Let R be $*$ -prime and central $*$ -reversible and $ab = ab^* = 0$ for some $a, b \in R$. We have $rab = rab^* = 0$ for every $r \in R$ and so bra and b^*ra are central. Since $bratb = 0$ and $bratb^* = 0$ for all $t \in R$, then $a = 0$ or $b = 0$ and R is a $*$ -domain. The converse is obvious by **Corollary 3**. $\square$

It is well known from [3, Corollary 3.7] that every $*$ -reversible $*$ -ring is $*$ -Abelian. Similarly, we have the same result for central $*$ -reversible case.

Proposition 9. *A central $*$ -reversible $*$ -ring R is $*$ -Abelian.*

Proof. Let $e^2 = e = e^* \in R$. for any $r \in R$, $(re - ere)(1 - e) = (re - ere)(1 - e)^* = 0$ implies $(1 - e)(re - ere) = re - ere$ is central. Commuting $re - ere$ by e we get $re - ere = 0$. Similarly for any $r \in R$, $(r^*e - er^*e)(1 - e) = (r^*e - er^*e)(1 - e)^* = 0$ implies $r^*e - er^*e = 0$. Therefore $re = ere = er$ and R is $*$ -Abelian. $\square$

The next example shows that the reverse implication of **Proposition 9** is not true in general; that is there exists a $*$ -Abelian $*$ -ring which is not central $*$ -reversible, and hence is not $*$ -reversible.

Example 5. The only projections of the $*$ -ring

$R = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid a \equiv c \pmod{2}, b \equiv 0 \pmod{2}, a, b, c \in \mathbb{Z} \right\}$ under adjoint involution $*$ are $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$ and $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and so R is $*$ -Abelian. On the other hand, for $x = \begin{pmatrix} 0 & 0 \\ 0 & 2 \end{pmatrix}$ and $y = \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix} \in R$ with $xy = xy^* = 0$, we have $yx = \begin{pmatrix} 0 & 4 \\ 0 & 0 \end{pmatrix}$ is not central and so R is not central $*$ -reversible.

3 Weakly $*$ -Reversible $*$ -Rings

In this section, we introduce another generalization for $*$ -reversible; namely weakly $*$ -reversible $*$ -rings.

Definition. A $*$ -ring R is called *weakly $*$ -reversible* if for all $a, b, r \in R$, $ab = ab^* = 0$, implies $Rbra$ is a nil left (equivalently, $braR$ is a nil right) ideal of R . Consequently, Rb^*ra is a nil left (equivalently, b^*raR is a nil right) ideal of R .

Each commutative $*$ -ring is weakly reversible. Clearly, each weakly reversible $*$ -ring is weakly $*$ -reversible. The converse is true when the ring has $*$ -IFP as shown in the following.

Proposition 10. *Let R be a $*$ -ring. If R is weakly $*$ -reversible and has $*$ -IFP, then R is weakly reversible.*

Proof. Obvious, since $ab = 0$, implies $aRb^* = 0$, by the $*$ -IFP property, and R is weakly reversible. $\square$

Moreover, we can easily prove the following result.

Proposition 11. *The class of weakly $*$ -reversible $*$ -ring is closed under direct sums (with changeless involution) and under taking $*$ -subrings.*

Proposition 12. For a commutative $*$ -ring R , $T_{nE}(R)$ is a weakly $*$ -reversible $*$ -ring, with involution $*$ defined by fixing the two diagonals considering the diagonal right / left lower as symmetric ones and interchange the symmetric elements about it; that is

$$\begin{pmatrix} a & a_{12} & a_{13} & \cdots & a_{1(n-1)} & a_{1n} \\ 0 & a & a_{23} & \cdots & a_{2(n-1)} & a_{2n} \\ 0 & 0 & a & \cdots & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \cdots & a_{(n-1)n} \\ 0 & 0 & 0 & 0 & \cdots & a \end{pmatrix}^* = \begin{pmatrix} a & a_{(n-1)n} & a_{(n-2)n} & \cdots & a_{2n} & a_{1n} \\ 0 & a & a_{23} & \cdots & a_{2(n-1)} & a_{1(n-1)} \\ 0 & 0 & a & \cdots & \cdots & a_{1(n-2)} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \cdots & a_{12} \\ 0 & 0 & 0 & 0 & \cdots & a \end{pmatrix}$$

Proof. Let R be weakly $*$ -reversible and $A = \begin{pmatrix} a & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a & a_{23} & \cdots & a_{2n} \\ 0 & 0 & a & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a \end{pmatrix}$,

$$B = \begin{pmatrix} b & b_{12} & b_{13} & \cdots & b_{1n} \\ 0 & b & b_{23} & \cdots & b_{2n} \\ 0 & 0 & b & \cdots & b_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & b \end{pmatrix} \in T_{nE}(R) \text{ satisfy } AB = 0 = AB^*. \text{ Hence } ab =$$

$$0 = ab^* \text{ and Since } R \text{ is weakly } *-reversible, \text{ then for } C = \begin{pmatrix} c & c_{12} & c_{13} & \cdots & c_{1n} \\ 0 & c & c_{23} & \cdots & c_{2n} \\ 0 & 0 & c & \cdots & c_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & c \end{pmatrix}$$

$$\text{and } D = \begin{pmatrix} d & d_{12} & d_{13} & \cdots & d_{1n} \\ 0 & d & d_{23} & \cdots & d_{2n} \\ 0 & 0 & d & \cdots & d_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & d \end{pmatrix} \in T_{nE}(R), \text{ there exists } k \in \mathbb{N}, \text{ with } (cbda)^k =$$

0. Thus

$$(CBDA)^k = \begin{pmatrix} 0 & \star & \star & \cdots & \star \\ 0 & 0 & \star & \cdots & \star \\ 0 & 0 & 0 & \cdots & \star \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix} \text{ and } (CBDA)^{kn} = 0 \text{ follows and}$$

$T_{nE}(R)$ is weakly $*$ -reversible. [$\star$ denotes an element of R]

□

Next, the given example shows that there exists a weakly $*$ -reversible and quasi $*$ -IFP $*$ -ring which is not $*$ -reversible.

Example 6. Let R be a commutative $*$ -ring. Then the $*$ -ring

$$T_{3E}(R) = \left\{ \begin{pmatrix} a & b & c \\ 0 & a & d \\ 0 & 0 & a \end{pmatrix} \mid a, b, c, d \in R \right\},$$

is weakly $*$ -reversible by **Proposition 12**, for some $a \neq 0$. For $A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$, $B = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$, we have $AB = 0 = AB^*$ and $BA = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$, so $T_{3E}(R)$ is not $*$ -reversible, while it has quasi- $*$ -IFP.

We note that if R is a commutative then the $*$ -ring.

$$T_n(R) = \left\{ \begin{pmatrix} a & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a & a_{23} & \cdots & a_{2n} \\ 0 & 0 & a & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a \end{pmatrix} \mid a, a_{ij} \in R, n \geq 3 \right\},$$

is not $*$ -reversible by [3, Example 3.8] and is weakly $*$ -reversible by **Proposition 12**. Moreover, it is clear that $T_{4E}(R)$ is not quasi- $*$ -IFP and so $T_n(R)$ is not quasi- $*$ -IFP for $n \geq 4$.

The next example demonstrates that the condition $T_n(R)$ in **Proposition 12**, cannot be weakened to the full matrix $*$ -ring $\mathbb{M}_n(R)$, where n is any integer bigger than 1.

Example 7. Let R be a weakly $*$ -reversible $*$ -ring and n any integer bigger than 1, then $\mathbb{M}_2(R)$, with adjoint involution, is not weakly $*$ -reversible. For $A = \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, we have $AB = 0 = AB^*$ and for $C = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in \mathbb{M}_2(R)$, we see that $RBCA = \begin{pmatrix} 0 & a \\ 0 & c \end{pmatrix}$ is not nil.

The following result shows that the class of central $*$ -reversible $*$ -rings lies properly between the classes of $*$ -reversible and weakly $*$ -reversible $*$ -rings.

Theorem 1. Let R be a $*$ -ring and consider the following conditions.

1. R is $*$ -reversible.
2. R is central $*$ -reversible.

3. R is weakly $*$ -reversible.
Then $(1) \implies (2) \implies (3)$.

Proof.

$(1) \implies (2)$: Clearly.

$(2) \implies (3)$: Let $a, b \in R$ with $ab = ab^* = 0$. Then for all $s \in R$, $sab = sab^* = 0$ and bsa is central, since R is central $*$ -reversible. Hence $(rbsa)^2 = (rbsa)(rbsa) = r(bsa)r(bsa) = rr(bs(ab)sa) = 0$, for all $r, s \in R$ and R is weakly $*$ -reversible.

□

The converse of **Theorem 1** is not true by **Examples 1** and **6**. However, from **Corollary 3** and **Theorem 1** we get the following corollary.

Corollary 4. *Every $*$ -domain is a weakly $*$ -reversible $*$ -ring.*

4 Weakly quasi- $*$ -IFP

Here, weakly quasi- $*$ -IFP $*$ -rings are introduced as generalization for the class of quasi- $*$ -IFP $*$ -rings. First, we introduce weakly $*$ -IFP $*$ -rings.

Definition. A $*$ -ring R is called *weakly $*$ -IFP* if for all $a, b \in R$, $ab = 0$ implies $arb^* \in \text{nil}(R)$ for all $r \in R$.

Each commutative $*$ -ring is weakly $*$ -IFP. As before, one can easily prove the following result.

Proposition 13. *The class of weakly $*$ -IFP $*$ -ring is closed under direct sums (with changeless involution) and under taking $*$ -subrings.*

Proposition 14. *For a commutative $*$ -ring R , $T_{nE}(R)$ is weakly $*$ -IFP, with involution $*$ given in **Proposition 12**.*

Proof. Let $A = (a_{ij})$ and $B = (b_{ij}) \in T_{nE}(R)$ with $AB = 0$, where $1 \leq i \leq j \leq n$, then we have $ab = 0$, where a and b are the diagonal elements of A and B , respectively. Since R is weakly $*$ -IFP, there exists $k \in \mathbb{N}$ such that $(acb)^k = 0$ for all $C = (c_{ij}) \in T_{nE}(R)$, where c is the diagonal element of C . Hence $((ACB^*)^k)^n = 0$ and $T_{nE}(R)$ is weakly $*$ -IFP. □

It is clear that every $*$ -ring having $*$ -IFP is weakly $*$ -IFP while the converse is not always true as shown by the following example.

Example 8. The $*$ -ring $T_{3E}(\mathbb{Z})$ with the involution $*$ given by: $\begin{pmatrix} a & b & c \\ 0 & a & d \\ 0 & 0 & a \end{pmatrix}^* = \begin{pmatrix} a & d & c \\ 0 & a & b \\ 0 & 0 & a \end{pmatrix}$ is weakly $*$ -IFP by **Proposition 14**. For $A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$, we have $AB = 0$ and $ARB^* = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \neq 0$, so $T_{3E}(\mathbb{Z})$ has not $*$ -IFP.

By the way, there exists a weakly IFP $*$ -ring which is not weakly $*$ -IFP as in the next example.

Example 9. Let $\mathbb{F}$ be a field and consider the $*$ -ring $R = \mathbb{F} \oplus \mathbb{F}$, with the exchange involution $(a, b)^* = (b, a)$, for all $a, b \in \mathbb{F}$. R is clearly weakly IFP and is not weakly $*$ -IFP.

Next, we define weakly quasi- $*$ -IFP $*$ -rings

Definition. A $*$ -ring R is said to be *weakly quasi- $*$ -IFP* if for all $a, b \in R$, $ab = 0 = ab^*$ implies arb is a nilpotent element for each $r \in R$. Consequently arb^* is also nilpotent.

Each commutative $*$ -ring is weakly quasi $*$ -IFP. Clearly, each weakly IFP $*$ -ring is weakly quasi- $*$ -IFP. The converse is true when the ring has $*$ -IFP as shown in the following.

Proposition 15. *Let R be a $*$ -ring. If R is weakly quasi- $*$ -IFP and has $*$ -IFP, then R is weakly IFP.*

Proof. Clearly, since $ab = 0$, implies $aRb^* = 0$, by the $*$ -IFP property, and R is weakly quasi- $*$ -IFP. $\square$

Moreover, the class of weakly quasi- $*$ -IFP $*$ -ring is closed under direct sums (using changeless involution) and under taking $*$ -subrings.

Proposition 16. *The class of weakly quasi- $*$ -IFP $*$ -ring is closed under direct sums and under taking $*$ -subrings.*

By a proof similar to **Proposition 12**, we get the following.

Proposition 17. *If R is a commutative $*$ -ring, then $T_{nE}(R)$ is weakly quasi- $*$ -IFP, with involution $*$ given in **Proposition 12**.*

Note that if R is a commutative $*$ -ring then the $*$ -ring.

$$T_{nE}(R) = \left\{ \begin{pmatrix} a & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a & a_{23} & \cdots & a_{2n} \\ 0 & 0 & a & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a \end{pmatrix} \mid a, a_{ij} \in R, n \geq 3 \right\},$$

is not $*$ -reversible by [3, Example 3.8] and is weakly quasi- $*$ -IFP by **Proposition 17**. However, It is clearly that $T_{4E}(R)$ is not quasi- $*$ -IFP and so $T_{nE}(R)$ is not quasi- $*$ -IFP for $n \geq 4$.

The next example demonstrates that the condition $T_{nE}(R)$ in **Proposition 17**, cannot be weakened to the full matrix $*$ -ring $\mathbb{M}_n(R)$, where $n > 1$.

Example 10. $\mathbb{Z}$ is weakly quasi- $*$ -IFP $*$ -ring with identical involution, while the $*$ -ring $\mathbb{M}_2(\mathbb{Z})$ with adjoint involution $*$ is not weakly quasi- $*$ -IFP. Indeed, the matrices $A = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix}$ satisfy $AB = 0 = AB^*$ and for $C = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \in \mathbb{M}_2(R)$, we have $ACB = \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix}$ is not nilpotent.

It is well known that every $*$ -reversible $*$ -ring has quasi- $*$ -IFP by [3, Proposition 3.6]. Next, we prove that central $*$ -reversible $*$ -rings are weakly quasi- $*$ -IFP.

Theorem 2. *Let R be a $*$ -ring and consider the following conditions.*

1. R is $*$ -reversible.
 2. R is central $*$ -reversible.
 3. R is weakly quasi- $*$ -IFP.
- Then (1) $\implies$ (2) $\implies$ (3).*

Proof.

(1) $\implies$ (2). Is clear.

(2) $\implies$ (3). If $a, b \in R$ satisfy $ab = ab^* = 0$, then ba is central and $(arb)^2 = 0$. Hence arb is nilpotent for all $r \in R$ and R is weakly quasi- $*$ -IFP.

□

The converse of **Theorem 2** is not true by **Examples 1** and **6**. Moreover, from **Corollary 3** and **Theorem 2** we have the following result.

Corollary 5. *Every $*$ -domain is a weakly quasi- $*$ -IFP $*$ -ring.*

From **Proposition 4** we have immediately the following corollary.

Corollary 6. *If R is a $*$ -Baer and central $*$ -reversible $*$ -ring, then R has quasi $*$ -IFP.*

From [8, Proposition 2.20], if R is central reduced (that is every nilpotent element is central), then $T(R, R)$ is central reversible and from [3, Proposition 3.14], if R is $*$ -reduced and $*$ -reversible, then $T(R, R)$, with componentwise involution, is $*$ -reversible. Accordingly, we have the following corollaries.

Corollary 7. *If the $*$ -ring R is central reduced $*$ -ring then $T(R, R)$ is central $*$ -reversible.*

Corollary 8. *If the $*$ -ring R is reduced then $T(R, R)$ is central $*$ -reversible.*

Corollary 9. *If the $*$ -ring R is $*$ -reduced and $*$ -reversible then $T(R, R)$, with componentwise involution, is central $*$ -reversible.*

Corollary 10. *If the $*$ -ring R is reduced and $*$ -reversible then $T(R, R)$, with componentwise involution, is central $*$ -reversible.*

By [11, Corollary 2.4], R is weakly reversible if and only if its trivial extension $T(R, R)$ is weakly reversible and from **Proposition 12**, we have the following corollaries.

Corollary 11. *If R is weakly reversible then $T(R, R)$ is weakly $*$ -reversible.*

Corollary 12. *If $T(R, R)$ is weakly reversible then R is weakly $*$ -reversible.*

Corollary 13. *A commutative $*$ -ring R is weakly $*$ -reversible if and only if $T(R, R)$, with adjoint involution, is weakly $*$ -reversible.*

From [13, Corollary 2.1], R is weakly IFP if and only if $T(R, R)$ is weakly IFP and by **Proposition 17**, we have the following corollaries.

Corollary 14. *If R is weakly IFP then $T(R, R)$ is weakly quasi $*$ -IFP.*

Corollary 15. *If $T(R, R)$ is weakly IFP then R is weakly quasi $*$ -IFP.*

Corollary 16. *A commutative $*$ -ring R is weakly quasi $*$ -IFP if and only if $T(R, R)$, with adjoint involution, is weakly quasi $*$ -IFP.*

5 Extensions of $*$ -Reversible and Weakly quasi- $*$ -IFP $*$ -Rings

In this section, the properties of $*$ -reversible, central $*$ -reversible and weakly quasi- $*$ -IFP are shown to be extended from $*$ -ring to its localization, polynomial, Laurent polynomial, Dorroh extension and from Ore $*$ -ring to its classical

Quotient.

Let R be a $*$ -ring and S be a multiplicatively closed subset of R consisting of nonzero central regular elements, then the localization of R to S is $S^{-1}R = \{u^{-1}a | u \in S, a \in R\}$ is a $*$ -ring with involution $\diamond$ defined as:

$$(u^{-1}a)^\diamond = u^{-1*}a^* = u^{*-1}a^*.$$

Proposition 18. *A $*$ -ring R is $*$ -reversible if and only if $S^{-1}R$ is $*$ -reversible.*

Proof. Let R be a $*$ -reversible $*$ -ring and $\alpha\beta = 0 = \alpha\beta^\diamond$ with $\alpha = u^{-1}a$, $\beta = v^{-1}b$ where $a, b \in R$ and $u, v \in S$. Hence $\alpha\beta = u^{-1}av^{-1}b = u^{-1}v^{-1}ab = (vu)^{-1}ab = 0$ and $\alpha\beta^\diamond = u^{-1}a(v^*)^{-1}b^* = u^{-1}(v^*)^{-1}ab^* = (v^*u)^{-1}ab^* = 0$, since S is contained in the center of R , so $ab = 0 = ab^*$. By hypothesis $ba = 0$ which implies $\beta\alpha = v^{-1}bu^{-1}a = v^{-1}u^{-1}ba = (uv)^{-1}ba = 0$ and $S^{-1}R$ is $*$ -reversible. The converse is clear. $\square$

By a similar proof, we get analogous results for central $*$ -reversible and weakly quasi- $*$ -IFP $*$ -rings.

Proposition 19. *A $*$ -ring R is central $*$ -reversible if and only if $S^{-1}R$ is central $*$ -reversible.*

Proposition 20. *A $*$ -ring R is weakly quasi- $*$ -IFP, if and only if $S^{-1}R$ is weakly quasi- $*$ -IFP.*

From **Propositions 18, 19 and 20** we get the following corollaries.

Corollary 17. *If R is a reversible $*$ -ring, then $S^{-1}R$ is $*$ -reversible.*

Corollary 18. *If $S^{-1}R$ is a reversible $*$ -ring, then R is $*$ -reversible.*

Corollary 19. *If R is a central reversible $*$ -ring, then $S^{-1}R$ is central $*$ -reversible.*

Corollary 20. *If $S^{-1}R$ is a central reversible $*$ -ring, then R is central $*$ -reversible.*

Corollary 21. *If R has quasi- $*$ -IFP, then $S^{-1}R$ is weakly quasi- $*$ -IFP.*

Corollary 22. *If $S^{-1}R$ has quasi- $*$ -IFP, then R is weakly quasi- $*$ -IFP.*

The $*$ -ring of Laurent polynomials in x , with coefficients in a $*$ -ring R , consists of all formal sum $f(x) = \sum_{i=k}^n a_i x^i$ with obvious addition and multiplication, where $a_i \in R$ and k, n are (possibly negative) integers and with involution $*$ defined as $f^*(x) = \sum_{i=k}^n a_i^* x^i$. We denote this ring as usual by $R[x; x^{-1}]$.

Corollary 23. *Let R be a $*$ -ring. Then $R[x]$ is $*$ -reversible if and only if $R[x; x^{-1}]$ is $*$ -reversible.*

Proof. By [3, Proposition 3.15], it suffices to establish necessity. Clearly $S = \{1, x, x^2, \dots\}$ is a multiplicatively closed subset of $R[x]$. Since $R[x; x^{-1}] = S^{-1}R[x]$, it follows that $R[x; x^{-1}]$ is $*$ -reversible, by **Proposition 18**. $\square$

Corollary 24. *Let R be a $*$ -ring. Then $R[x]$ is central $*$ -reversible if and only if $R[x; x^{-1}]$ is central $*$ -reversible.*

Proof. By **Proposition 5**, it suffices to prove necessity which can be done as the proof of **Corollary 23** using **Proposition 19**. $\square$

Corollary 25. *For a $*$ -ring, $R[x]$ is weakly quasi- $*$ -IFP if and only if $R[x; x^{-1}]$ is weakly quasi- $*$ -IFP.*

Proof. By **Proposition 16**, it suffices to establish necessity which can be done as the proof of **Corollary 23** using **Proposition 20**. $\square$

From **Corollary 25** we have the following results.

Corollary 26. *If $R[x]$ has quasi- $*$ -IFP, then $R[x; x^{-1}]$ is weakly quasi- $*$ -IFP.*

Corollary 27. *If $R[x; x^{-1}]$ has quasi- $*$ -IFP, then $R[x]$ is weakly quasi- $*$ -IFP.*

Corollary 28. *If $R[x]$ has IFP, then $R[x; x^{-1}]$ is weakly quasi- $*$ -IFP.*

Corollary 29. *If $R[x; x^{-1}]$ has IFP, then $R[x]$ is weakly quasi- $*$ -IFP.*

Corollary 30. *If $R[x]$ has $*$ -IFP, then $R[x; x^{-1}]$ is weakly quasi- $*$ -IFP.*

Corollary 31. *If $R[x; x^{-1}]$ has $*$ -IFP, then $R[x]$ is weakly quasi- $*$ -IFP.*

A $*$ -ring R is called a $*$ -Armendariz $*$ -ring if whenever the polynomials $f(x) = \sum_{i=0}^m a_i x^i, g(x) = \sum_{j=0}^n b_j x^j \in R[x]$ satisfy $f(x)g(x) = f(x)g^*(x) = 0$, then $a_i b_j = 0$ for all i, j . Consequently $a_i b_j^* = 0$.

Theorem 3. *Let R be a $*$ -Armendariz $*$ -ring. Then the following statements are equivalent.*

1. R is $*$ -reversible (central $*$ -reversible).
2. $R[x]$ is $*$ -reversible (central $*$ -reversible).
3. $R[x; x^{-1}]$ is $*$ -reversible (central $*$ -reversible).

Proof.

(1) $\implies$ (2): Let $f(x) = \sum_{i=0}^m a_i x^i$ and $g(x) = \sum_{j=0}^n b_j x^j \in R[x]$ with $f(x)g(x) = 0 = f(x)g^*(x)$. Since R is $*$ -Armendariz, $a_i b_j = 0 = a_i b_j^*$ for each i and j . But R is $*$ -reversible (central $*$ -reversible), hence $b_j a_i = 0$ ($b_j a_i$ is central) for each i and j . It follows that $g(x)f(x) = 0$ ($g(x)f(x)$ is central) and $R[x]$ is $*$ -reversible (central $*$ -reversible).

(2) $\implies$ (1): Clear from [3, Proposition 3.15] (**Proposition 5**).

(2) $\iff$ (3): Follows from **Corollary 23** (**Corollary 24**).

□

The following corollary is an immediate from **Theorem 3**.

Corollary 32. *Let R be an Armendariz $*$ -ring. Then the following statements are equivalent.*

1. R is $*$ -reversible (central $*$ -reversible).
2. $R[x]$ is $*$ -reversible (central $*$ -reversible).
3. $R[x; x^{-1}]$ is $*$ -reversible (central $*$ -reversible).

The Dorroh extension $D(R, \mathbb{Z}) = \{(r, n) : r \in R, n \in \mathbb{Z}\}$ of a $*$ -ring R is a ring with componentwise addition and multiplication $(r_1, n_1)(r_2, n_2) = (r_1 r_2 + n_1 r_2 + n_2 r_1, n_1 n_2)$. The involution of R can be extended naturally to $D(R, \mathbb{Z})$ as $(r, n)^* = (r^*, n)$ (see [2]). We have the following:

Proposition 21. *A $*$ -ring R is $*$ -reversible if and only if its Dorroh extension $D(R, \mathbb{Z})$ of R is $*$ -reversible.*

Proof. The sufficiency is clear. For necessity, let $(r_1, n_1), (r_2, n_2) \in D(R, \mathbb{Z})$ with $(r_1, n_1)(r_2, n_2) = 0 = (r_1, n_1)(r_2^*, n_2)$, then from $0 = (r_1, n_1)(r_2, n_2) = (r_1 r_2 + n_1 r_2 + n_2 r_1, n_1 n_2)$ and $0 = (r_1, n_1)(r_2^*, n_2) = (r_1 r_2^* + n_1 r_2^* + n_2 r_1, n_1 n_2)$, we have $r_1 r_2 + n_1 r_2 + n_2 r_1 = 0$, $r_1 r_2^* + n_1 r_2^* + n_2 r_1 = 0$ and $n_1 n_2 = 0$. Since $\mathbb{Z}$ is $*$ -domain, $n_1 = 0$ or $n_2 = 0$. If $n_1 = 0$, we get $0 = r_1 r_2 + n_2 r_1 = r_1(r_2 + n_2)$ and $0 = r_1 r_2^* + n_2 r_1 = r_1(r_2^* + n_2)$. From the $*$ -reversibility of R it follows that $0 = (r_2 + n_2)r_1 = r_2 r_1 + n_2 r_1 = (r_2, n_2)(r_1, 0)$ and so $D(R, \mathbb{Z})$ is $*$ -reversible. □

By a similar proof to the previous proposition, we get the following.

Proposition 22. *A $*$ -ring R is central $*$ -reversible if and only if its Dorroh extension $D(R, \mathbb{Z})$ of R is central $*$ -reversible.*

Recall that a ring R is called right Ore if given $a, b \in R$ with b regular there exist $a_1, b_1 \in R$ with b_1 regular such that $ab_1 = ba_1$. Left Ore is defined

similarly and R is *Ore ring* if it is both right and left Ore. For $*$ -rings, right Ore implies left Ore and vice versa. It is a known fact that R is Ore if and only if its classical quotient ring Q of R exists and for $*$ -rings, $*$ can be extended to Q by $(a^{-1}b)^* = b^*(a^*)^{-1}$ (see[12, Lamme 4]).

Theorem 4. *Let R be an Ore $*$ -ring and Q be its classical quotient $*$ -ring, then R is $*$ -reversible if and only if Q is $*$ -reversible.*

Proof. The sufficiency is clear by [3, Proposition 3.15]. The proof of necessity is similar to that of [10, Theorem 2.6]. $\square$

From [10, Theorem 2.6] and **Theorem 4**, we have the following corollaries.

Corollary 33. *If R is a reversible $*$ -ring, then Q is $*$ -reversible.*

Corollary 34. *If Q is a reversible $*$ -ring, then R is $*$ -reversible.*

Corollary 35. *If R is a $*$ -reversible $*$ -ring, then Q is central $*$ -reversible (weakly $*$ -reversible).*

Corollary 36. *If Q is a $*$ -reversible $*$ -ring, then R is central $*$ -reversible (weakly $*$ -reversible).*

Conclusion

Finally, we can state following implications in the class of rings with involution.

$$\begin{array}{ccccc}
 & & \text{weakly IFP} & & \\
 & & \uparrow & & \\
 \text{reversible} & \implies & \text{central reversible} & \implies & \text{weakly reversible} \\
 \downarrow & & \downarrow & & \downarrow \\
 * - \text{reversible} & \implies & \text{central } * - \text{reversible} & \implies & \text{weakly } * - \text{reversible} \\
 \downarrow & & \downarrow & & \\
 \text{quasi } * - \text{IFP} & \implies & * - \text{Abelian} & & \\
 \downarrow & & & & \\
 \text{weakly quasi } * - \text{IFP} & & & &
 \end{array}$$

References

- [1] U. A. Aburawash. Semiprime involution rings and chain conditions. *Contr. to General Algebra* 7, Hölder-Pichler-Tempsky, Wien and B. G. Teubner, Stuttgart, :7–11, 1991.
- [2] U. A. Aburawash. On embedding of involution rings. *Mathematica Pannonica*, 8(2):245–250, 1997.
- [3] U. A. Aburawash and M. Saad. $*$ -reversible and $*$ -reflexive properties for rings with involution. *Submitted for publication*.
- [4] U. A. Aburawash and M. Saad. On biregular, IFP and quasi-Baer $*$ -rings. *East-West Journal of Mathematics*, 16(2):182–192, 2014.

- [5] U. A. Aburawash and M. Saad. $*$ -Baer property for rings with involution. *Studia Sci. Math. Hungar.*, 53(2):243–255, 2016.
- [6] U. A. Aburawash and Khadija B. Sola. $*$ -zero divisors and $*$ -prime ideals. *East-West J.Math.*, 12(1):27–31, 2010.
- [7] P. M. Cohn. Reversible rings. *Bull. London Math. Soc.*, 31:641–648, 1999.
- [8] H. Kose, B. Ungor, S. Halicioglu and A. Harmanci. A generalization of reversible rings. *Iranian Journal of science and Technology*, 38A(1):43–48, 2014.
- [9] I. Kaplansky. *Ring of operators*. Benjamin, New york, 1968.
- [10] N. K. Kim and Y. Lee. Extensions of reversible rings. *Journal of Pure and Applied Algebra*, 185:207–223, 2003.
- [11] Z. Liang and Y. Gang. On weakly reversible rings. *Acat Math. Unvi. Comenianae*, LXXVI(2):189–192, 2007.
- [12] W. S. Martindale. Rings with involution and polynomial identities. *Journal of Algebra*, 11:186–194, 1969.
- [13] L. Liang, L. Wang and Z. Liu. On a generalization of semicommutative rings. *Taiwanese Journal of Mathematics*, 11(5):1359–1368, 2007.