

ON FUNCTORIAL PROPERTIES OF BCK AND BCI-STRUCTURES

S.M.A. Zaidi, Shabbir Khan and Gulam Muhiuddin

*Department of Mathematics
Aligarh Muslim University
Aligarh - 202002, India.
zaidimath@math.com, skhanmu@rediffmail.com, gmchishty@math.com*

Abstract

In this paper, we study functors by retractions and co-retractions on BCI-algebras. Further, some functors by filters, prime ideals and self maps on BCK-algebras are discussed.

1. Introduction

The notion of BCK-algebra was originated by Imai and Iseki [6]. Later on, Iseki [7] introduced the notion of a BCI-algebra as a generalization of a BCK-algebra. A series of interesting notions concerning BCI-algebras were introduced. Several papers have been written on various aspects of these algebras and their relations to other algebraic structures have been studied (see [9,10,11,12]).

Iseki and Thaheem [14] proved that if X is an associative BCI-algebra, then $End(X)$; the set of all BCI-homomorphisms on X , is also a BCI-algebra. They also proved that for any $\theta_1, \theta_2 \in End(X)$, $\theta_1 \star \theta_2 \in End(X)$ together with the mapping $\theta_1 \star \theta_2 : X \rightarrow X$ defined by $(\theta_1 \star \theta_2)(x) = \theta_1(x) \star \theta_2(x)$ for all $x \in X$.

In developing an algebraic theory of BCK-algebras, the notion of ideals has played an important role. The theory of ideals has been developed by Iseki [8] and extensively studied by several authors. Deeba [4] introduced the notion of filters as a dual to the concept of ideals of BCK-algebras and proved that if P_1, P_2 and P_3 are prime ideals of a bounded commutative BCK-algebra X , then

Key words and phrases: BCK-homomorphism, bounded commutative BCK-algebra and $BCK(i_+)$; category of positive implicative BCK-algebras
2000 AMS Mathematics Subject Classification: 18A20, 18D05, 06F35

$X-P_1$, $X-P_2$ and $X-P_3$ will be the corresponding filters of X . Thus there is a one to one correspondence between the ideals and filters of a bounded commutative BCK-algebra. Moreover, if P is a prime ideal of X and $x \in P$, then $Nx \in (X-P)$, the corresponding filter of the ideal P .

Kondo [15] defined a left map l_x on a positive implicative BCK-algebra X and proved that if X is a positive implicative BCK-algebra, then the set of all left maps $L(X)$ is also a positive implicative BCK-algebra.

Motivated from [4, 14, 15], we study functorial properties of BCK and BCI-structures in view of the fact that a structure has a functorial property if a functor can be defined by it. In particular, we construct some functors using the functorial properties of BCK and BCI-structures such as retractions, co-retractions, filters, prime ideals and self maps.

2. Preliminaries

In this section, we recall the following aspects of the theory of BCK and BCI-algebras that are necessary for the development of this paper from [9, 13, 14, 16]. For background information on categories and functors we refer the reader to [1].

Let X be a set with binary operation ' \star ' and a constant 0, then X is called BCI-algebra if the following axioms are satisfied for all $x, y, z \in X$:

$$(i) \quad (x \star y) \star (x \star z) \leq z \star y,$$

$$(ii) \quad x \star (x \star y) \leq y,$$

$$(iii) \quad x \leq x,$$

$$(iv) \quad x \leq 0 \implies x = 0,$$

$$(v) \quad x \leq y \text{ and } y \leq x \implies x = y,$$

$$(vi) \quad x \leq y \iff x \star y = 0.$$

If we replace axiom (iv) by $0 \leq x$, X is called BCK-algebra.

A BCK-algebra X is said to be commutative if $x \wedge y = y \wedge x$, where $x \wedge y = y \star (y \star x)$.

A BCK-algebra X is said to be bounded if there is an element 1 in X such that $x \leq 1$ for all $x \in X$. In a bounded BCK-algebra, we denote $1 \star x$ by N_x .

A BCK-algebra X is said to be positive implicative if $(x \star z) \star (y \star z) = (x \star y) \star z$ for all $x, y, z \in X$.

Let X be a positive implicative BCK-algebra. A self map $l_x : X \rightarrow X$ defined by $l_x(t) = x \star t$ for all $t \in X$, is called a left map of X . The composition of left maps is defined by $l_x o l_y = l_{x \star y}$ for all $x, y \in X$.

A non-empty subset I of BCK-algebra X is said to be an ideal of X if

- (i) $0 \in I$,
- (ii) $x \in I$ and $y \star x \in I$ imply that $y \in I$.

An ideal I of a commutative BCK-algebra X is said to be prime if $x \wedge y \in I$ implies $x \in I$ or $y \in I$.

A non-empty set F of BCK-algebra X is said to be a filter of X if

- (i) $x \in F$ and $y \geq x$ imply that $y \in F$
- (ii) $x \in F$ and $y \in F$ imply that $glb\{x, y\} \in F$.

Let X and Y be BCI-algebras (BCK-algebras). Then a mapping $f : X \rightarrow Y$ is called BCI-homomorphism (BCK-homomorphism) if

$$f(x \star y) = f(x) \star f(y) \quad \text{for all } x, y \in X.$$

The category of BCI-algebras can be constructed by taking the class of all BCI-algebras as the class of objects of the category and the class of all BCI-homomorphisms as the class of morphisms of the category. We shall denote the category of BCI-algebras by BCI .

Similarly, the category of BCK-algebras is constructed and denote it by BCK .

3. Functors by Retractions and Co-retractions

Consider the category $B_{(r)}$ in which the objects of the category are associative BCI-algebras and morphisms of the category are those BCI-homomorphisms which are retractions.

Theorem 3.1. Let $f : X \rightarrow Y$ be a morphism in the category $B_{(r)}$ and g be any right inverse of f . Then the mapping $T_f : End(X) \rightarrow End(Y)$ defined by $T_f(\theta) = f o \theta o g$ for all $\theta \in End(X)$, is a BCI-homomorphism.

Proof. If $\theta_1, \theta_2 : X \rightarrow X$, then we get $T_f(\theta_1 \star \theta_2) = fo(\theta_1 \star \theta_2)og$ and $T_f(\theta_1) \star T_f(\theta_2) = (fo\theta_1og) \star (fo\theta_2og)$. Clearly, $T_f(\theta_1 \star \theta_2) : Y \rightarrow Y$ and for any $y \in Y$, one gets

$$\begin{aligned} T_f(\theta_1 \star \theta_2)(y) &= (fo(\theta_1 \star \theta_2)og)(y) \\ &= (fo(\theta_1 \star \theta_2)o(g(y))) \\ &= f[(\theta_1 \star \theta_2)(g(y))] \quad (\text{as } g(y) \in X) \\ &= (fo\theta_1og)(y) \star (fo\theta_2og)(y) \\ &= T_f(\theta_1)(y) \star T_f(\theta_2)(y) \end{aligned}$$

which yields $T_f(\theta_1 \star \theta_2) = T_f(\theta_1) \star T_f(\theta_2)$ and henceforth T_f is a BCI-homomorphism.

Proposition 3.1. If $g : X \rightarrow Y$ and $h : Y \rightarrow Z$ are retractions, then $hog : X \rightarrow Z$ is also a retraction.

Proof Since $g : X \rightarrow Y$ and $h : Y \rightarrow Z$ are retractions, then there exist $g' : Y \rightarrow X$ and $h' : Z \rightarrow Y$ such that $gog' = I_Y$ and $hoh' = I_Z$. Now, $(hog)o(g'oh') = ho(gog')oh' = ho(I_Y)oh' = (hoI_Y)oh' = hoh' = I_Z$ which implies $(hog)o(g'oh') = I_Z$ and so $hog : X \rightarrow Z$ is a retraction.

Proposition 3.2. If $g : X \rightarrow Y$ and $h : Y \rightarrow Z$ are morphisms in $B_{(r)}$, then $T_{hog} = T_h o T_g$.

Proof In view of Theorem 3.1 and Proposition 3.1, $T_{hog} : End(X) \rightarrow End(Z)$ is a BCI-homomorphism. For any $\theta \in Hom(X)$, we get

$$\begin{aligned} T_{hog}(\theta) &= (hog)o\theta o(g'oh') \\ &= h(go\theta)o(g'oh') \\ &= ho(go\thetaog')oh' \\ &= ho(T_g(\theta))oh' \\ &= T_h(T_g(\theta)) \\ &= (T_h o T_g)(\theta) \end{aligned}$$

implying thereby $T_{hog} = T_h o T_g$.

Corollary 3.1. If $I_X : X \rightarrow X$ is an identity morphism in $B_{(r)}$, then $T_{I_X} : End(X) \rightarrow End(X)$ is an identity morphism in B_{CI} .

In view of Theorem 3.1, Propositions 3.1, 3.2 and Corollary 3.1, we can define a functor $T : B_{(r)} \rightarrow B_{CI}$ such that

- (i) $T(X) = End(X)$ for all $X \in B_{(r)}$,
- (ii) for any morphism $f : X \rightarrow Y$ in $B_{(r)}$, we have $T(f) = T_f : End(X) \rightarrow End(Y)$ in B_{CI} .

Now, consider the category $B_{(cr)}$ in which the objects of the category are associative BCI-algebras and morphisms of the category are those BCI-homomorphisms which are co-retractions.

Theorem 3.2. Let $f : X \rightarrow Y$ be a morphism in the category $B_{(cr)}$ and g be any left inverse of f . Then the mapping $G_f : End(Y) \rightarrow End(X)$ defined by $G_f(\beta) = go\beta of$ for all $\beta \in End(Y)$, is a BCI-homomorphism.

Proof If $\beta_1, \beta_2 \in End(Y)$, then $\beta_1 \star \beta_2 \in End(Y)$. For any $x \in X$, we get

$$\begin{aligned}
 G_f(\beta_1 \star \beta_2)(x) &= (go(\beta_1 \star \beta_2)of)(x) \\
 &= [go(\beta_1 \star \beta_2)]f(x) \\
 &= g[\beta_1(f(x)) \star \beta_2(f(x))] \\
 &= g(\beta_1(f(x))) \star g(\beta_2(f(x))) \\
 &= (go\beta_1 of)(x) \star (go\beta_2 of)(x) \\
 &= G_f(\beta_1)(x) \star G_f(\beta_2)(x) \\
 &= (G_f(\beta_1) \star G_f(\beta_2))(x)
 \end{aligned}$$

which yields $G_f(\beta_1 \star \beta_2) = G_f(\beta_1) \star G_f(\beta_2)$. Therefore, G_f is a BCI-homomorphism.

Proposition 3.3. If $g : X \rightarrow Y$ and $h : Y \rightarrow Z$ are co-retractions, then $hog : X \rightarrow Z$ is a co-retraction.

Proof The proof follows on the similar lines of Proposition 3.1.

Proposition 3.4. If $g : X \rightarrow Y$ and $h : Y \rightarrow Z$ are morphisms in $B_{(cr)}$, then $G_{hog} = G_g \circ G_h$.

Proof In view of Theorem 3.2 and Proposition 3.3, $G_{hog} : End(Z) \rightarrow End(X)$ is a BCI-homomorphism and for any $\mu \in End(Z)$, one gets

$$\begin{aligned}
 G_{hog}(\mu) &= (g'oh')o\mu o(hog) && \text{(as } g, h \text{ are co-retractions)} \\
 &= (g'oh')o(\mu oh)og \\
 &= g'o(h'o\mu oh)og \\
 &= g'o(G_h(\mu))og \\
 &= G_g(G_h(\mu)) \\
 &= (G_g o G_h)(\mu)
 \end{aligned}$$

which implies $G_{hog} = G_g o G_h$.

Corollary 3.2. If $I_X : X \rightarrow X$ is an identity morphism in $B_{(cr)}$, then $G_{I_X} : End(X) \rightarrow End(X)$ is an identity morphism in B_{CI} .

In view of Theorem 3.2, Propositions 3.3, 3.4 and Corollary 3.2, we can define a contravariant functor $G : B_{(cr)} \rightarrow B_{CI}$ such that

- (i) $G(X) = End(X)$ for all $X \in B_{(cr)}$,
- (ii) for any morphism $f : X \rightarrow Y$ in $B_{(cr)}$, we have $G(f) = G_f : End(Y) \rightarrow End(X)$ in B_{CI} .

4. Functors by Filters and Prime Ideals

Let $P(X)$ represents the category of prime ideals of X in which objects of the category are the prime ideals and morphisms of the category are the BCK-homomorphisms between them. Further, let $F(X)$ denotes the category of filters of X in which objects of the category are the filters and the morphisms of the category are the functions between them.

Let P_1, P_2 be the prime ideals. Then for any morphism $f : P_1 \rightarrow P_2$ in $P(X)$, we can define a function $S_f : (X - P_1) \rightarrow (X - P_2)$ in $F(X)$ by $S_f(N_x) = N_{f(x)}$ for all $x \in X$.

Proposition 4.1. If $f : P_1 \rightarrow P_2$ and $g : P_2 \rightarrow P_3$ are BCK-homomorphisms where P_1, P_2 and P_3 are prime ideals of the bounded commutative BCK-algebra X , then $S_{g \circ f} = S_g \circ S_f$.

Proof Let X be a bounded commutative BCK-algebra. Let $f : P_1 \rightarrow P_2$ and $g : P_2 \rightarrow P_3$ be the morphisms in $P(X)$. Then for the composite morphism $gof : P_1 \rightarrow P_3$ in $P(X)$, the mapping $S_{gof} : (X-P_1) \rightarrow (X-P_3)$ is a morphism in $F(X)$. For any $N_x \in (X-P_1)$, we get

$$\begin{aligned} S_{gof}(N_x) &= N_{(gof)(x)} \\ &= N_{g(f(x))} \\ &= S_g(N_{f(x)}) \\ &= S_g(S_f(N_x)) \\ &= (S_g \circ S_f)(N_x) \end{aligned}$$

yielding thereby $S_{gof} = S_g \circ S_f$.

Corollary 4.1. If $I_P : P \rightarrow P$ is an identity homomorphism in $P(X)$, then $S_{I_P} : (X-P) \rightarrow (X-P)$ is an identity morphism in $F(X)$.

In view of the above discussion, we can define a covariant functor $S : P(X) \rightarrow F(X)$ such that

- (i) $S(P) = (X-P)$, for all $P \in P(X)$,
- (ii) for any morphism $f : P_1 \rightarrow P_2$ in $P(X)$, we have $S(f) = S_f : (X-P_1) \rightarrow (X-P_2)$ in $F(X)$.

5. Functors by Self Maps

Consider the category $BCK(i_+)$ in which the objects of the category are positive implicative BCK-algebras and morphisms of the category are the BCI-homomorphisms between them.

Proposition 5.1. If $f : X \rightarrow Y$ is a BCK-homomorphism, then the map $L_f : L(X) \rightarrow L(Y)$ defined by $L_f(l_x) = l_{f(x)}$ for all $x \in X$, is a BCK-homomorphism.

Proof For any $l_{x_1}, l_{x_2} \in L(X)$, we have

$$\begin{aligned}
L_f(l_{x_1} o l_{x_2}) &= L_f(l_{x_1 \star x_2}) \\
&= l_{f(x_1 \star x_2)} \\
&= l_{f(x_1) \star f(x_2)} \\
&= l_{f(x_1)} o l_{f(x_2)} \text{ [By the definition of composition} \\
&\quad \text{of maps in } L(X)] \\
&= L_f(l_{x_1}) o L_f(l_{x_2})
\end{aligned}$$

which shows that $L_f : L(X) \rightarrow L(Y)$ is a BCK-homomorphism.

Theorem 5.1. If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are BCK-homomorphisms in $B_{CK}(i_+)$, then $L_{gof} = L_g o L_f$.

Proof In view of Proposition 5.1, the maps $L_f : L(X) \rightarrow L(Y)$, $L_g : L(Y) \rightarrow L(Z)$ and their composition $L_{gof} : L(X) \rightarrow L(Z)$ are morphisms in the category $B_{CK}(i_+)$.

For any $l_x \in L(X)$, we have

$$\begin{aligned}
L_{gof}(l_x) &= l_{gof(x)} \\
&= l_{g(f(x))} \\
&= L_g(l_{f(x)}) \\
&= L_g(L_f(l_x)) \\
&= (L_g o L_f)(l_x)
\end{aligned}$$

yielding thereby $L_{gof} = L_g o L_f$.

Corollary 5.1. If $I_X : X \rightarrow X$ is an identity homomorphism in $B_{CK}(i_+)$, then $L_{I_X} : L(X) \rightarrow L(X)$ is also an identity homomorphism in $B_{CK}(i_+)$.

With the help of Proposition 5.1, Theorem 5.1 and Corollary 5.1, we can define a functor $F : B_{CK}(i_+) \rightarrow B_{CK}(i_+)$ such that

- (i) $F(X) = L(X)$ for all $X \in B_{CK}(i_+)$,
- (ii) for any morphism $f : X \rightarrow Y$ in $B_{CK}(i_+)$, we have $F(f) = L_f : L(X) \rightarrow L(Y)$ in $B_{CK}(i_+)$.

Acknowledgement: The authors are greatly indebted to the referee for several helpful suggestions and healthy comments.

References

- [1] T.S. Blyth, *Categories*, Longman, New York 1986.
- [2] M.A. Chaudhry and S.A. Bhatti, *On the category of BCI-algebras*, Punjab Univ. J. Math. 21 (1988), 34-39.
- [3] Z.M. Chen and H.X. Wang, *Some universal properties of BCI-algebras*, Kobe J. Math. 6 (1989), 43-48.
- [4] E.Y. Deeba, *Filter theory of BCK-algebras*, Math. Japon. 25 (1980), 631-639.
- [5] C.S. Hoo, *Injectives in the category of BCI-algebras*, Math. Japon. 33 (1988), 237-246.
- [6] Y. Imai and K. Iseki, *On axiom systems of propositional calculi XIV*, Proc. Japan Acad. 42 (1966), 19-22.
- [7] K. Iseki, *An algebra related with a propositional calculus*, Proc. Japan Acad. 42 (1966), 26-29.
- [8] K. Iseki, *On ideals in BCI-algebras*, Math. Sem. Notes, Kobe Univ. 3 (1975), 1-12.
- [9] K. Iseki, *On BCI-algebras*, Math. Sem. Notes, Kobe Univ. 8 (1980), 125-130.
- [10] K. Iseki, *On the existence of quasi-commutative BCI-algebras*, Math. Sem. Notes, Kobe Univ. 8 (1980), 181-186.
- [11] K. Iseki, *On variety of BCI-algebras*, Math. Sem. Notes, Kobe Univ. 8 (1980), 225-226.
- [12] K. Iseki, *A problem on BCI-algebras*, Math. Sem. Notes, Kobe Univ. 8 (1980), 235-236.
- [13] K. Iseki and S. Tanaka, *An introduction to the theory of BCK-algebras*, Math. Japon. 23 (1978), 1-26.
- [14] K. Iseki and A.B. Thaheem, *Note on BCI-algebras*, Math. Japon. 29 (1984), 255-258.
- [15] M. Kondo, *Positive implicative BCK-algebra and its dual algebra*, Math. Japon. 35 (1990), 289-291.
- [16] S.M.A. Zaidi and S. Khan, *A note on the category of BCK-algebras with regular morphisms*, Proc. Internat. Conf. Algebra and its Applications, A.M.U. Aligarh (2001), 305-315.