

## ON FUNCTORIAL PROPERTIES OF BCK AND BCI-STRUCTURES

S.M.A. Zaidi, Shabbir Khan and Gulam Muhiuddin

*Department of Mathematics  
Aligarh Muslim University  
Aligarh - 202002, India.  
zaidimath@math.com, skhanmu@rediffmail.com, gmchishty@math.com*

### Abstract

In this paper, we study functors by retractions and co-retractions on BCI-algebras. Further, some functors by filters, prime ideals and self maps on BCK-algebras are discussed.

### 1. Introduction

The notion of BCK-algebra was originated by Imai and Iseki [6]. Later on, Iseki [7] introduced the notion of a BCI-algebra as a generalization of a BCK-algebra. A series of interesting notions concerning BCI-algebras were introduced. Several papers have been written on various aspects of these algebras and their relations to other algebraic structures have been studied (see [9,10,11,12]).

Iseki and Thaheem [14] proved that if  $X$  is an associative BCI-algebra, then  $End(X)$ ; the set of all BCI-homomorphisms on  $X$ , is also a BCI-algebra. They also proved that for any  $\theta_1, \theta_2 \in End(X)$ ,  $\theta_1 \star \theta_2 \in End(X)$  together with the mapping  $\theta_1 \star \theta_2 : X \rightarrow X$  defined by  $(\theta_1 \star \theta_2)(x) = \theta_1(x) \star \theta_2(x)$  for all  $x \in X$ .

In developing an algebraic theory of BCK-algebras, the notion of ideals has played an important role. The theory of ideals has been developed by Iseki [8] and extensively studied by several authors. Deeba [4] introduced the notion of filters as a dual to the concept of ideals of BCK-algebras and proved that if  $P_1, P_2$  and  $P_3$  are prime ideals of a bounded commutative BCK-algebra  $X$ , then

---

**Key words and phrases:** BCK-homomorphism, bounded commutative BCK-algebra and  $BCK(i_+)$ ; category of positive implicative BCK-algebras  
2000 AMS Mathematics Subject Classification: 18A20, 18D05, 06F35

$X-P_1$ ,  $X-P_2$  and  $X-P_3$  will be the corresponding filters of  $X$ . Thus there is a one to one correspondence between the ideals and filters of a bounded commutative BCK-algebra. Moreover, if  $P$  is a prime ideal of  $X$  and  $x \in P$ , then  $Nx \in (X-P)$ , the corresponding filter of the ideal  $P$ .

Kondo [15] defined a left map  $l_x$  on a positive implicative BCK-algebra  $X$  and proved that if  $X$  is a positive implicative BCK-algebra, then the set of all left maps  $L(X)$  is also a positive implicative BCK-algebra.

Motivated from [4, 14, 15], we study functorial properties of BCK and BCI-structures in view of the fact that a structure has a functorial property if a functor can be defined by it. In particular, we construct some functors using the functorial properties of BCK and BCI-structures such as retractions, co-retractions, filters, prime ideals and self maps.

## 2. Preliminaries

In this section, we recall the following aspects of the theory of BCK and BCI-algebras that are necessary for the development of this paper from [9, 13, 14, 16]. For background information on categories and functors we refer the reader to [1].

Let  $X$  be a set with binary operation ' $\star$ ' and a constant 0, then  $X$  is called BCI-algebra if the following axioms are satisfied for all  $x, y, z \in X$ :

$$(i) \quad (x \star y) \star (x \star z) \leq z \star y,$$

$$(ii) \quad x \star (x \star y) \leq y,$$

$$(iii) \quad x \leq x,$$

$$(iv) \quad x \leq 0 \implies x = 0,$$

$$(v) \quad x \leq y \text{ and } y \leq x \implies x = y,$$

$$(vi) \quad x \leq y \iff x \star y = 0.$$

If we replace axiom (iv) by  $0 \leq x$ ,  $X$  is called BCK-algebra.

A BCK-algebra  $X$  is said to be commutative if  $x \wedge y = y \wedge x$ , where  $x \wedge y = y \star (y \star x)$ .

A BCK-algebra  $X$  is said to be bounded if there is an element 1 in  $X$  such that  $x \leq 1$  for all  $x \in X$ . In a bounded BCK-algebra, we denote  $1 \star x$  by  $N_x$ .

A BCK-algebra  $X$  is said to be positive implicative if  $(x \star z) \star (y \star z) = (x \star y) \star z$  for all  $x, y, z \in X$ .

Let  $X$  be a positive implicative BCK-algebra. A self map  $l_x : X \rightarrow X$  defined by  $l_x(t) = x \star t$  for all  $t \in X$ , is called a left map of  $X$ . The composition of left maps is defined by  $l_x o l_y = l_{x \star y}$  for all  $x, y \in X$ .

A non-empty subset  $I$  of BCK-algebra  $X$  is said to be an ideal of  $X$  if

- (i)  $0 \in I$ ,
- (ii)  $x \in I$  and  $y \star x \in I$  imply that  $y \in I$ .

An ideal  $I$  of a commutative BCK-algebra  $X$  is said to be prime if  $x \wedge y \in I$  implies  $x \in I$  or  $y \in I$ .

A non-empty set  $F$  of BCK-algebra  $X$  is said to be a filter of  $X$  if

- (i)  $x \in F$  and  $y \geq x$  imply that  $y \in F$
- (ii)  $x \in F$  and  $y \in F$  imply that  $glb\{x, y\} \in F$ .

Let  $X$  and  $Y$  be BCI-algebras (BCK-algebras). Then a mapping  $f : X \rightarrow Y$  is called BCI-homomorphism (BCK-homomorphism) if

$$f(x \star y) = f(x) \star f(y) \quad \text{for all } x, y \in X.$$

The category of BCI-algebras can be constructed by taking the class of all BCI-algebras as the class of objects of the category and the class of all BCI-homomorphisms as the class of morphisms of the category. We shall denote the category of BCI-algebras by  $BCI$ .

Similarly, the category of BCK-algebras is constructed and denote it by  $BCK$ .

### 3. Functors by Retractions and Co-retractions

Consider the category  $B_{(r)}$  in which the objects of the category are associative BCI-algebras and morphisms of the category are those BCI-homomorphisms which are retractions.

**Theorem 3.1.** Let  $f : X \rightarrow Y$  be a morphism in the category  $B_{(r)}$  and  $g$  be any right inverse of  $f$ . Then the mapping  $T_f : End(X) \rightarrow End(Y)$  defined by  $T_f(\theta) = f o \theta o g$  for all  $\theta \in End(X)$ , is a BCI-homomorphism.

**Proof.** If  $\theta_1, \theta_2 : X \rightarrow X$ , then we get  $T_f(\theta_1 \star \theta_2) = fo(\theta_1 \star \theta_2)og$  and  $T_f(\theta_1) \star T_f(\theta_2) = (fo\theta_1og) \star (fo\theta_2og)$ . Clearly,  $T_f(\theta_1 \star \theta_2) : Y \rightarrow Y$  and for any  $y \in Y$ , one gets

$$\begin{aligned} T_f(\theta_1 \star \theta_2)(y) &= (fo(\theta_1 \star \theta_2)og)(y) \\ &= (fo(\theta_1 \star \theta_2)o)(g(y)) \\ &= f[(\theta_1 \star \theta_2)(g(y))] \quad (\text{as } g(y) \in X) \\ &= (fo\theta_1og)(y) \star (fo\theta_2og)(y) \\ &= T_f(\theta_1)(y) \star T_f(\theta_2)(y) \end{aligned}$$

which yields  $T_f(\theta_1 \star \theta_2) = T_f(\theta_1) \star T_f(\theta_2)$  and henceforth  $T_f$  is a BCI-homomorphism.

**Proposition 3.1.** If  $g : X \rightarrow Y$  and  $h : Y \rightarrow Z$  are retractions, then  $hog : X \rightarrow Z$  is also a retraction.

**Proof** Since  $g : X \rightarrow Y$  and  $h : Y \rightarrow Z$  are retractions, then there exist  $g' : Y \rightarrow X$  and  $h' : Z \rightarrow Y$  such that  $gog' = I_Y$  and  $hoh' = I_Z$ . Now,  $(hog)o(g'oh') = ho(gog')oh' = ho(I_Y)oh' = (hoI_Y)oh' = hoh' = I_Z$  which implies  $(hog)o(g'oh') = I_Z$  and so  $hog : X \rightarrow Z$  is a retraction.

**Proposition 3.2.** If  $g : X \rightarrow Y$  and  $h : Y \rightarrow Z$  are morphisms in  $B_{(r)}$ , then  $T_{hog} = T_h o T_g$ .

**Proof** In view of Theorem 3.1 and Proposition 3.1,  $T_{hog} : End(X) \rightarrow End(Z)$  is a BCI-homomorphism. For any  $\theta \in Hom(X)$ , we get

$$\begin{aligned} T_{hog}(\theta) &= (hog)o\theta o(g'oh') \\ &= h(go\theta)o(g'oh') \\ &= ho(go\theta o g')oh' \\ &= ho(T_g(\theta))oh' \\ &= T_h(T_g(\theta)) \\ &= (T_h o T_g)(\theta) \end{aligned}$$

implying thereby  $T_{hog} = T_h o T_g$ .

**Corollary 3.1.** If  $I_X : X \rightarrow X$  is an identity morphism in  $B_{(r)}$ , then  $T_{I_X} : End(X) \rightarrow End(X)$  is an identity morphism in  $B_{CI}$ .

In view of Theorem 3.1, Propositions 3.1, 3.2 and Corollary 3.1, we can define a functor  $T : B_{(r)} \rightarrow B_{CI}$  such that

- (i)  $T(X) = End(X)$  for all  $X \in B_{(r)}$ ,
- (ii) for any morphism  $f : X \rightarrow Y$  in  $B_{(r)}$ , we have  $T(f) = T_f : End(X) \rightarrow End(Y)$  in  $B_{CI}$ .

Now, consider the category  $B_{(cr)}$  in which the objects of the category are associative BCI-algebras and morphisms of the category are those BCI-homomorphisms which are co-retractions.

**Theorem 3.2.** Let  $f : X \rightarrow Y$  be a morphism in the category  $B_{(cr)}$  and  $g$  be any left inverse of  $f$ . Then the mapping  $G_f : End(Y) \rightarrow End(X)$  defined by  $G_f(\beta) = go\beta of$  for all  $\beta \in End(Y)$ , is a BCI-homomorphism.

**Proof** If  $\beta_1, \beta_2 \in End(Y)$ , then  $\beta_1 \star \beta_2 \in End(Y)$ . For any  $x \in X$ , we get

$$\begin{aligned}
 G_f(\beta_1 \star \beta_2)(x) &= (go(\beta_1 \star \beta_2)of)(x) \\
 &= [go(\beta_1 \star \beta_2)]f(x) \\
 &= g[\beta_1(f(x)) \star \beta_2(f(x))] \\
 &= g(\beta_1(f(x))) \star g(\beta_2(f(x))) \\
 &= (go\beta_1 of)(x) \star (go\beta_2 of)(x) \\
 &= G_f(\beta_1)(x) \star G_f(\beta_2)(x) \\
 &= (G_f(\beta_1) \star G_f(\beta_2))(x)
 \end{aligned}$$

which yields  $G_f(\beta_1 \star \beta_2) = G_f(\beta_1) \star G_f(\beta_2)$ . Therefore,  $G_f$  is a BCI-homomorphism.

**Proposition 3.3.** If  $g : X \rightarrow Y$  and  $h : Y \rightarrow Z$  are co-retractions, then  $hog : X \rightarrow Z$  is a co-retraction.

**Proof** The proof follows on the similar lines of Proposition 3.1.

**Proposition 3.4.** If  $g : X \rightarrow Y$  and  $h : Y \rightarrow Z$  are morphisms in  $B_{(cr)}$ , then  $G_{hog} = G_g \circ G_h$ .

**Proof** In view of Theorem 3.2 and Proposition 3.3,  $G_{hog} : End(Z) \rightarrow End(X)$  is a BCI-homomorphism and for any  $\mu \in End(Z)$ , one gets

$$\begin{aligned}
 G_{hog}(\mu) &= (g'oh')o\mu o(hog) && \text{(as } g, h \text{ are co-retractions)} \\
 &= (g'oh')o(\mu oh)og \\
 &= g'o(h'o\mu oh)og \\
 &= g'o(G_h(\mu))og \\
 &= G_g(G_h(\mu)) \\
 &= (G_g o G_h)(\mu)
 \end{aligned}$$

which implies  $G_{hog} = G_g o G_h$ .

**Corollary 3.2.** If  $I_X : X \rightarrow X$  is an identity morphism in  $B_{(cr)}$ , then  $G_{I_X} : End(X) \rightarrow End(X)$  is an identity morphism in  $B_{CI}$ .

In view of Theorem 3.2, Propositions 3.3, 3.4 and Corollary 3.2, we can define a contravariant functor  $G : B_{(cr)} \rightarrow B_{CI}$  such that

- (i)  $G(X) = End(X)$  for all  $X \in B_{(cr)}$ ,
- (ii) for any morphism  $f : X \rightarrow Y$  in  $B_{(cr)}$ , we have  $G(f) = G_f : End(Y) \rightarrow End(X)$  in  $B_{CI}$ .

#### 4. Functors by Filters and Prime Ideals

Let  $P(X)$  represents the category of prime ideals of  $X$  in which objects of the category are the prime ideals and morphisms of the category are the BCK-homomorphisms between them. Further, let  $F(X)$  denotes the category of filters of  $X$  in which objects of the category are the filters and the morphisms of the category are the functions between them.

Let  $P_1, P_2$  be the prime ideals. Then for any morphism  $f : P_1 \rightarrow P_2$  in  $P(X)$ , we can define a function  $S_f : (X - P_1) \rightarrow (X - P_2)$  in  $F(X)$  by  $S_f(N_x) = N_{f(x)}$  for all  $x \in X$ .

**Proposition 4.1.** If  $f : P_1 \rightarrow P_2$  and  $g : P_2 \rightarrow P_3$  are BCK-homomorphisms where  $P_1, P_2$  and  $P_3$  are prime ideals of the bounded commutative BCK-algebra  $X$ , then  $S_{g \circ f} = S_g \circ S_f$ .

**Proof** Let  $X$  be a bounded commutative BCK-algebra. Let  $f : P_1 \rightarrow P_2$  and  $g : P_2 \rightarrow P_3$  be the morphisms in  $P(X)$ . Then for the composite morphism  $gof : P_1 \rightarrow P_3$  in  $P(X)$ , the mapping  $S_{gof} : (X-P_1) \rightarrow (X-P_3)$  is a morphism in  $F(X)$ . For any  $N_x \in (X-P_1)$ , we get

$$\begin{aligned} S_{gof}(N_x) &= N_{(gof)(x)} \\ &= N_{g(f(x))} \\ &= S_g(N_{f(x)}) \\ &= S_g(S_f(N_x)) \\ &= (S_g \circ S_f)(N_x) \end{aligned}$$

yielding thereby  $S_{gof} = S_g \circ S_f$ .

**Corollary 4.1.** If  $I_P : P \rightarrow P$  is an identity homomorphism in  $P(X)$ , then  $S_{I_P} : (X-P) \rightarrow (X-P)$  is an identity morphism in  $F(X)$ .

In view of the above discussion, we can define a covariant functor  $S : P(X) \rightarrow F(X)$  such that

- (i)  $S(P) = (X-P)$ , for all  $P \in P(X)$ ,
- (ii) for any morphism  $f : P_1 \rightarrow P_2$  in  $P(X)$ , we have  $S(f) = S_f : (X-P_1) \rightarrow (X-P_2)$  in  $F(X)$ .

## 5. Functors by Self Maps

Consider the category  $BCK(i_+)$  in which the objects of the category are positive implicative BCK-algebras and morphisms of the category are the BCI-homomorphisms between them.

**Proposition 5.1.** If  $f : X \rightarrow Y$  is a BCK-homomorphism, then the map  $L_f : L(X) \rightarrow L(Y)$  defined by  $L_f(l_x) = l_{f(x)}$  for all  $x \in X$ , is a BCK-homomorphism.

**Proof** For any  $l_{x_1}, l_{x_2} \in L(X)$ , we have

$$\begin{aligned}
 L_f(l_{x_1} o l_{x_2}) &= L_f(l_{x_1 \star x_2}) \\
 &= l_{f(x_1 \star x_2)} \\
 &= l_{f(x_1) \star f(x_2)} \\
 &= l_{f(x_1)} o l_{f(x_2)} \text{ [By the definition of composition} \\
 &\quad \text{of maps in } L(X)] \\
 &= L_f(l_{x_1}) o L_f(l_{x_2})
 \end{aligned}$$

which shows that  $L_f : L(X) \rightarrow L(Y)$  is a BCK-homomorphism.

**Theorem 5.1.** If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are BCK-homomorphisms in  $B_{CK}(i_+)$ , then  $L_{gof} = L_g o L_f$ .

**Proof** In view of Proposition 5.1, the maps  $L_f : L(X) \rightarrow L(Y)$ ,  $L_g : L(Y) \rightarrow L(Z)$  and their composition  $L_{gof} : L(X) \rightarrow L(Z)$  are morphisms in the category  $B_{CK}(i_+)$ .

For any  $l_x \in L(X)$ , we have

$$\begin{aligned}
 L_{gof}(l_x) &= l_{gof(x)} \\
 &= l_{g(f(x))} \\
 &= L_g(l_{f(x)}) \\
 &= L_g(L_f(l_x)) \\
 &= (L_g o L_f)(l_x)
 \end{aligned}$$

yielding thereby  $L_{gof} = L_g o L_f$ .

**Corollary 5.1.** If  $I_X : X \rightarrow X$  is an identity homomorphism in  $B_{CK}(i_+)$ , then  $L_{I_X} : L(X) \rightarrow L(X)$  is also an identity homomorphism in  $B_{CK}(i_+)$ .

With the help of Proposition 5.1, Theorem 5.1 and Corollary 5.1, we can define a functor  $F : B_{CK}(i_+) \rightarrow B_{CK}(i_+)$  such that

- (i)  $F(X) = L(X)$  for all  $X \in B_{CK}(i_+)$ ,
- (ii) for any morphism  $f : X \rightarrow Y$  in  $B_{CK}(i_+)$ , we have  $F(f) = L_f : L(X) \rightarrow L(Y)$  in  $B_{CK}(i_+)$ .



**Acknowledgement:** The authors are greatly indebted to the referee for several helpful suggestions and healthy comments.

## References

- [1] T.S. Blyth, *Categories*, Longman, New York 1986.
- [2] M.A. Chaudhry and S.A. Bhatti, *On the category of BCI-algebras*, Punjab Univ. J. Math. 21 (1988), 34-39.
- [3] Z.M. Chen and H.X. Wang, *Some universal properties of BCI-algebras*, Kobe J. Math. 6 (1989), 43-48.
- [4] E.Y. Deeba, *Filter theory of BCK-algebras*, Math. Japon. 25 (1980), 631-639.
- [5] C.S. Hoo, *Injectives in the category of BCI-algebras*, Math. Japon. 33 (1988), 237-246.
- [6] Y. Imai and K. Iseki, *On axiom systems of propositional calculi XIV*, Proc. Japan Acad. 42 (1966), 19-22.
- [7] K. Iseki, *An algebra related with a propositional calculus*, Proc. Japan Acad. 42 (1966), 26-29.
- [8] K. Iseki, *On ideals in BCI-algebras*, Math. Sem. Notes, Kobe Univ. 3 (1975), 1-12.
- [9] K. Iseki, *On BCI-algebras*, Math. Sem. Notes, Kobe Univ. 8 (1980), 125-130.
- [10] K. Iseki, *On the existence of quasi-commutative BCI-algebras*, Math. Sem. Notes, Kobe Univ. 8 (1980), 181-186.
- [11] K. Iseki, *On variety of BCI-algebras*, Math. Sem. Notes, Kobe Univ. 8 (1980), 225-226.
- [12] K. Iseki, *A problem on BCI-algebras*, Math. Sem. Notes, Kobe Univ. 8 (1980), 235-236.
- [13] K. Iseki and S. Tanaka, *An introduction to the theory of BCK-algebras*, Math. Japon. 23 (1978), 1-26.
- [14] K. Iseki and A.B. Thaheem, *Note on BCI-algebras*, Math. Japon. 29 (1984), 255-258.
- [15] M. Kondo, *Positive implicative BCK-algebra and its dual algebra*, Math. Japon. 35 (1990), 289-291.
- [16] S.M.A. Zaidi and S. Khan, *A note on the category of BCK-algebras with regular morphisms*, Proc. Internat. Conf. Algebra and its Applications, A.M.U. Aligarh (2001), 305-315.