

# THE P-ADIC FIELD CASE OF THE FUNCTIONAL EQUATION $P(f) = Q(g)$

Nguyen Trong Hoa

*Daklak Pedagogical College, Buon Ma Thuot, Vietnam  
email:ntronghoa@math.ac.vn*

## Abstract

In this paper, we study the existence of non constant meromorphic solutions  $f$  and  $g$  of the functional equation  $P(f) = Q(g)$ , where  $P(z)$  and  $Q(z)$  are two given nonlinear polynomials with coefficients in the non-Archimedean field  $\mathbb{K}$ .

## 1. Introduction

The Tenth problem of Hilbert is asked to establish an algorithm for finding all integer solutions of  $F(x, y) = 0$ , where  $F(x, y)$  is a polynomial with integer coefficients. It is natural to study the analogous of Hilbert's Tenth problem in the field of meromorphic functions. More specifically, we ask what forms of equation  $F(x, y) = 0$ , where  $F(x, y)$  is a polynomial with complex coefficients, may or may not have non constant meromorphic functions  $f$  and  $g$  that satisfy  $F(f, g) = 0$ ? Earlier in 1920s, as a simple application of his value distribution theory, Nevanlinna proved that a non constant meromorphic function ( in the complex plane) is uniquely determined by the inverse image of five distinct values (including infinity), ignoring multiplicity. Gross [11] extended this study by considering pre-images of a set and posed the question: *Is there a finite set  $A$  so that an entire (meromorphic) function is uniquely determined by the pre-image of the set  $A$ , counting multiplicities?* Let  $f$  be a non-constant meromorphic

---

The author is partially supported by the National Basic Research Program of Vietnam

**Key words:** non-Archimedean field, Hilbert's Tenth problem, meromorphic function, hyperbolicity, genus.

2000 Mathematics Subject Classification:

function and  $S$  be a subset of distinct elements. Define

$$E_f(S) = \bigcup_{a \in S} \{(z, m) \mid f(z) = a \text{ with multiplicity } m\}.$$

Two functions  $f$  and  $g$  of the same type are said to *share*  $S$ , *counting multiplicity*, if  $E_f(S) = E_g(S)$ . A subset  $S$  is called a *unique range set* (a *URS* in short) for entire (or meromorphic) functions if for any two non-constant entire (or meromorphic) functions  $f$  and  $g$  such that  $E_f(S) = E_g(S)$ , one has  $f=g$ . Assume that  $S$  be a finite set, we set :

$$P_S(z) = \prod_{a \in S} (x - a).$$

As a connection to the study of the uniqueness problem, Li and Yang [3] introduced the following definition:

**Definition 1** A non-constant polynomial  $P(z)$  is said to be a *unique polynomial* for entire (or meromorphic) functions if for two non-constant entire (or meromorphic) functions  $f$  and  $g$ ,  $P(f) = P(g)$  implies that  $f = g$ .

$P(z)$  is said to be a *strong uniqueness polynomial* for entire (or meromorphic) functions if for two non-constant entire (or meromorphic) functions  $f, g$  and some nonzero constant  $c$ , the condition  $P(f) = cP(g)$  implies that  $c = 1$  and  $f = g$ .

To demonstrate that the finite set  $S$  be a URS for entire (or meromorphic) functions, we prove that the polynomial  $P_S(z)$  is a strong uniqueness polynomial. If  $P$  is a strong uniqueness polynomial for entire (or meromorphic) functions, then the set of the zeros of  $P$  can be a URS.

Recently, H.H. Khoai and C.C. Yang [1] generalized the above studies by considering a pair of two nonlinear polynomials  $P(z)$  and  $Q(z)$  such that the only meromorphic solutions  $f, g$  satisfying  $P(f) = Q(g)$  are constants. This problem is considered in the complex plane  $\mathbb{C}$  by H.H. Khoai and C.C. Yang [1] as well as by C.C. Yang and P. Li [2].

In this paper, we find the conditions for the existence of non-constant meromorphic function solutions  $f$  and  $g$  of the functional equation  $P(f) = Q(g)$  in  $\mathbb{K}$  with  $\mathbb{K}$  being an algebraically closed field, complete for a non-trivial non-Archimedean absolute value. To solve the functional equation, we study the hyperbolicity of the algebraic curve  $\{P(x) - Q(y) = 0\}$  by estimating its genus. We shall do this by giving sufficiently many linear independent regular 1-forms of Wronskian type on that curve.

## 2. Main theorems

**Theorem 2.1** *Let  $P(z)$  and  $Q(z)$  be two nonlinear polynomials of degrees  $n$  and  $m$ , respectively, with  $n \geq m$ . Suppose that  $P(\alpha) \neq Q(\beta)$  for all zeros  $\alpha$*

of  $P'$  and  $\beta$  of  $Q'$ . Then there exist no non-constant meromorphic functions  $f$  and  $g$  such that  $P(f) = Q(g)$ , if  $n$  and  $m$  satisfy the following condition:

$$n \geq m \geq 2 \text{ and } n \geq 3.$$

**Theorem 2.2** Let  $P(z)$  and  $Q(z)$  be two nonlinear polynomials of degrees  $n$  and  $m$ , respectively. Then there exist no non-constant meromorphic functions  $f$  and  $g$  such that  $P(f) = Q(g)$  provided that  $P$  and  $Q$  satisfy one of the following conditions:

1. There exists a zero  $\beta_1$  of  $Q'$  with multiplicity  $m_1$  at least 2 and  $P(\alpha) \neq Q(\beta_1)$ , for all zeros  $\alpha$  of  $P'$ .
2. There exists two simple zeros  $\beta_1, \beta_2$  of  $Q'$  such that  $P(\alpha) \neq Q(\beta_i)$  for all zeros  $\alpha$  of  $P'$  and  $i = 1, 2$ .

**Definition 2** Let  $P(z)$  be a nonlinear polynomial of degree  $n$  whose derivative is given by:

$$P'(z) = c(z - \alpha_1)^{n_1} \dots (z - \alpha_k)^{n_k},$$

where  $n_1 + \dots + n_k = n - 1$  and  $\alpha_1, \dots, \alpha_k$  are distinct zeros of  $P'$ . The number  $k$  is called the *derivative index* of  $P$ .

A polynomial  $P(z)$  is said to satisfy the *condition of separating the roots of  $P'$*  (separation condition) if  $P(\alpha_i) \neq P(\alpha_j)$  for all  $k \geq i \neq j \geq 1$ .

**Theorem 2.3** Let  $P(z)$  and  $Q(z)$  be two polynomials defined by (1) and let  $P$  satisfy the separation condition. Suppose that  $\beta_1, \dots, \beta_J$  are distinct zeros of  $Q'$  with multiplicity  $m_j$ , respectively, such that for every  $\beta_j$ ,  $j = 1, 2, \dots, J$ , there exists zeros  $\alpha_i$  of  $P'$  with  $P(\alpha_i) = Q(\beta_j)$ . Then there exist no non-constant meromorphic functions  $f$  and  $g$  such that  $P(f) = Q(g)$  if

$$m - \sum_{j=1}^J m_j \geq 3.$$

**Remark 2.4** In case  $\deg P = \deg Q = 2$ , the equation  $P(f) = Q(g)$  has some non-constant meromorphic functions. Indeed, in this case we can rewrite the equation  $P(f) = Q(g)$  in the form:

$$(f - a)^2 = (bg - c)^2 + d,$$

where  $a, b, c, d \in \mathbb{K}$  and  $b \neq 0$ . Hence

$$(f - bg - a + c)(f + bg - a - c) = d.$$

Assume that  $h$  is a non-constant meromorphic function, we set

$$f = \frac{1}{2}\left(h + \frac{d}{h}\right) + a, \quad g = \frac{1}{2b}\left(-h + \frac{d}{h}\right) + \frac{c}{b}.$$

Then  $f$  and  $g$  are non-constant meromorphic solutions of equation  $P(f) = Q(g)$ .

### 3. Lemmas and Proofs

Let  $\mathbb{K}$  be an algebraically closed field, complete for a non-trivial non-Archimedean absolute value with characteristic zero. Suppose that  $H(X, Y, Z)$  is a homogeneous polynomial of degree  $n$  and

$$C := \{(X, Y, Z) \in \mathbb{P}^2(\mathbb{K}) \mid H(X, Y, Z) = 0\}.$$

Denote

$$\begin{aligned} W_1 &= W(X, Y) = \begin{vmatrix} X & Y \\ dX & dY \end{vmatrix}, \\ W_2 &= W(Y, Z) = \begin{vmatrix} Y & Z \\ dY & dZ \end{vmatrix}, \\ W_3 &= W(X, Z) = \begin{vmatrix} X & Z \\ dX & dZ \end{vmatrix}. \end{aligned}$$

Assume that  $R(X, Y, Z)$  and  $S(X, Y, Z)$  are two homogeneous polynomials in  $\mathbb{P}^2(\mathbb{K})$ . Let

$$\omega_i = \frac{R(X, Y, Z)}{S(X, Y, Z)} W_i,$$

with  $i = 1, 2, 3$ . If  $R(X, Y, Z)$  and  $S(X, Y, Z)$  satisfy  $\deg S = \deg R + 2$ , then  $\omega_i$  is a well-defined rational 1-form on  $\mathbb{P}^2(\mathbb{K})$ .

**Definition 3** Let  $C$  be an algebraic curve in  $\mathbb{P}^2(\mathbb{K})$ . A 1-form  $\omega$  on  $C$  is said to be *regular* if it is the pull-back of a rational 1-form on  $\mathbb{P}^2(\mathbb{K})$  such that the pole set of  $\omega$  does not intersect  $C$ . A well-defined rational regular 1-form on  $C$  is said to be an *1-form of Wronskian type*.

Notice that to solve the functional equation  $P(f) = Q(g)$ , is the same as to find meromorphic functions  $f, g$  on  $\mathbb{K}$  such that  $(f(z), g(z))$  in curve  $\{P(x) - Q(y) = 0\}$ . On the other hand, if  $C$  is hyperbolic on  $\mathbb{K}$  and suppose that  $f, g$  be meromorphic functions such that  $(f(z), g(z)) \in C$ , where  $z \in \mathbb{K}$ , then  $f$  and  $g$  are constant (see.[6]). Therefore, to show that this equation has no non-constant solutions, we shall prove the hyperbolicity of  $\{P(x) - Q(y) = 0\}$ . By Picard-Berkovich's theorem in the  $p$ -adic case, a curve  $C$  in  $\mathbb{K}$  is Brody hyperbolic if and only if the genus of the curve  $C$  is at least 1.

It is well-known that the genus  $g$  of an algebraic curve  $C$  is equal to the dimension of the space of regular 1-forms on  $C$ . Therefore, to compute the genus, we have to construct a basis of the space of regular 1-forms on  $C$ .

Now, let  $P$  and  $Q$  be two nonlinear polynomials of degrees  $n$  and  $m$  in  $\mathbb{K}$ , respectively, with

$$\begin{aligned} P(x) &= a_n x^n + \dots + a_1 x + a_0 \\ Q(y) &= b_m y^m + \dots + b_1 y + b_0. \end{aligned} \tag{1}$$

Without loss of generality, we assume that  $m \geq n$ . Set

$$F_1(x, y) := P(x) - Q(y),$$

$$F(X, Y, Z) := Z^n \left\{ P\left(\frac{X}{Z}\right) - Q\left(\frac{Y}{Z}\right) \right\}. \quad (2)$$

$$C := \{(X, Y, Z) \in \mathbb{P}^2(\mathbb{K}) \mid F(X, Y, Z) = 0\}. \quad (3)$$

We conclude

$$P'(x) = na_n x^{n-1} + \dots + 2a_2 x + a_1 = na_n (x - \alpha_1)^{n_1} \dots (x - \alpha_k)^{n_k}, \quad (4)$$

$$Q'(y) = mb_m y^{m-1} + \dots + 2b_2 y + b_1 = mb_m (y - \beta_1)^{m_1} \dots (y - \beta_l)^{m_l}, \quad (5)$$

where  $n_1 + \dots + n_k = n - 1$ ;  $m_1 + \dots + m_l = m - 1$ ,  $\alpha_1, \dots, \alpha_k$  are distinct zeros of  $P'$ ; and  $\beta_1, \dots, \beta_l$  are distinct zeros of  $Q'$ . Define

$$P'(X, Z) := Z^{n-1} P'\left(\frac{X}{Z}\right); \quad Q'(Y, Z) := Z^{m-1} Q'\left(\frac{Y}{Z}\right).$$

Then

$$\begin{aligned} \frac{\partial F}{\partial X} &= P'(X, Z), \\ \frac{\partial F}{\partial Y} &= -Z^{n-m} Q'(Y, Z), \\ \frac{\partial F}{\partial Z} &= \sum_{i=0}^{n-1} (n-i) a_i X^i Z^{n-1-i} - \sum_{j=0}^{m'} (n-j) b_j Y^j Z^{n-1-j}, \end{aligned}$$

where

$$m' = \begin{cases} n-1 & \text{if } n = m \\ m & \text{if } n > m. \end{cases}$$

Then, by Euler's theorem, for all points  $(X, Y, Z) \in C$ , we have

$$\frac{\partial F}{\partial X} X + \frac{\partial F}{\partial Y} Y + \frac{\partial F}{\partial Z} Z = 0. \quad (6)$$

The equation of the tangent space of  $C$  at the point  $(X, Y, Z) \in C$  is defined by

$$\frac{\partial F}{\partial X} dX + \frac{\partial F}{\partial Y} dY + \frac{\partial F}{\partial Z} dZ = 0. \quad (7)$$

From (6) and (7), we obtain

$$\begin{aligned} \frac{\partial F}{\partial X} &= \frac{\begin{vmatrix} Y & Z \\ dY & dZ \end{vmatrix}}{\begin{vmatrix} X & Y \\ dX & dY \end{vmatrix}} \frac{\partial F}{\partial Z}, \\ \frac{\partial F}{\partial Y} &= \frac{\begin{vmatrix} Z & X \\ dZ & dX \end{vmatrix}}{\begin{vmatrix} X & Y \\ dX & dY \end{vmatrix}} \frac{\partial F}{\partial Z}. \end{aligned}$$

Hence, we have

$$\frac{\frac{W(Y, Z)}{\frac{\partial F}{\partial X}}}{\frac{\partial F}{\partial X}} = \frac{\frac{W(Z, X)}{\frac{\partial F}{\partial Y}}}{\frac{\partial F}{\partial Y}} = \frac{\frac{W(X, Y)}{\frac{\partial F}{\partial Z}}}{\frac{\partial F}{\partial Z}}. \quad (8)$$

Set

$$\eta := \frac{W(Y, Z)}{\frac{\partial F}{\partial X}} = \frac{W(Z, X)}{\frac{\partial F}{\partial Y}} = \frac{W(X, Y)}{\frac{\partial F}{\partial Z}},$$

we obtain

$$\begin{aligned} \eta &= \frac{W(Y, Z)}{P'(X, Z)} = \frac{W(X, Z)}{Z^{n-m}Q'(Y, Z)} \\ &= \frac{W(X, Y)}{\sum_{i=0}^{n-1} (n-i)a_i X^i Z^{n-1-i} - \sum_{j=0}^{m'} (n-j)b_j Y^j Z^{n-1-j}}. \end{aligned} \quad (9)$$

In order to prove the main results, we need the following lemmas.

**Lemma 3.1** *Let  $P$  and  $Q$  be two nonlinear polynomials of degrees  $n$  and  $m$ , respectively, (defined by (1)) with  $n \geq m$ , and let  $C$  be a projective curve defined by (3). If  $P(\alpha_i) \neq Q(\beta_j)$  for all zeros  $\alpha_i$  of  $P'$  and  $\beta_j$  of  $Q'$ , (defined by (4), (5)), then we have the following assertions:*

1. *If  $n = m$  or  $n = m + 1$ , then  $C$  is non-singular in  $\mathbb{P}^2(\mathbb{K})$ .*
2. *If  $n - m \geq 2$ , then the point  $(0, 1, 0)$  is a unique singular point of  $C$  in  $\mathbb{P}^2(\mathbb{K})$ .*

**Proof** By the hypothesis of the lemma,  $P(\alpha_i) \neq Q(\beta_j)$  for all zeros  $\alpha_i$  of  $P'$  and  $\beta_j$  of  $Q'$ , we conclude that  $C$  is non-singular in  $\mathbb{P}^2(\mathbb{K}) \setminus \{Z = 0\}$ . Now we consider the singularity of  $C$  in  $\{Z = 0\}$ . Assume that  $(X, Y, 0)$  is a singular point of  $C$ . We obtain

$$\frac{\partial F}{\partial X}(X, Y, 0) = 0; \frac{\partial F}{\partial Y}(X, Y, 0) = 0; \frac{\partial F}{\partial Z}(X, Y, 0) = 0 \text{ and } F(X, Y, 0) = 0.$$

We consider the following three cases:

If  $n = m$ , then

$$\begin{aligned} na_n X^{n-1} &= 0 \\ nb_n Y^{n-1} &= 0 \\ a_{n-1} X^{n-1} - b_{n-1} Y^{n-1} &= 0 \\ a_n X^n - b_n Y^n &= 0, \end{aligned}$$

has no root in  $\mathbb{P}^2(\mathbb{K})$ . If  $n = m + 1$ , then

$$\begin{aligned} na_n X^{n-1} &= 0 \\ a_n X^n - b_{n-1} Y^{n-1} &= 0 \\ a_n X^n &= 0, \end{aligned}$$

has no root in  $\mathbb{P}^2(\mathbb{K})$ .

If  $n - m \geq 2$ , then

$$\begin{aligned} na_n X^{n-1} &= 0 \\ a_{n-1} X^{n-1} &= 0 \\ a_n X^n &= 0, \end{aligned}$$

has a unique root  $(0, 1, 0)$  in  $\mathbb{P}^2(\mathbb{K})$  and  $C$  is singular with a unique singular point at  $(0, 1, 0)$ .

Thus, if  $n = m$  or  $n = m + 1$ , then  $C$  is a smooth curve. If  $n - m \geq 2$ , then  $C$  is singular with a unique singular point at  $(0, 1, 0)$ .  $\square$

**Proposition 3.2** *If  $n = m$  or  $n = m + 1$  and the hypotheses of Lemma 3.1 are satisfied, then  $C$  is Brody hyperbolic on  $\mathbb{K}$  for all  $n \geq 3$ .*

**Proof** Consider the 1-form defined by (9). Since  $C$  is smooth, then by Lemma 3.1, if one of the expressions of  $\eta$  is regular, we can conclude that the other expressions are also regular. Let

$$\begin{aligned} \pi : \mathbb{K}^3 \setminus \{(0, 0, 0)\} &\implies \mathbb{P}^2(\mathbb{K}) \\ (X, Y, Z) &\longmapsto [X : Y : Z], \end{aligned}$$

where  $[X : Y : Z] := \{(\lambda X, \lambda Y, \lambda Z) \mid \lambda \in \mathbb{K}^* = \mathbb{K} \setminus \{0\}\}$ . Note that  $\eta$  is regular on  $\pi^{-1}(C)$ . Then the 1-form

$$\omega := \frac{W(X, Z)}{Z^{n-m} Q'(Y, Z)} Z^{n-3} = Z^{n-3} \eta,$$

is a well-defined rational regular 1-form on  $C$  if  $n \geq 3$ . Thus, if  $n \geq 4$ , then for any homogeneous polynomial  $R(X, Y, Z)$  of degree  $n - 3$ , the 1-form  $\omega := R(X, Y, Z) \eta$ , is regular on  $C$  and vanishes along  $\{(X, Y, Z) \mid R(X, Y, Z) = 0\}$ . Note that the set of monomials of degree  $n - 3$  in  $X, Y, Z$  is a basis of the vector space of homogeneous polynomials of degree  $n - 3$ . This basis only has  $\frac{(n-1)(n-2)}{2}$  vectors. Hence, the genus of  $C$  is

$$g = \frac{(n-1)(n-2)}{2},$$

Therefore,  $C$  is Brody hyperbolic if  $n \geq 3$ .  $\square$

**Remark 3.3** We also require that the 1-form defined by (9) is non trivial when it restricts to a component of  $\pi^{-1}(C)$ . This is equivalent to the condition that the nominators are not identically zero when they restrict to a component of  $\pi^{-1}(C)$ , i.e. the Wronskians  $W_i$ , (with  $i = 1, 2, 3$ ) are not identically zero. It means that the homogeneous polynomial defining  $C$  has no linear factor of the

form  $aX - bY$ ,  $aY - bZ$  or  $aX - bZ$ , with  $a, b \in \mathbb{K}$ . Indeed, we assume on the contrary that, for example,  $aX - bZ$  is a factor of curve  $C$  defined by (3). Without loss of generality, we can take  $a \neq 0$ . Since  $aX - bZ$  is a factor of  $F(X, Y, Z)$ , we have

$$0 = F\left(\frac{b}{a}Z, Y, Z\right) = Z^n \left\{ P\left(\frac{\frac{b}{a}Z}{Z}\right) - Q\left(\frac{Y}{Z}\right) \right\} = Z^n \left\{ P\left(\frac{b}{a}\right) - Q\left(\frac{Y}{Z}\right) \right\},$$

and then we obtain  $P\left(\frac{b}{a}\right) \equiv Q\left(\frac{Y}{Z}\right)$  for all  $Y, Z$ . It is impossible.

**Proposition 3.4** *If  $n \geq m + 2$  and the hypotheses of Lemma 3.1 are satisfied, then  $C$  is Brody hyperbolic on  $\mathbb{K}$  for all  $m \geq 3$ .*

**Proof** Since  $n \geq m + 2$  by Lemma 3.1, we conclude that  $C$  is singular with a unique singular point at  $(0, 1, 0)$ . Considering the 1-form defined by (9)

$$\begin{aligned} \eta &= \frac{W(Y, Z)}{P'(X, Z)} = \frac{W(X, Z)}{Z^{n-m}Q'(Y, Z)} \\ &= \frac{W(X, Y)}{\sum_{i=0}^{n-1} (n-i)a_i X^i Z^{n-1-i} - \sum_{j=0}^{m'} (n-j)b_j Y^j Z^{n-1-j}}. \end{aligned}$$

Due to Remark 3.3, we deduce that  $\eta$  is not identically zero when it restricts to a component of  $\pi^{-1}(C)$  and it is not regular on  $C$ . We take

$$\begin{aligned} \xi &:= \frac{Z^{n-m}W(Y, Z)}{P'(X, Z)} = \frac{W(X, Z)}{Q'(Y, Z)} \\ &= \frac{Z^{n-m}W(X, Y)}{\sum_{i=0}^{n-1} (n-i)a_i X^i Z^{n-1-i} - \sum_{j=0}^{m'} (n-j)b_j Y^j Z^{n-1-j}} \\ &= Z^{n-m}\eta. \end{aligned}$$

Then,  $\xi$  is regular on  $\pi^{-1}(C)$  because the denominators of  $\xi$  have no common zeros. Therefore

$$\omega := \frac{W(X, Z)}{Q'(Y, Z)} Z^{m-3} = Z^{m-3}\xi,$$

is well-defined on  $C$ , regular and vanishing along  $(n-3)\{Z=0\}$ . This implies that, if  $m \geq 3$ , then  $\omega$  is a 1-form of Wronskian type on  $C$ .

If  $m = 3$  then  $\omega = \xi$  is a linear independent regular 1-form of Wronskian type on the curve  $C$ .

If  $m \geq 4$ , we take  $R_1, R_2, \dots, R_{\frac{(m-2)(m-1)}{2}}$  as a basis of monomials of degree  $m-3$  in  $\{X, Y, Z\}$ , then

$$\{R_i \omega \mid i = 1, 2, \dots, \frac{(m-2)(m-1)}{2}\},$$



are linearly independent and global regular 1-forms of Wronskian type on the curve  $C$ . Thus, the genus  $g_C$  of  $C$  is

$$g_C \geq \frac{(m-2)(m-1)}{2}.$$

Therefore,  $C$  is Brody hyperbolic if  $m \geq 3$ . □

Next, we recall some notations. Let  $C$  be a curve on  $\mathbb{K}$  defined by a homogeneous polynomial  $F(X, Y, Z) = 0$  and let  $\rho$  be a point of  $C$ . A holomorphic map

$$\phi = (\phi_1, \phi_2, \phi_3) : \Delta_\epsilon = \{t \in \mathbb{K} \mid |t| < \epsilon\} \longrightarrow C,$$

with  $\phi(0) = \rho$ , is referred to a *holomorphic parameterization* of  $C$  at  $\rho$ . Local holomorphic parameterization always exists for sufficiently small  $\epsilon$ . If  $\phi$  is a local holomorphic parameterization of  $C$  at  $\rho$ , then the Laurent expansion of  $F \circ \phi(t)$  at  $\rho$  has the form

$$F \circ \phi(t) = \sum_{i=p}^q c_i t^i, \quad c_p \neq 0, \quad p < q < \infty.$$

The order of  $F$  at  $\rho$ , (it is also the order of  $F \circ \phi(t)$  at  $t = 0$ ) is defined by  $p$  and denoted by

$$p := \text{ord}_{\rho, \phi} F = \text{ord}_{t=0} F(\phi(t)).$$

Assume that  $\varphi(x, y)$  is an analytic function in  $x, y$  and it is singular at  $\rho = (a, b)$ . The Puiseux expansion of  $\varphi(x, y)$  at  $\rho$  is given by

$$x - a = \sum_{i=0}^{\infty} a_{\alpha+i} t^{\alpha+i}, \quad y - b = \sum_{j=0}^{\infty} b_{\beta+j} t^{\beta+j},$$

$$\varphi\left(a + \sum_{i=0}^{\infty} a_{\alpha+i} t^{\alpha+i}, b + \sum_{j=0}^{\infty} b_{\beta+j} t^{\beta+j}\right) = 0.$$

The  $\alpha$  (respectively,  $\beta$ ) is the order (also the multiplicity number) of  $x$  at  $\rho$ , (respectively, the order of  $y$  at  $\rho$ ) for  $\varphi$  and is denoted by

$$\alpha := \text{ord}_{\rho, \varphi} x \quad (\text{respectively, } \beta := \text{ord}_{\rho, \varphi} y).$$

We now have the following result

**Proposition 3.5** *Let  $P$  be a nonlinear polynomial of degree  $n \geq 4$  and  $Q = b_2 y^2 + b_1 y + b_0$ , where  $b_2 \neq 0$  and  $C$  a projective curve defined by (3). If  $P(\alpha_i) \neq Q(\beta)$  for all zeros  $\alpha_i$  of  $P'$  and  $\beta$  of  $Q'$  defined by (4), then  $C$  is Brody hyperbolic on  $\mathbb{K}$  for all  $n \geq 4$ .*

**Proof** Since  $n \geq 4$  and  $m = 2$ , by Lemma 3.1, we conclude that  $C$  is singular with a unique singular point at  $(0, 1, 0)$ . Considering polynomial

$$H(X, Z) := F(X, 1, Z) = \sum_{i=0}^n a_{n-i} X^{n-i} Z^i - Z^{n-2}(b_2 + b_1 Z + b_0 Z^2), \text{ with } a_n, b_2 \neq 0.$$

Since polynomial  $H(X, Z)$  is singular at  $(0, 0)$ , by the Newton diagram of the Puiseux expansion of  $H$  at  $(0, 0)$ , we have

$$\text{ord}_{0,H} X = \frac{n-2}{d}, \quad \text{ord}_{0,H} Z = \frac{n}{d},$$

where  $d$  is the greatest common divisor of  $n$  and  $n-2$ .

It follows from (9) that

$$\begin{aligned} \eta &= \frac{W(Y, Z)}{P'(X, Z)} \\ &= \frac{W(X, Z)}{Z^{n-2}(2b_2 Y + b_1)} \\ &= \frac{W(X, Y)}{\left\{ \sum_{i=0}^{n-1} (n-i) a_i X^i Z^{n-1-i} \right\} - Z^{n-3}(b_2 Y^2 + b_1 Y Z + b_0 Z^2)}. \end{aligned}$$

By Remark 3.3,  $\eta$  is non trivial on  $C$ . The  $\eta$  is regular on  $\pi^{-1}(C \setminus \{Z = 0\})$  because the denominators of  $\eta$  have no common zeros for  $Z \neq 0$ . Let

$$\omega := Z^{n-3} \eta = \frac{W(X, Z)}{Z(2b_2 Y + b_1)}.$$

By the proof above,  $\omega$  is regular on  $\pi^{-1}(C \setminus \{Z = 0\})$ . In  $C \cap \{Z = 0\}$ , the denominators of  $\omega$  only vanish at  $\rho = (0, 1, 0)$ . Note that at this point  $H(X, Z) = F(X, 1, Z)$  is singular, and therefore

$$\begin{aligned} \text{ord}_{t=0} \omega &= \text{ord}_{t=0} W(X, Z) - \text{ord}_{0,H} Z \\ &= \text{ord}_{t=0} (XdZ - ZdX) - \text{ord}_{0,H} Z \\ &= \frac{n-2-d}{d}. \end{aligned}$$

Since  $d = (n, n-2)$ , it follows that either  $d = 1$  or  $d = 2$ . If  $d = 1$ , then  $\text{ord}_{t=0} \omega = n-3$ . If  $d = 2$ , then  $\text{ord}_{t=0} \omega = \frac{n-4}{2}$ .

Consequently, if  $n \geq 4$ , then  $\omega$  is a non trivial well-defined regular 1-form of Wronskian type on  $C$  and the curve  $C$  is Brody hyperbolic.  $\square$

We now consider the case when the curve  $C$  is singular in  $\mathbb{P}^2(\mathbb{K}) \setminus \{Z = 0\}$ .

From Lemma 3.1, the curve  $C$  is only singular in  $\mathbb{P}^2(\mathbb{K}) \setminus \{Z = 0\}$  at  $(\alpha_i, \beta_j, 1)$ , with  $\alpha_1, \dots, \alpha_k$  are distinct zeros of  $P'$  and  $\beta_1, \dots, \beta_l$  are distinct

zeros of  $Q'$ . By (4),(5), the distinct zeros  $\alpha_1, \dots, \alpha_k$  of  $P'$  have multiplicities  $n_1, \dots, n_k$  and the distinct zeros  $\beta_1, \dots, \beta_l$  of  $Q'$  have multiplicities  $m_1, \dots, m_l$ . Let

$$\begin{aligned}\Gamma &:= \{(\alpha_i, \beta_j, 1) \text{ be singular points of } C\}, \\ \Delta &:= \{\alpha_i | (\alpha_i, \beta_j, 1) \text{ be singular points of } C\}\end{aligned}$$

and

$$\Lambda := \{\beta_j | (\alpha_i, \beta_j, 1) \text{ be singular points of } C\}.$$

Setting  $I = \#\Delta$ ,  $J = \#\Lambda$ , then we have  $k \geq I$  and  $l \geq J$ . Without loss of generality, we can take

$$\Delta = \{\alpha_1, \dots, \alpha_I\}, \Lambda = \{\beta_1, \dots, \beta_J\}.$$

Assume that  $P$  satisfies the separation condition. Then, we conclude  $J \geq I$  and for every  $j$ ,  $j = 1, 2, \dots, J$ , there exists a unique value  $i_j$  such that  $(\alpha_{i_j}, \beta_j, 1)$  is a singular point of  $C$  (these  $\alpha_{i_j}$  can be equal to each other). Hence,

$$\Gamma = \{(\alpha_{i_j}, \beta_j, 1) | j = 1, 2, \dots, J\}, \text{ with } l \geq J. \quad (10)$$

Let

$$H_j = a(X - \alpha_{i_j}Z) + b(Y - \beta_jZ), \text{ with } a, b \neq 0.$$

We obtain  $H_j(\alpha_{i_j}, \beta_j, 1) = 0$ . Putting

$$\xi := \frac{H_j^{s_j}}{(Y - \beta_jZ)^{m_j}} W(X, Z), \text{ for } s_j = \max\{n_{i_j}, m_j\},$$

it follows that  $\xi$  is regular on  $C$ . Therefore, from (9) we obtain

$$\xi = \frac{mb_m Z^{n-m} H_j^{s_j} \prod_{j \neq t}^l (Y - \beta_t Z)^{m_t}}{na_n \prod_{\nu=1}^k (X - \alpha_\nu Z)^{n_\nu}} W(Y, Z).$$

Note that  $P$  satisfies the separation condition, then  $(\alpha_\nu, \beta_j, 1) \notin C$  with  $\nu \neq i_j$ .

On the other hand,  $\text{ord}_{\rho, \phi} H_j^{s_j} = s_j \geq n_{i_j}, m_j$  and hence  $\text{ord}_{\rho, \phi} \frac{H_j^{s_j}}{(X - \alpha_{i_j} Z)^{n_{i_j}}} \geq 0$ , for  $\rho = (\alpha_{i_j}, \beta_j, 1)$ .

Now, we set

$$\zeta := \frac{\prod_{j=1}^J H_j^{s_j}}{Q'(Y, Z)} W(X, Z).$$

Then  $\zeta$  is regular on  $C$ . Let

$$\omega := \left\{ \prod_{j=1}^J H_j^{s_j} \right\} Z^{n - \sum_{j=1}^J s_j - 3} \eta = Z^{m - \sum_{j=1}^J s_j - 3} \zeta,$$

we can easily see that  $\omega$  is a non trivial, well-defined regular 1-form of Wronskian type on  $C$  and the curve  $C$  is Brody hyperbolic if  $m - \sum_{j=1}^J s_j \geq 3$ .

We can summarize these facts in the following result.

**Proposition 3.6** *Let  $P(z)$  and  $Q(z)$  (defined by (1)) be two nonlinear polynomials of degrees  $n, m$ , respectively, with  $n \geq m$ , and let  $P$  satisfy the separation condition. Suppose that  $\Gamma = \{(\alpha_{i_j}, \beta_j, 1) | j = 1, 2, \dots, J\}$  (defined by (10)) is the set of singular points of  $C$ . Then the curve  $C$  is Brody hyperbolic if  $m - \sum_{j=1}^J s_j \geq 3$ , with  $s_j = \max(n_{i_j}, m_j), j = 1, 2, \dots, J$ .*

*Lemma 3.7* *Let  $P(z)$  and  $Q(z)$  be two polynomials defined by (1) and  $P$  satisfy the separation condition. Let  $\Gamma := \{(\alpha_i, \beta_j, 1) | j = 1, 2, \dots, J\}$  be the set of singular points of  $C$ . If  $m_j - n_i \geq 2$  at all points  $\rho_{ij} = (\alpha_i, \beta_j, 1)$ , then  $C$  is Brody hyperbolic.*

**Proof** The condition of Lemma 3.7 implies that the curve  $C$  is singular at any point  $\rho_{ij}$  and  $m_j \geq 3$ . Let

$$\zeta := \frac{(X - \alpha_i Z)^{m_j - 2}}{(Y - \beta_j Z)^{m_j}} W(X, Z)$$

be a well-defined rational 1-form on  $C$ . It follows from (9) that

$$\begin{aligned} \zeta &= \frac{mb_m Z^{n-m} (X - \alpha_i Z)^{m_j - 2} \prod_{1=t \neq j}^l (Y - \beta_t Z)^{m_t}}{na_n \prod_{\nu=1}^k (X - \alpha_\nu Z)^{n_\nu}} W(Y, Z) \\ &= \frac{mb_m Z^{n-m} (X - \alpha_i Z)^{m_j - n_i - 2} na_n \prod_{i \neq \nu=1}^k (X - \alpha_\nu Z)^{n_\nu}}{na_n \prod_{i \neq \nu=1}^k (X - \alpha_\nu Z)^{n_\nu}} W(Y, Z). \end{aligned}$$

The first expression implies that all poles of  $\zeta$  must belong to  $C \cap \{Y = \beta_j\}$ . The second expression implies that all poles of  $\zeta$  must belong to  $C \cap \{X = \alpha_\nu\}$  for all  $\nu \neq i$ . Since  $P$  satisfies the separation condition, we can conclude that  $Q(\beta_j) = P(\alpha_i) \neq P(\alpha_\nu)$  for all  $\nu \neq i$ . This shows that  $\zeta$  is regular on  $C$  if  $m_j - n_i \geq 2$ . The proof of the lemma is now complete.  $\square$

**Proof of Theorem 2.1** It follows immediately from Proposition 3.2, 3.4 and 3.5.  $\square$

**Proof of Theorem 2.2** By (9), we see that

$$\eta = \frac{W(Y, Z)}{na_n \prod_{\nu=1}^k (X - \alpha_\nu Z)^{n_\nu}} = \frac{W(X, Z)}{mb_m Z^{n-m} \prod_{t=1}^l (Y - \beta_t Z)^{m_t}}.$$

Set

$$\theta := \frac{W(X, Z)}{(Y - \beta_1 Z)^2}.$$

We then have

$$\theta = \frac{mb_m Z^{n-m} (Y - \beta_1 Z)^{m_1-2} \prod_{t=2}^l (Y - \beta_t Z)^{m_t}}{na_n \prod_{\nu=1}^k (X - \alpha_\nu Z)^{n_\nu}} W(Y, Z).$$

Since  $(\alpha_i, \beta_1, 1) \notin C$  with for all  $\alpha_i$ , we see that  $\theta$  is a well-defined regular 1-form of Wronskian type on  $C$ , proving the first assertion of the theorem.

Similary, let

$$\xi := \frac{W(X, Z)}{(Y - \beta_1 Z)(Y - \beta_2 Z)}.$$

Then  $\xi$  is a well-defined regular 1-form of Wronskian type on  $C$  and therefore the second assertion of theorem is proved.  $\square$

**Proof of Theorem 2.3** By applying Proposition 3.6 and Lemma 3.7, the Theorem is followed.  $\square$

## References

- [1] H.H. Khoai and C.C. Yang, *On the functional equation  $P(f) = Q(g)$* , Value Distribution Theory, Kluwer Academic Publishers (2004), 201 - 207.
- [2] C.C. Yang and P. Li, *Some further results on the functional equation  $P(f) = Q(g)$* , Value distribution theory and its trends, ACAA, Kluwer Academic Publisher (2004).
- [3] C.C. Yang and P. Li, *On the unique range sets of meromorphic functions*, Proc. Amer. Math. Soc., 124(1996), 177-195.
- [4] A. Escassut and C.C. Yang, *The functional equation  $P(f) = Q(g)$  in a  $p$ -adic field*, J. Number Theory, 105(2004) No.2, 344-360.
- [5] F. Gross and C.C. Yang, *On pre-images and range sets of meromorphic functions*, Proc. Japan Acad., 58(1982), 17-20.
- [6] P.C. Hu and C.C. Yang, "Meromorphic functions over non-Archimedean field", Kluwer Academic Publisher, (2000).
- [7] H. Fujimoto, *On uniqueness of meromorphic functions sharing finite sets*, Amer. J. Math., 122(2000), 1175-1203.
- [8] B. Shiffman, *Uniqueness of entire and meromorphic functions sharing finite sets*, Complex Variables, vol.4, (2001), 433-450.
- [9] T.T.H. An, J.T-Y. Wang and P-M. Wong, *Strong uniqueness polynomials: The complex case*, Complex Var. Theory and Appl. 49 (2004), N.1, 25-54.
- [10] F. Gross, *Factorization of meromorphic functions and some open problems*, Complex analysis (Proc. Conf., Univ. Kentucky, Lexington, Ky., 1976), Lecture Notes in Math., Vol.599, Springer, Berlin (1977) 51-67.
- [11] A. Boutabaa and A. Escassut, *Applications of the  $p$ -adic Nevanlinna theory to functional equations*, Annale de l' Institut Fourier', T50 (3) (2000), 751-766.