

## A COMPACT EMBEDDING OF SEMISIMPLE SYMMETRIC SPACES

Tran Dao Dong and Tran Vui

*\*\*Department of Mathematics,  
Faculty of Education, Hue University  
32 Le Loi Hue, Vietnam  
email:tddong@yahoo.com*

### Abstract

Let  $G$  be a connected real semisimple Lie group with finite center and  $\sigma$  be an involutive automorphism of  $G$ . Suppose that  $H$  is a closed subgroup of  $G$  with  $G_e^\sigma \subset H \subset G^\sigma$ , where  $G^\sigma$  is the fixed points group of  $\sigma$  and  $G_e^\sigma$  denotes its identity component. The coset space  $\mathbf{X} = G/H$  is then a semisimple symmetric space. Let  $\theta$  be a Cartan involution which commutes with  $\sigma$  and  $K$  be the set of all fixed points of  $\theta$ . Then  $K$  is a  $\sigma$ -stable maximal compact subgroup of  $G$  and the coset space  $G/K$  becomes a Riemannian symmetric space of noncompact type. By using the action of the Weyl group, we have constructed a compact real analytic manifold in which the Riemannian symmetric space  $G/K$  is realized as an open subset and that  $G$  acts analytically on it. The purpose of this note is to apply the above construction to the case of semisimple symmetric spaces  $\mathbf{X} = G/H$ . Our construction is similar to those of Schlichtkrull, Lizhen Ji, Oshima for Riemannian symmetric spaces and similar to those of Kosters, Sekiguchi, Oshima for semisimple symmetric spaces.

## 1 Introduction

Let  $G$  be a connected real semisimple Lie group with finite center,  $\sigma$  be an involutive automorphism of  $G$  and  $\mathbf{X} = G/H$  be the corresponding semisimple symmetric space. Here  $H$  is a closed subgroup of  $G$  with  $G_e^\sigma \subset H \subset G^\sigma$ , where  $G^\sigma$  is the fixed points group of  $\sigma$  and  $G_e^\sigma$  denotes its identity component.

Denote by  $\theta$  the Cartan involution which commutes with  $\sigma$  and  $K$  the set of all fixed points of  $\theta$ . Then  $K$  is a  $\sigma$ -stable maximal compact subgroup of  $G$ .

---

**Key words:** symmetric space, Weyl group, Cartan decomposition, compactification.  
2000 Mathematics Subject Classification:

Let  $\mathcal{G}$  be the Lie algebra of  $G$ . The involutions of  $\mathcal{G}$  induced by  $\sigma$  and  $\theta$  are denoted by the same letters, respectively.

Suppose that  $\mathcal{G} = \mathcal{H} \oplus \mathcal{Q} = \mathcal{K} \oplus \mathcal{P}$  are the decompositions of  $\mathcal{G}$  into +1 and -1 eigenspaces for  $\sigma$  and  $\theta$ , respectively, where  $\mathcal{H}$  (resp.  $\mathcal{K}$ ) is the Lie algebra of  $H$  (resp.  $K$ ). Fix a maximal Abelian subspace  $\mathcal{A}$  in  $\mathcal{P} \cap \mathcal{Q}$  and let  $\mathcal{A}^*$  denote the dual space of  $\mathcal{A}$ . The corresponding analytic subgroup  $A$  of  $\mathcal{A}$  in  $G$  is then called the vectorial part of  $X$ . For a  $\alpha \in \mathcal{A}^*$ , put

$$\mathcal{G}_\alpha = \{Y \in \mathcal{G} \mid [H, Y] = \alpha(H)Y, \forall H \in \mathcal{A}\}.$$

Then the set  $\Sigma = \{\alpha \in \mathcal{A}^* \mid \mathcal{G}_\alpha \neq \{0\}, \alpha \neq 0\}$  defines a root system with the inner product induced by the Killing form  $\langle, \rangle$  of  $\mathcal{G}$ . Moreover, the Weyl group  $W$  of  $\Sigma$  is defined with the normalizer  $N_K(\mathcal{A})$  of  $\mathcal{A}$  in  $K$  modulo the centralizer  $M = Z_K(\mathcal{A})$  of  $\mathcal{A}$  in  $K$ . It acts naturally on  $\mathcal{A}$  and coincides via this action with the reflection group of the root system  $\Sigma$ .

Choose a fundamental system  $\Delta = \{\alpha_1, \dots, \alpha_l\}$  of  $\Sigma$ , where the number  $l$  which equals  $\dim \mathcal{A}$  is called the split rank of the symmetric space  $X$  and denote  $\Sigma^+$  the corresponding set of all positive roots in  $\Sigma$ . Denote by  $W_{K \cap H}$  the normalizer  $N_{K \cap H}(\mathcal{A})$  of  $\mathcal{A}$  in  $K$  modulo the centralizer  $Z_{K \cap H}(\mathcal{A})$  of  $\mathcal{A}$  in  $K$ . We see that  $W_{K \cap H}$  is a subgroup of  $W$ . For each element  $\omega$  of  $W$  we fix a representative  $\underline{\omega}$  in  $N_K(\mathcal{A})$  so that  $\underline{\omega} \in N_{K \cap H}(\mathcal{A})$  if  $\omega \in W_{K \cap H}$ .

The purpose of this note is to construct a compact real analytic manifold  $\widehat{\mathbf{X}}$  in which the semisimple symmetric space  $\mathbf{X} = G/H$  is realized as an open subset and that  $G$  acts analytically on it. By the Cartan decomposition  $G = KAH$ , we must compactify the vectorial part  $A$ . Our construction is a motivation of Oshima's construction. In [5] we proposed a construction for the Riemannian symmetric spaces  $G/K$ . Here we apply the construction for the semisimple symmetric spaces  $\mathbf{X} = G/H$ .

Denote by  $\mathcal{G}_{\mathbf{C}}$  the complexification of  $\mathcal{G}$  and  $G_{\mathbf{C}}$  the corresponding analytic group. For simplicity, we assume that  $G$  is the real form of the complex Lie group  $G_{\mathbf{C}}$ . Let  $\mathcal{A}_{\mathbf{C}}$  be the complexification of  $\mathcal{A}$  and  $A_{\mathbf{C}}$  be the analytic subgroup of  $\mathcal{A}_{\mathbf{C}}$  in  $G_{\mathbf{C}}$ . For each  $a \in A_{\mathbf{C}}$  and  $\alpha \in \Sigma$  we define  $a^\alpha = e^{\alpha \cdot \log a} \in \mathbf{C}^* = \mathbf{C} \setminus \{0\}$  and consider the subset

$$A_{\mathbb{R}} = \{a \in A_{\mathbf{C}} \mid a^\alpha \in \mathbb{R}, \forall \alpha \in \Sigma\}.$$

This note is organized as follows. In Section 1, we consider the compactification of the vectorial part based on the construction in [5]. By this way, we firstly construct an embedding of  $A_{\mathbb{R}}$  into a compact real analytic manifold  $\widehat{A}_{\mathbb{R}}$  which is called a compactification of  $A_{\mathbb{R}}$ . Then we illustrate the construction via an example. In Section 2, we construct the compact manifold  $\widehat{\mathbf{X}}$  based on the action of Weyl group on  $\widehat{A}_{\mathbb{R}}$  and consider the real analytic structure of  $\widehat{\mathbf{X}}$  induced from the real analytic structure of  $\widehat{A}_{\mathbb{R}}$ .

## 2 Compactification of the vectorial part

Let  $(\mathbf{C}^*)^\Sigma$  be the set of complexes  $z = (z_\beta)_{\beta \in \Sigma}$ , where  $z_\beta \in \mathbf{C}^*$  and  $\mathbf{C}\mathbb{P}^1$  be the 1-dimensional complex projective space. Consider the map  $\varphi : A_{\mathbf{C}} \longrightarrow (\mathbf{C}^*)^\Sigma$  defined by  $\varphi(a) = (a^\alpha)_{\alpha \in \Sigma}$ ,  $\forall a \in A_{\mathbf{C}}$ . Then, for every  $z = (z_\alpha)_{\alpha \in \Sigma} \in \varphi(A_{\mathbf{C}})$  we have

$$z_{-\alpha} = (z_\alpha)^{-1}, \forall \alpha \in \Sigma \quad (2.1)$$

$$z_\alpha = \prod_{\gamma \in \Delta} (z_\gamma)^{k(\alpha, \gamma)}, \forall \alpha \in \Sigma^+, \alpha = \sum_{\gamma \in \Delta} k(\alpha, \gamma) \cdot \gamma. \quad (2.2)$$

By using the natural embedding of  $(\mathbf{C}^*)^\Sigma$  into  $(\mathbf{C}\mathbb{P}^1)^\Sigma$ , we get an embedding map of  $A_{\mathbf{C}}$  into  $(\mathbf{C}\mathbb{P}^1)^\Sigma$  denoted also by  $\varphi$ .

Let  $\mathbf{M} = \{z \in (\mathbb{R}\mathbb{P}^1)^\Sigma \mid z_{-\alpha} = z_\alpha^{-1}, \forall \alpha \in \Sigma\}$ , where  $\mathbb{R}\mathbb{P}^1$  is the 1-dimensional real projective space. By definition,  $\mathbf{M}$  is compact. Moreover, the subset

$$\mathcal{U}_{\Sigma^+} = \left\{ m = (m_\alpha, m_{-\alpha}) \in \mathbf{M} \mid m_\alpha \in \mathbb{R}, m_{-\alpha} \in \mathbb{R}^* \cup \{\infty\}, \forall \alpha \in \Sigma^+ \right\}$$

is an open subset in  $(\mathbb{R}\mathbb{P}^1)^\Sigma$  and we get a homeomorphism  $\chi_{\Sigma^+} : \mathcal{U}_{\Sigma^+} \longrightarrow \mathbb{R}^{\Sigma^+}$  defined by  $\chi_{\Sigma^+}(m) = (m_\alpha)_{\alpha \in \Sigma^+}$ ,  $\forall m \in \mathcal{U}_{\Sigma^+}$ .

Recall that  $W$  acts on  $\mathbf{M}$  by  $(w.z)_\alpha = z_{w^{-1}\alpha}$ ,  $\forall \alpha \in \Sigma$ ,  $w \in W$ ,  $z \in \mathbf{M}$ . So we have  $\mathcal{U}_{w(\Sigma^+)} = w(\mathcal{U}_{\Sigma^+})$ ,  $\forall w \in W$ . By a similar way as in [5, Lemma 1.2], we see that the pair  $\{\mathcal{U}_{\Sigma^+}, \chi_{\Sigma^+}\}$  is a chart on  $\mathbf{M}$  and  $\{\mathcal{U}_{w(\Sigma^+)}, \chi_{w(\Sigma^+)}\}_{w \in W}$  defines an atlas of charts on  $\mathbf{M}$  such that  $\mathbf{M}$  becomes a real analytic submanifold.

By definition,  $\varphi(A_{\mathbb{R}})$  is a subset of  $(\mathbb{R}\mathbb{P}^1)^\Sigma$ . Denote by  $\widehat{A}_{\mathbb{R}}$  the closure of  $\varphi(A_{\mathbb{R}})$  in  $(\mathbb{R}\mathbb{P}^1)^\Sigma$ . It follows from (2.1) and (2.2) that  $\widehat{A}_{\mathbb{R}}$  is a compact subset of  $\mathbf{M}$ . We now define an atlas of charts on  $\widehat{A}_{\mathbb{R}}$  induced from the atlas on  $\mathbf{M}$ .

Let  $\mathcal{U}_\Delta$  be the subset of  $\mathcal{U}_{\Sigma^+}$  consists of elements  $m = (m_\alpha, m_{-\alpha})$  such that  $m_\alpha = \prod_{\gamma \in \Delta} (m_\gamma)^{k(\alpha, \gamma)}$ ,  $\forall \alpha \in \Sigma^+$ ,  $\alpha = \sum_{\gamma \in \Delta} k(\alpha, \gamma) \cdot \gamma$ . Then  $\mathcal{U}_\Delta$  is an open subset in  $\widehat{A}_{\mathbb{R}}$ . It follows that  $\chi_{\Sigma^+}(\mathcal{U}_\Delta) = \{x \in \mathbb{R}^{\Sigma^+} \mid x_\alpha = \prod_{\gamma \in \Delta} (x_\gamma)^{k(\alpha, \gamma)}\}$  and we get

a homeomorphism  $\chi_\Delta : \mathcal{U}_\Delta \longrightarrow \mathbb{R}^\Delta$  defined by  $\chi_\Delta(m) = (m_\gamma)_{\gamma \in \Delta}$ ,  $\forall m \in \mathcal{U}_\Delta$ . Moreover, by a similar argument as that given in [5, Theorem 1.4], we have

**Theorem 2.1.**  *$\widehat{A}_{\mathbb{R}}$  is a compact real analytic manifold that is called a compactification of  $A_{\mathbb{R}}$ . The set of charts  $\{\mathcal{U}_{w(\Delta)}, \chi_{w(\Delta)}\}_{w \in W}$  defines an atlas of charts on  $\widehat{A}_{\mathbb{R}}$  so that the manifold  $\widehat{A}_{\mathbb{R}}$  is covered by  $|W|$ -many charts.*

**Remark 2.2.** Denote by  $\mathcal{G}_+ = \mathcal{G}^{\sigma\theta} = \mathcal{K} \cap \mathcal{H} \oplus \mathcal{P} \cap \mathcal{Q}$  and let

$$\Sigma_+ = \Sigma(\mathcal{A}, \mathcal{G}_+) = \{\alpha \in \Sigma \mid \mathcal{G}_\alpha \cap \mathcal{G}_+ \neq \{0\}, \alpha \neq 0\}$$

be the set of restricted roots of  $\mathcal{A}$  in  $\mathcal{G}_+$ .

Suppose that  $H$  is connected. Since  $H = (H \cap K).exp(\mathcal{H} \cap \mathcal{P})$  we have that  $H \cap K$  is connected. Then  $G_+ = (H \cap K).exp(\mathcal{P} \cap \mathcal{Q})$  is connected and the corresponding Weyl group  $W_{K \cap H}$  of  $\mathcal{A}$  in  $\mathcal{G}_+$  defined by

$$W_{K \cap H} = W_{(K \cap H)_e} = W(\mathcal{A}, \mathcal{G}_+) = \langle s_\alpha \mid \alpha \in \Sigma_+ \rangle.$$

Since  $\mathcal{G}_+ = \mathcal{K} \cap \mathcal{H} \oplus \mathcal{P} \cap \mathcal{Q}$  is the Cartan decomposition and  $K \cap H$  is a maximal compact subgroup of  $G_+$ , the coset  $\mathbf{X}_+ := G_+ / (K \cap H)$  becomes a Riemannian symmetric space of non-compact type. We then can apply the construction for the vectorial part of  $\mathbf{X}_+$ .

Denote  $A_{\mathbb{R},+} = \left\{ a \in A_{\mathbb{C}} \mid a^\gamma \in \mathbb{R}, \forall \gamma \in \Sigma_+ \right\}$ . By construction, the corresponding compactification  $\widehat{A}_{\mathbb{R},+}$  is contained in the compact manifold

$$\mathbf{M}_+ = \left\{ z \in (\mathbb{P}^1(\mathbb{R}))^{\Sigma_+} \mid z_{-\gamma} = (z_\gamma)^{-1}, \forall \gamma \in \Sigma_+ \right\}.$$

Moreover,  $\widehat{A}_{\mathbb{R},+}$  is a compact real analytic manifold which is covered by  $|W_{K \cap H}|$ -many charts. Here, each Weyl chamber of  $\mathcal{A}$  for  $\Sigma_+$  contains  $|W_{K \cap H} \setminus W|$ -corresponding Weyl chambers of  $\mathcal{A}$  for  $\Sigma$  and these subchambers are parameterized by  $W_{K \cap H} \setminus W$ .

**Example** Consider the real semi-simple Lie group  $G = SL(3, \mathbb{R})$  and denote by  $\mathcal{G} = sl(3, \mathbb{R})$  the corresponding Lie algebra of  $G$ . Suppose that  $\theta$  is the Cartan involution defined by  $\theta(X) = ({}^t X)^{-1}$ ,  $\forall X \in \mathcal{G}$  and  $K = SO(3, \mathbb{R})$  is the maximal compact subgroup in  $G$  with respect to  $\theta$ . Then  $\mathcal{G} = \mathcal{K} \oplus \mathcal{P}$  is the Cartan decomposition of  $\mathcal{G}$  with respect to  $\theta$ , where  $\mathcal{K} = so(3, \mathbb{R})$  is the Lie algebra of  $K$ .

Let  $\sigma$  be the involution of  $\mathcal{G}$  defined by

$$\sigma(X) = J\theta(X)J, \forall X \in \mathcal{G}, \text{ where } J = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

and  $\mathcal{G} = \mathcal{H} \oplus \mathcal{Q}$  be the decomposition of  $\mathcal{G}$  with respect to  $\sigma$ , where  $\mathcal{H} = so(1, 2)$  is the Lie algebra of the corresponding fixed points group  $H = SO_0(1, 2)$ .

It follows that  $\mathbf{X} = G/H = SL(3, \mathbb{R})/SO_0(1, 2)$  is a semisimple symmetric space of rank two and we get a maximal Abelian subspace of  $\mathcal{P} \cap \mathcal{Q}$  defined by

$$\mathcal{A} = \left\{ \begin{pmatrix} t_1 & 0 & 0 \\ 0 & t_2 & 0 \\ 0 & 0 & t_3 \end{pmatrix} \mid t_1 + t_2 + t_3 = 0 \right\}.$$

By definition, the root system  $\Sigma$  of  $\mathcal{A}$  in  $\mathcal{G}$  is  $\Sigma = \{ e_i - e_j \mid 1 \leq i \neq j \leq 3 \}$  and the Weyl group  $W$  is isomorphic to  $S_3$ , the symmetric group of order three.

Moreover, the corresponding analytic subgroup in  $G$  of  $\mathcal{A}$  defined by

$$A = \left\{ \begin{pmatrix} a_1 & 0 & 0 \\ 0 & a_2 & 0 \\ 0 & 0 & a_3 \end{pmatrix} \mid a_1 a_2 a_3 = 1, a_i > 0 \right\} \cong (0, \infty)^2$$

and we get

$$A_{\mathbb{R}} = \left\{ \begin{pmatrix} a_1 & 0 & 0 \\ 0 & a_2 & 0 \\ 0 & 0 & a_3 \end{pmatrix} \mid a_1 a_2 a_3 = 1 \right\} \cong (\mathbb{R}^*)^2.$$

Denote  $\alpha = e_1 - e_2$ ,  $\beta = e_2 - e_3$ . We see that  $\Delta = \{\alpha, \beta\}$  and  $\Sigma = \Sigma^+ \cup (-\Sigma^+)$ , where  $\Sigma^+ = \{\alpha, \beta, \alpha + \beta\}$ . Then we have

$$\mathbf{M} = \left\{ (z_\gamma, z_{-\gamma}) \mid z_\gamma \in \mathbb{P}^1(\mathbb{R}), \gamma \in \Sigma^+ \right\} \cong \mathbb{P}^1(\mathbb{R})^3.$$

It follows that the pair  $\{\mathcal{U}_{\Sigma^+}, \chi_{\Sigma^+}\}$  is a chart on  $\mathbf{M}$ , where

$$\mathcal{U}_{\Sigma^+} = \left\{ (m_\alpha, m_\beta, m_{\alpha+\beta}, m_{-\alpha}, m_{-\beta}, m_{-\alpha-\beta}) \mid m_\alpha, m_\beta, m_{\alpha+\beta} \in \mathbb{R} \right\} \cong \mathbb{R}^3$$

is an open subset in  $(\mathbb{R}\mathbb{P}^1)^{\Sigma^+}$  and  $\chi_{\Sigma^+} : \mathcal{U}_{\Sigma^+} \rightarrow \mathbb{R}^{\Sigma^+}$  is the homeomorphism defined by  $\chi_{\Sigma^+}(m) = (m_\alpha, m_\beta, m_{\alpha+\beta})$ ,  $\forall m \in \mathcal{U}_{\Sigma^+}$ . Then we get

$$\mathcal{U}_\Delta = \left\{ (m_\alpha, m_\beta, m_{\alpha+\beta}, m_{-\alpha}, m_{-\beta}, m_{-\alpha-\beta}) \mid m_\alpha, m_\beta, m_{\alpha+\beta} \in \mathbb{R}, m_{\alpha+\beta} = m_\alpha \cdot m_\beta \right\}$$

and  $\chi_{\Sigma^+}(\mathcal{U}_\Delta) = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_3 = x_1 \cdot x_2\} \cong \mathbb{R}^2$ . Hence  $\widehat{A}_{\mathbb{R}} \cong \mathbb{R}^2 \cup \{\infty\} \cong S^2$  is a compact smooth manifold that is covered by 6-many charts.

By definition, we see that

$$G_+ = \left\{ \begin{pmatrix} a & 0 & 0 \\ 0 & a_{11} & a_{12} \\ 0 & a_{21} & a_{22} \end{pmatrix} \mid \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \in GL(2, \mathbb{R}), a^{-1} = \det(A) > 0 \right\}.$$

Then  $X_+ = G_+/K \cap H$  is a Riemannian symmetric space of non-compact type.

Moreover, we have  $\Sigma_+ = \{e_i - e_j \mid 2 \leq i \neq j \leq 3\} = \{\beta, -\beta\}$  and the Weyl group  $W_{K \cap H}$  is isomorphic to  $S_2$ , where  $S_2$  is the symmetric group of order two. It follows that the coset  $W_{K \cap H} \setminus W$  has three elements. Note that

$$A_{\mathbb{R},+} = \left\{ a \in A_{\mathbb{C}} \mid a^\gamma \in \mathbb{R}, \forall \gamma \in \Sigma_+ \right\} \cong \mathbb{R}^*.$$

Then the corresponding compactification  $\widehat{A}_{\mathbb{R},+}$  is contained in the compact manifold

$$\mathbf{M}_+ = \left\{ (z_\gamma, z_\gamma^{-1}) \mid z_\gamma \in \mathbb{P}^1(\mathbb{R}), \forall \gamma \in \Sigma_+ \right\} \cong \mathbb{P}^1(\mathbb{R}).$$

It follows that  $\widehat{A}_{\mathbb{R},+} \cong \mathbb{R} \cup \{\infty\} \cong S^1$  is a compact smooth manifold that is covered by 2-many charts.

Here each Weyl chamber of  $\mathcal{A}$  for  $\Sigma_+$  contains three corresponding Weyl chambers of  $\mathcal{A}$  for  $\Sigma$  and these subchambers are parameterized by  $W_{K \cap H} \setminus W$ .

### 3 Construction of a compact embedding

In this Section, we will construct an embedding of the semisimple symmetric space  $\mathbf{X}$  into a compact real analytic manifold  $\widehat{\mathbf{X}}$  such that the action of  $G$  on  $\widehat{\mathbf{X}}$  is analytic. Our construction is based on the indicated construction in [5] for Riemannian symmetric spaces and similar to those of Kosters, Sekiguchi, Oshima for semisimple symmetric spaces.

First we recall some notations concerning the compactification  $\widehat{A}_{\mathbb{R}}$  as in [5]. Consider the subset  $\widehat{A}_{\mathbb{R}}^- = \{ \tilde{a} \in \widehat{A}_{\mathbb{R}} \mid (\tilde{a})^\alpha \in [-1, 1] \}$  and recall that the Weyl group  $W$  acts on  $\widehat{A}_{\mathbb{R}}$  as follows  $(w.\tilde{a})_\alpha = (\tilde{a})_{w^{-1}(\alpha)}$ ,  $\forall w \in W, \forall \tilde{a} \in \widehat{A}_{\mathbb{R}}$ . Then we have  $W.\widehat{A}_{\mathbb{R}}^- = \widehat{A}_{\mathbb{R}}$  (see [5, Lemma 2.1]). Moreover, for each element  $\tilde{a} \in \widehat{A}_{\mathbb{R}}$  we have a unique decomposition  $\tilde{a} = (\tilde{a})_{fin} \cdot \epsilon(\tilde{a})$ , where  $a_{fin} \in A_{\mathbb{R}}$  and  $\epsilon(\tilde{a}) \in \widehat{A}_{\mathbb{R}}$  such that  $\epsilon(\tilde{a})^\gamma \in \{-1, 0, +1, \infty\}$ ,  $\forall \gamma \in \Delta$ .

Note that  $\epsilon(\tilde{a}) \in \{-1, 0, +1, \infty\}^\Delta$  and for all  $\alpha = \sum_{\gamma \in \Delta} k(\alpha, \gamma) \cdot \gamma \in \Sigma$  we have

$$\epsilon(\tilde{a})^\alpha = \prod_{\gamma \in \Delta} (\epsilon(\tilde{a})^\gamma)^{|k(\alpha, \gamma)|}.$$

Motivating the Oshima's definition,  $\epsilon(\tilde{a})$  is called an extended signature of the element  $\tilde{a}$ . Now we define parabolic subalgebras with respect to extended signatures  $\epsilon(\tilde{a})$ , for all  $\tilde{a} \in \widehat{A}_{\mathbb{R}}$ .

First we consider  $\tilde{a}_t \in \widehat{A}_{\mathbb{R}}^-$  and let  $F_\epsilon = \{ \gamma \mid \epsilon_\gamma = \epsilon(\tilde{a}_t)^\gamma \neq 0 \}$  be a subset of the simple root system  $\Delta$  with respect to the extended signature  $\epsilon = \epsilon(\tilde{a}_t)$ . Denote  $\Sigma_\epsilon = (\sum_{\gamma \in F_\epsilon} \mathbb{R}\gamma) \cap \Sigma$  and suppose that  $W_\epsilon$  is the subgroup of  $W$  generated by reflections with respect to  $\gamma$  in  $F_\epsilon$ . Let  $P_\sigma$  be the parabolic subgroup of  $G$  with the corresponding Langlands decomposition  $P_\sigma = M_\sigma A_\sigma N_\sigma$  so that  $M_\sigma A_\sigma$  is the centralizer of  $\mathcal{A}$  in  $G$  and the Lie algebra  $\mathcal{N}_\sigma$  of  $N_\sigma$  equals  $\sum_{\alpha \in \Sigma^+} \mathcal{G}_\alpha$ . Then

we can define a parabolic subalgebra

$$\mathcal{P}_\epsilon = \mathcal{M}_\sigma + \mathcal{A}_\sigma + \sum_{\alpha \in \Sigma_\epsilon} \mathcal{G}_\alpha + \sum_{\alpha \in \Sigma^+ - \Sigma_\epsilon} \mathcal{G}_\alpha$$

of  $\mathcal{G}$  and its Langlands decomposition  $\mathcal{P}_\epsilon = \mathcal{M}_\epsilon + \mathcal{A}_\epsilon + \mathcal{N}_\epsilon$  so that  $\mathcal{A}_\epsilon \subset \mathcal{A}_\sigma$  (see [11]).

Let  $P_\epsilon$  denote the corresponding parabolic subgroup of  $\mathcal{P}_\epsilon$  in  $G$ . It follows that  $P_\epsilon = M_\epsilon A_\epsilon N_\epsilon$  is the corresponding Langlands decomposition of  $P_\epsilon$  and we

define a closed subgroup  $P(\epsilon)$  of  $G$  by  $P(\epsilon) = (M_\epsilon \cap \underline{\omega}^{-1}H\underline{\omega})A_\epsilon N_\epsilon$ , where  $\underline{\omega}$  is a representative of  $\omega \in W$  in  $N_K(\mathcal{A})$  so that  $\underline{\omega} \in N_{K \cap H}(\mathcal{A})$  if  $\omega \in W_{K \cap H}$ .

In general, for each  $\eta = \eta(\tilde{a})$  with  $\tilde{a} = w.\tilde{a}_t \in \widehat{A}_\mathbb{R}$ , we firstly consider the parabolic subalgebra  $P_\epsilon$  with respect to the element  $\epsilon = \epsilon(\tilde{a}_t)$ . Then we define a parabolic subalgebra  $P_\eta = w.P_\epsilon.w^{-1}$  based on the action of the Weyl group  $W$  on the parabolic subalgebra  $P_\epsilon$  (see [3]).

We now define an equivalent relation on the product manifold  $G \times \widehat{A}_\mathbb{R}$ . Let  $x = (g, \tilde{a})$  be an element of  $G \times \widehat{A}_\mathbb{R}$ , where  $\tilde{a} = w.\tilde{a}_t$  and  $t = (t_\gamma)_{\gamma \in \Delta}$ ,  $t_\gamma \in [-1, 1]$ . Then we put  $\text{sgn } x = \epsilon(\tilde{a}_t) = \text{sgn } t$ , which is an element of  $\{-1, 0, 1\}^\Delta$ . Here  $\text{sgn } t = (\text{sgn } t_\gamma)_{\gamma \in \Delta}$  and for an  $s$  in  $\mathbb{R}$  we define  $\text{sgn } s = 1$  (resp.  $0, -1$ ) if  $s > 0$  (resp.  $s = 0, s < 0$ ).

Denote  $F_x, \Sigma_x$  and  $W_x$  instead of  $F_\epsilon, \Sigma_\epsilon$  and  $W_\epsilon$ , respectively, we define a parabolic subalgebra

$$\mathcal{P}_x = \mathcal{M}_\sigma + \mathcal{A}_\sigma + \sum_{\alpha \in \Sigma_x} \mathcal{G}_\alpha + \sum_{\alpha \in \Sigma^+ - \Sigma_x} \mathcal{G}_\alpha$$

of  $\mathcal{G}$  and its Langlands decomposition  $\mathcal{P}_x = \mathcal{M}_x + \mathcal{A}_x + \mathcal{N}_x$  so that  $\mathcal{A}_\epsilon \subset \mathcal{A}_\sigma$ .

Moreover, denoting by  $P_x$  the corresponding subgroup of  $\mathcal{P}_x$  in  $G$ , we get the Langlands decomposition  $P_x = M_x A_x N_x$  and  $P(x) = (M_x \cap \underline{\omega}^{-1}H\underline{\omega})A_x N_x$  is a closed subgroup of  $G$ . Let  $\{H_1, H_2, \dots, H_l\}$  denote the dual basis of  $\Delta = \{\alpha_1, \dots, \alpha_l\}$ , that is,  $H_j \in \mathcal{A}$  and  $\alpha_i(H_j) = \delta_{ij}$ ,  $\forall i, j = 1, 2, \dots, l$  and put  $a(x) = \exp(-\sum_{\gamma \in F_x} \log|t_\gamma| H_\gamma)$ , where  $H_\gamma \in \{H_1, H_2, \dots, H_l\}$  with respect to  $\gamma$ .

Note that for all elements  $x = (g, \omega.\tilde{a}_t)$  and  $x' = (g', \omega'.\tilde{a}_{t'})$  of  $G \times \widehat{A}_\mathbb{R}$  such that  $\text{sgn } x = \text{sgn } x'$ , we have  $W_x = W_{x'}$ .

**Definition 3.1.** We say that two elements  $x = (g, \omega.\tilde{a}_t)$  and  $x' = (g', \omega'.\tilde{a}_{t'})$  of  $G \times \widehat{A}_\mathbb{R}$  are equivalent if and only if the following conditions hold:

- (i)  $\text{sgn } x = \text{sgn } x'$
- (ii)  $W_{K \cap H} \omega W_x = W_{K \cap H} \omega' W_x$
- (iii)  $ga(x)P(x) = g'a'(x')(M_x \cap \underline{\omega}'^{-1}H\underline{\omega}')A_x N_x$ .

**Remark 3.2.** Let  $x = (g, \omega.\tilde{a}_t)$  and  $x' = (g', \omega'.\tilde{a}_{t'})$  be elements of  $G \times \widehat{A}_\mathbb{R}$  and suppose that  $x$  and  $x'$  satisfy the above condition (iii). Then there exist  $u, u' \in W_{K \cap H}$  and  $v, v' \in W_x$  and  $m, m' \in M$  such that  $\underline{u} \underline{\omega} \underline{v} m = \underline{u}' \underline{\omega}' \underline{v}' m'$ . Moreover, (see [11, Lemma 1.2]) the condition (iii) in Definition 2.1 is equivalent to

$$ga(x)P(x)\underline{v}m = g'a'(x')P(x')\underline{v}'m'. \quad (3.1)$$

It follows from Remark 3.2 that the Definition 3.1 really gives an equivalent relation, and we write  $x \sim x'$ . The quotient space of  $G \times \widehat{A}_\mathbb{R}$  by this equivalent relation becomes a topological space with the quotient topology and is denoted by  $\widehat{\mathbf{X}}$ .

Let  $\pi : G \times \widehat{A}_{\mathbb{R}} \longrightarrow \widehat{\mathbf{X}}$  be the natural projection. Because the action of  $G$  on  $G \times \widehat{A}_{\mathbb{R}}$  are compatible with the equivalent relation, we can define an action of  $G$  on  $\widehat{\mathbf{X}}$  by

$$g_1 \pi(g, \tilde{a}) = \pi(g_1 g, \tilde{a}), \quad \forall g, g_1 \in G, \tilde{a} \in A_{\mathbb{R}}. \quad (3.2)$$

Put  $\widehat{A}_{\mathbb{R}, \epsilon} = \{ \tilde{a} \in \widehat{A}_{\mathbb{R}} \mid \epsilon(\tilde{a}) = \epsilon \}$  for each  $\epsilon \in \{ -1, 0, 1 \}^{\Delta}$  and denote  $\mathbf{X}_{\epsilon} = \pi(G \times \widehat{A}_{\mathbb{R}, \epsilon})$ . Then we have

**Proposition 3.3.** (i)  $\widehat{\mathbf{X}}$  is a compact connected  $G$ -space and  $\widehat{\mathbf{X}} = \bigcup_{\epsilon \in \{ -1, 0, 1 \}^{\Delta}} \mathbf{X}_{\epsilon}$

gives the orbital decomposition of  $\widehat{\mathbf{X}}$  for the action of  $G$  on it.

(ii) Each  $\mathbf{X}_{\epsilon}$  is homeomorphic to  $G/P(\epsilon)$ . There are just  $2^{|\Delta|}$  open orbits that are isomorphic to  $G/H$  and the number of compact orbits in  $\widehat{\mathbf{X}}$  equal that of the elements of the coset  $W_{K \cap H} \setminus W$ .

**Proof** (i) Since  $\pi(G \times \mathcal{U}_{\Delta})$  is connected and contains any open orbit, the connectedness of  $\widehat{\mathbf{X}}$  is clear. Denote by  $\mathcal{A}^+$  the positive chamber corresponding to  $\Sigma^+$  and put  $A^+ = \exp \mathcal{A}^+$ . Let  $\overline{A^+} = \{ \exp X \mid X \in \mathcal{A} \text{ with } \alpha(X) \geq 0 \text{ for all } \alpha \in \Sigma^+ \}$  be the closure of  $A^+$ , we see that

$$\overline{A^+} = \left\{ \exp \left( - \sum_{\gamma} (\log t_{\gamma}) H_{\gamma} \right) \mid t_{\gamma} \in (0, 1] \right\}.$$

Consider the compact subset  $K \times \widehat{A}_{\mathbb{R}} = K \times W \cdot \widehat{A}_{\mathbb{R}} \cong K \times [-1, 1]^{\Delta} \times W$  of  $G \times \widehat{A}_{\mathbb{R}}$ . Then the subset  $\pi(K \times \widehat{A}_{\mathbb{R}})$  is also compact because it is the image of a compact set under the continuous map. Moreover, it follows from the Cartan decomposition  $G = KAH$  that  $G = \bigcup_{w \in W} K \overline{A^+} w H$  (see [11]). Hence, Definition

3.1 and Remark 3.2 prove that the compact set  $\pi(K \times \widehat{A}_{\mathbb{R}})$  contains all open  $G$ -orbits in  $\widehat{\mathbf{X}}$ . In other words, this compact set is dense in  $\widehat{\mathbf{X}}$  and therefore it must be coincided with  $\widehat{\mathbf{X}}$ .

(ii) Put  $\tilde{a} \in \widehat{A}_{\mathbb{R}, \epsilon}$  for each  $\epsilon \in \{ -1, 0, 1 \}^{\Delta}$  and define a map  $\Psi : G/P(\epsilon) \longrightarrow \mathbf{X}_{\epsilon}$  by  $\Psi(gP(\epsilon)) = \pi(g, \tilde{a})$ ,  $\forall g \in G$ . Then by a similar way as in [5, Proposition 2.4], we get that the map is well defined and becomes an homeomorphism which is equivariant for the action of  $G$ .

We now define an analytic structure on  $\widehat{\mathbf{X}}$  based on the analytic structure on  $\widehat{A}_{\mathbb{R}}$ .

Let  $\mathcal{A}_{\mathcal{P}}$  be a maximal Abelian subspace of  $\mathcal{P}$  containing  $\mathcal{A}$  and let  $\Sigma(\mathcal{A}_{\mathcal{P}}, \mathcal{G})$  be the set of corresponding restricted roots. By [11, Lemma 1.4], we can assume that the representatives  $\underline{w}$  of elements  $w \in W$  satisfy  $Ad(\underline{w})(\mathcal{A}_{\mathcal{P}}) = \mathcal{A}_{\mathcal{P}}$ . Denote by  $\mathcal{G}(\sigma)$  the reductive Lie algebra generated by  $\{ \mathcal{G}(\mathcal{A}_{\mathcal{P}}; \lambda) \mid \lambda \in \Sigma(\mathcal{A}_{\mathcal{P}}, \mathcal{G}), \lambda|_{\mathcal{A}} = 0 \}$ , where  $\mathcal{G}(\mathcal{A}_{\mathcal{P}}; \lambda) = \{ X \in \mathcal{G} \mid [X, Y] = \lambda(Y)X \text{ for all } Y \in \mathcal{A}_{\mathcal{P}} \}$  and put

$$\mathcal{M}(\sigma) = \{ X \in \mathcal{M}_{\sigma} \mid [X, Y] = \lambda(Y)X \text{ for all } Y \in \mathcal{G}(\sigma) \}.$$



Let  $G(\sigma)$  and  $M(\sigma)_0$  be the analytic subgroups of  $G$  corresponding to  $\mathcal{G}(\sigma)$  and  $\mathcal{M}(\sigma)$ , respectively, and denote

$$M(\sigma) = M(\sigma)_0 \cdot Ad_G^{-1}(Ad(K) \cap \exp(\text{ad}(\sqrt{-1}\mathcal{A}_p))).$$

Then  $G(\sigma) \subset H$ ,  $M(\sigma) \subset N_K(\mathcal{A}_p)$  and the representative  $\underline{\omega}$  normalizes  $G(\sigma)$  and  $M(\sigma)$  for any  $\omega \in W$ . Moreover, it follows from [11, Lemma 1.5] that  $M_\sigma = M(\sigma)G(\sigma)$  and

$$M(\sigma)/(M(\sigma) \cap \underline{\omega}^{-1}H\underline{\omega}) \simeq M_\sigma/(M_\sigma \cap \underline{\omega}^{-1}H\underline{\omega}) \simeq M/(M \cap \underline{\omega}^{-1}H\underline{\omega}).$$

For every  $g \in G$  and  $w \in W$ , we put  $\Omega_g^w = \pi(g\overline{N}_\sigma M(\sigma) \times \mathcal{U}_{w(\Delta)})$ , where  $\overline{N}_\sigma$  is the analytic subgroup of  $G$  corresponding to  $\overline{N}_\sigma = \theta(\mathcal{N}_\sigma)$  and define a map

$$\Phi_g^w : \overline{N}_\sigma \times M(\sigma)/(M(\sigma) \cap \underline{\omega}^{-1}H\underline{\omega}) \times \mathbb{R}^\Delta \longrightarrow \Omega_g^w$$

by  $\Phi_g^w(n, m, t) = \pi(gnm, w.\tilde{a}_t)$ ,  $\forall (n, m, t) \in \overline{N}_\sigma \times M(\sigma)/(M(\sigma) \cap \underline{\omega}^{-1}H\underline{\omega}) \times \mathbb{R}^\Delta$ .  $\square$  By the same argument as that given in [5, Lemma 2.6], we get

**Lemma 3.4.** *For every  $g \in G$  and  $w \in W$ , the map  $\Phi_g^w$  is a homeomorphism of  $\overline{N}_\sigma \times M(\sigma)/(M(\sigma) \cap \underline{\omega}^{-1}H\underline{\omega}) \times \mathbb{R}^\Delta$  onto an open subset  $\Omega_g^w = \pi(g\overline{N}_\sigma M(\sigma) \times \mathcal{U}_{w(\Delta)})$  of  $\widehat{X}$ .*

For brevity, we denote  $\Omega^w = \overline{N}_\sigma \times M(\sigma)/(M(\sigma) \cap \underline{\omega}^{-1}H\underline{\omega}) \times \mathbb{R}^\Delta$ . Then we have

**Lemma 3.5.** *Let  $g, g' \in G$  and  $w, w' \in W$ . Then the map*

$$(\Phi_{g'}^{w'})^{-1} \circ (\Phi_g^w) : (\Phi_g^w)^{-1}(\Omega_g^w \cap \Omega_{g'}^{w'}) \longrightarrow (\Phi_{g'}^{w'})^{-1}(\Omega_g^w \cap \Omega_{g'}^{w'})$$

*define an analytic diffeomorphism between the open subsets of the set  $\Omega^w$ .*

**Proof** By definition,  $\Phi_g^w$  is bijective and continuous. Then the map  $(\Phi_{g'}^{w'})^{-1} \circ (\Phi_g^w)$  is bijective and its inverse is of the same form. So we need only to show that the map  $(\Phi_{g'}^{w'})^{-1} \circ (\Phi_g^w)$  is analytic. Since  $(\Phi_{g'}^{w'})^{-1} \circ (\Phi_g^w) = (\Phi_e^{w'})^{-1} \circ (\Phi_{g'^{-1}g}^w)$ , we can assume that  $g' = e$ .

Fix an arbitrary element  $u = (n_o, m_o, t_o)$  of the domain of  $(\Phi_e^{w'})^{-1} \circ (\Phi_g^w)$  and put  $x = (gn_o m_o, w\tilde{a}_{t_o}) \in \widehat{X}$  and  $u' = (n'_o, m'_o, t'_o) = ((\Phi_e^{w'})^{-1} \circ \Phi_g^w)(u)$  in  $\Omega^{w'}$ , we will show that the map is analytic in a neighborhood of  $u$ .

We first consider the case where  $w' = w$  and  $g \in \overline{N}_\sigma M(\sigma)A^x$ . Here  $A^x$  is the analytic subgroup of  $G$  corresponding to  $A^x = \sum_{\gamma \in F_x} \mathbb{R}H_\gamma$ . Suppose that  $g = n_1 m_1 a_1$ , where  $n_1 \in \overline{N}_\sigma, m_1 \in M(\sigma)$  and  $a_1 \in A^x$ . Then we have

$$(\Phi_e^w)^{-1} \circ (\Phi_g^w)(n, m, t) = (n_1 m_1 a_1 n(m_1 a_1))^{-1}, m_1 m, a_1 t,$$

where  $a_1 t = (\exp \langle -a_1, \log a_1 \rangle t_1, \dots, \exp \langle -a_l, \log a_1 \rangle t_l)$ . It follows that the map is analytic.

Next we assume that  $w' = w$  and  $u' = u = (e, 1, \epsilon)$  with an  $\epsilon \in \{-1, 0, 1\}^\Delta$ . Then we have  $g \in P(x)$  and [11, Lemma 1.9 (i)] there exist neighborhoods  $V$  of the origin in  $\mathcal{P}(x)$  and  $U_0$  of  $u$  in  $\Omega^w$  such that for any  $Y \in V$  and  $s \in [0, 1]$ , the map  $(\Phi_e^w)^{-1} \circ \exp(sY) \circ \Phi_e^w$  defines an analytic diffeomorphism of  $U_0$  onto a neighborhood of  $u$ . Hence we have the claim if  $g \in \exp V$ . Moreover, any  $g \in P(x)$  can be written in the form  $g = g_0 g_1 \dots g_k$  with  $g_0 \in M(\sigma) \cap \underline{\omega}^{-1} H \underline{\omega}$  and  $g_i \in \exp V (i = 1, \dots, k)$ , where  $k$  is a suitable positive integer. Then we have the relation

$$(\Phi_e^w)^{-1} \circ \Phi_g^w = ((\Phi_e^w)^{-1} \circ \Phi_{g_0}^w) \circ ((\Phi_e^w)^{-1} \circ \Phi_{g_1}^w) \circ \dots \circ ((\Phi_e^w)^{-1} \circ \Phi_{g_k}^w)$$

and  $(\Phi_e^w)^{-1} \circ \Phi_{g_i}^w$  map the point  $u$  to the same point for  $i = 0, 1, \dots, k$ . It follows that  $(\Phi_e^w)^{-1} \circ \Phi_g^w$  are analytic in some neighborhoods of  $u$  in  $\Omega^w$  and we have the claim.

We consider the case where  $w' \neq w, g = e$  and  $u = (e, 1, \epsilon)$ . Then it follows that  $u' = (e, 1, \epsilon)$  when  $g' = \underline{v}' m' m^{-1} \underline{v}^{-1}$ , with  $v, v' \in W_x$ . By a similar way as in [11, Lemma 1.9 (ii)], we can prove that the map  $\Phi_{g'}^{w'} \circ (\Phi_e^w)$  is analytic in the set

$$\Omega^w(\epsilon) = \overline{N}_\sigma \times M(\sigma) / (M(\sigma) \cap \underline{\omega}^{-1} H \underline{\omega}) \times \mathbb{R}_\epsilon^\Delta,$$

where  $\mathbb{R}_\epsilon^\Delta = \{t \in \mathbb{R}^\Delta \mid \text{sgn } t_\gamma = \epsilon_\gamma \text{ if } \epsilon_\gamma \neq 0\}$ .

Now we consider the general case. Put  $g_1 = (n_o \underline{m}_o a(t_o))^{-1}$ ,  $g_2 = g_3^{-1} g_4^{-1} g g_1^{-1}$ , where  $g_3 = \underline{v}' m' m^{-1} \underline{v}^{-1}$ ,  $g_4 = n'_o \underline{m}_o a(t'_o)$  and consider the maps

$$\Phi_1 = (\Phi_e^w)^{-1} \circ \Phi_{g_1}^w, \Phi_2 = (\Phi_e^w)^{-1} \circ \Phi_{g_2}^w, \Phi_3 = (\Phi_{g_3}^{w'})^{-1} \circ (\Phi_e^w), \Phi_4 = (\Phi_e^{w'})^{-1} \circ \Phi_{g_4}^{w'}.$$

Then we have

$$(\Phi_e^{w'})^{-1} \circ \Phi_g^w = \Phi_4 \circ \Phi_3 \circ \Phi_2 \circ \Phi_1.$$

Since  $\Phi_1(u) = (e, 1, \text{sgn } x)$  and  $g_2 \in P(x)$ , it follows from what we have proved that the maps  $\Phi_1, \Phi_2, \Phi_3$  and  $\Phi_4$  are analytic in a neighborhood of  $u = (n_o, m_o, t_o)$ . This implies that the map  $(\Phi_e^{w'})^{-1} \circ \Phi_g^w$  is analytic in a neighborhood of  $u$  and we have the Lemma.  $\square$  Lemma 3.4 and Lemma 3.5

assures that we can define an analytic structure on  $\widehat{\mathbf{X}}$  through the maps  $\Phi_g^w$  so that they define analytic diffeomorphisms onto open subsets  $\Omega_g^w$  of  $\widehat{\mathbf{X}}$  and the action of  $G$  on  $\widehat{\mathbf{X}}$  is analytic. On the other hand  $\widehat{\mathbf{X}}$  is Hausdorff because  $\Phi_g^w$  is homeomorphic and  $\overline{N}_\sigma \times M(\sigma) / (M(\sigma) \cap \underline{\omega}^{-1} H \underline{\omega}) \times \mathbb{R}^\Delta$  is Hausdorff. Combining this with Proposition 3.3 we get

**Theorem 3.6.** (i)  $\widehat{\mathbf{X}}$  is a compact connected real analytic manifold and  $\bigcup_{w \in W, g \in G} \Omega_g^w$  is an open covering of  $\widehat{\mathbf{X}}$  such that the maps  $\Phi_g^w$  are real analytic diffeomorphisms.

(ii) The action of  $G$  on  $\widehat{\mathbf{X}}$  is analytic and the orbit  $G\pi(x)$  for a point  $x$  in  $\widehat{\mathbf{X}}$  is isomorphic to  $G/P(x)$ .

(iii) There are just  $2^{|\Delta|}$  open orbits that are isomorphic to  $G/H$  and the number of compact orbits in  $\widehat{\mathbf{X}}$  equal that of the elements of the coset  $W_{K \cap H} \backslash W$ .

**Remark 3.7.** (i) By a similar way, we can construct the compactification of the corresponding Riemannian symmetric spaces  $\mathbf{X}_+ = G^+/K \cap H$ .

(ii) For the special case  $H = K$ , i.e., for the case of Riemannian symmetric spaces of non-compact type, we obtain the corresponding compactification indicated in [5].

**Acknowledgment:** The authors would like to thanks Professor van den Ban, Professor Do Ngoc Diep and Professor Toshio Oshima for many helpful discussions.

## References

- [1] E. van den Ban and H. Schlichtkrull, *Harmonic analysis on reductive symmetric spaces*, Proc. 3<sup>rd</sup> European Congress of Mathematics, 2000.
- [2] A. Borel and Lizhen Ji, *Compactifications of locally symmetric spaces symmetric spaces*, Lectures for the European School of Group Theory, Luminy, France, 2001.
- [3] A. Borel and Lizhen Ji, *Compactifications of symmetric spaces I*, Lectures for the European School of Group Theory, Luminy, France, 2001.
- [4] C. De Concini and C. Procesi, *Complete symmetric varieties*, Lecture Notes in Math., 996 (1983), 1-44, Springer.
- [5] Tran Dao Dong and Tran Vui, *A realization of Riemannian symmetric spaces in compact manifolds*, Proc. of the ICAA 2002, 188-196, Bangkok.
- [6] M. Flensted-Jensen, *Discrete series for semisimple symmetric spaces*, Ann. of Math., 111 (1980), 253-311.
- [7] Lizhen Ji, *Introduction to symmetric spaces and their compactifications*, Lectures for the European School of Group Theory, Luminy, France, 2001.
- [8] W.A. Kosters, *Eigenspaces of the Laplace-Beltrami operator on  $SL(n, \mathbb{R})/S(G(1) \times GL(n-1))$* , Part I, Proc. of Math., University of Leiden (1984).
- [9] T. Matsuki and T. Oshima, *A description of discrete series for semisimple symmetric Spaces*, Adv. Studies in Pure Math., 4(1984), 331-390.
- [10] T. Oshima, *A realization of Riemannian symmetric spaces*, J. Math. Soc. Japan, vol 30 (1978), 117-132.
- [11] T. Oshima, *A realization of semisimple symmetric spaces and construction of boundary value maps*, Advanced Studies in Pure Math., vol 14 (1988), 603-650.

- [12] T. Oshima and J. Sekiguchi, *Eigenspaces of invariant differential operators on an affine symmetric spaces*, Invent Math., vol 57 (1980), 1-81.
- [13] H. Schlichtkrull, “Hyperfunctions and Harmonic Analysis on Symmetric Spaces”, Birkhäuser, Boston, 1984.
- [14] H. Schlichtkrull, *Harmonic analysis on semisimple symmetric spaces*, Lectures for the European School of Group Theory, University of Twente, The Netherlands, 1992.
- [15] J. Sekiguchi, *Eigenspaces of the Laplace-Beltrami operator on a hyperboloid*, Nagoya Math. J., vol 79 (1980), 151-185.