

SYSTEM OF PARAMETERS FOR PSEUDO COHEN-MACAULAY MODULES

Nguyen Thai Hoa and Nguyen Duc Minh*

*Department of Mathematics, Quynhon University, Vietnam
e-mail: minhnd45@hotmail.com*

1 Introduction

Throughout, (A, \mathfrak{m}) denotes a commutative Noetherian local ring with the maximal ideal \mathfrak{m} and M a finitely generated A -module with $\dim M = d$. We denote by $Q_M(\underline{x})$ the submodule of M defined by

$$Q_M(\underline{x}) = \bigcup_{n>0} \left((x_1^{n+1}, \dots, x_d^{n+1})M : x_1^n \cdots x_d^n \right),$$

where $\underline{x} = (x_1, \dots, x_d)$ is a system of parameters on M . The submodule $Q_M(\underline{x})$ is a useful tool in the study of Monomial Conjecture, determinant maps, top local cohomology modules, modules of generalized fractions... (see [25], [4], [6] and [8]).

1.1 Definition The module M is called *pseudo Cohen-Macaulay* if there exists an system of parameters \underline{x} on M such that $e(\underline{x}; M) = \ell_A(M/Q_M(\underline{x}))$

1.2 Example

If $\dim M = 1$, then M is pseudo Cohen-Macaulay by [22].

If M is Cohen-Macaulay, then one can easily see that $Q_M(\underline{x}) = (x_1, \dots, x_d)M$ for an arbitrary system of parameters $\underline{x} = (x_1, \dots, x_d)$ of M . Thus every Cohen-Macaulay module is a pseudo Cohen-Macaulay module. The converse may be not true in general.

*The author is supported by the Swedish International Development Cooperation Agency (SIDA) and The Abdus Salam International Centre Theoretical of Physics (ICTP), Trieste, Italy.

Key words: pseudo Cohen-Macaulay module, system of parameters, Noetherian dimension, coregular element, residual length,
2000 Mathematics Subject Classification: 13C14, 13H14, 16E65, 16G50

This note presents some properties on systems of parameters of pseudo Cohen-Macaulay modules.

2 Preliminaries

2.1 Secondary representation, cosequences, width, Noetherian dimension of Artinian modules

Let L be an Artinian A -module with a minimal secondary representation

$$L = C_1 + \cdots + C_n,$$

where each C_i is \mathfrak{p}_i -secondary. The finite set $\text{Att}(L) = \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$ is called the set of attached prime ideals of L . Set $L_0 = \sum_{\mathfrak{p}_i \in \text{Att}(L) \setminus \{\mathfrak{m}\}} C_i$. Then L_0 is independent of the choice of the minimal secondary representation of L and is called *the residuum of L* . Moreover, the length of the quotient module L/L_0 is finite. This length is called *the residual length of L* and denoted by $R\ell(L)$.

An element $a \in A$ is called *L -coregular element* if $L = aL$. The sequence of elements a_1, \dots, a_n of A is called *an L -cosequence* if $0 :_L (a_1, \dots, a_n) \neq 0$ and a_i is $0 :_L (a_1, \dots, a_{i-1})$ -coregular element for every $i = 1, \dots, n$. We denote by $\text{Width}(L)$ the supremum of lengths of all L -cosequences in \mathfrak{m} .

An element $a \in \mathfrak{m}$ is called *pseudo- L -coregular* if $a \notin \bigcup_{\mathfrak{p} \in \text{Att}(L) \setminus \{\mathfrak{m}\}} \mathfrak{p}$. Note that for each pseudo- L -coregular element $a \in \mathfrak{m}$, there exists $s \in \mathbb{N}$ such that $a^s L = L_0$.

2.1 Lemma (cf. [2, (11.3.9) and (11.3.10)]). *Let $\mathfrak{p} \in \text{Ass}(M)$. Then, $H_{\mathfrak{m}}^{\dim A/\mathfrak{p}}(M) \neq 0$ and $\mathfrak{p} \in \text{Att}(H_{\mathfrak{m}}^{\dim A/\mathfrak{p}}(M))$. Moreover, $\text{Att}(H_{\mathfrak{m}}^d(M)) = \{\mathfrak{p} \in \text{Ass}(M) \mid \dim A/\mathfrak{p} = d\}$.*

The *Noetherian dimension* of L , denoted by $\text{N-dim}_A L$, is defined inductively as follows: when $L = 0$, put $\text{N-dim}_A L = -1$. For an integer $d \geq 0$, we put $\text{N-dim}_A L = d$ if $\text{N-dim}_A L < d$ is false and for every ascending sequence $L_0 \subseteq L_1 \subseteq \cdots$ of submodules of L , there exists n_0 such that $\text{N-dim}_A(L_{n+1}/L_n) < d$ for all $n > n_0$.

It is easy to see that $\text{N-dim}_A L = 0$ if and only if L is a non-zero Noetherian module.

2.2 Lemma ([7]).

(i) *For any exact sequence of Artinian A -modules*

$$0 \longrightarrow L' \longrightarrow L \longrightarrow L'' \longrightarrow 0$$

we have $N\text{-dim } L = \max\{N\text{-dim } L', N\text{-dim } L''\}$.

(ii) $N\text{-dim}(L) \leq \dim(L)$. The equality holds if A is complete.

(iii) $N\text{-dim}_A(L) = N\text{-dim}_{\hat{A}}(L) = \dim_{\hat{A}}(L)$.

(iv) $N\text{-dim}(H_{\mathfrak{m}}^i(M)) \leq i, \forall i = 0, \dots, d-1$ and $N\text{-dim}(H_{\mathfrak{m}}^d(M)) = d$.

2.2 The invariants $p(M), pf(M)$ and pseudo Cohen-Macaulay modules.

Let $\underline{x} = (x_1, \dots, x_d)$ be an system of parameters of M and $\underline{n} = (n_1, \dots, n_d)$ a d -tuple of positive integers. Set $\underline{x}(\underline{n}) = (x_1^{n_1}, \dots, x_d^{n_d})A$. Consider the differences

$$\begin{aligned} I_{M, \underline{x}}(\underline{n}) &= \ell(M/\underline{x}(\underline{n})M) - n_1 \dots n_d e(\underline{x}; M); \\ J_{M, \underline{x}}(\underline{n}) &= n_1 \dots n_d e(\underline{x}; M) - \ell(M/Q_M(\underline{x}(\underline{n}))) \end{aligned}$$

as functions in n_1, \dots, n_d , where $e(\underline{x}; M)$ is the multiplicity of M with respect to \underline{x} and

$$Q_M(\underline{x}) = \bigcup_{t>0} ((x_1^{t+1}, \dots, x_d^{t+1})M : x_1^t \dots x_d^t).$$

In general, $I_{M, \underline{x}}(\underline{n})$ and $J_{M, \underline{x}}(\underline{n})$ are not polynomials for n_1, \dots, n_d large enough (see [3], [6]). However they are bounded above by polynomials and the least degree of all polynomials in \underline{n} bounding above $I_{M, \underline{x}}(\underline{n})$ (resp. $J_{M, \underline{x}}(\underline{n})$) is independent of the choice of \underline{x} , and it is denoted by $p(M)$ (resp. $pf(M)$). The invariant $p(M)$ is called *the polynomial type* of M (see [3]) and the invariant $pf(M)$ is called *the polynomial type of fractions* of M (see [16], [5] and [4]). For convenience we stipulate that the degree of the zero-polynomial is equal to $-\infty$. One can easy to see that following the conditions are equivalent:

- (i) M is pseudo Cohen-Macaulay
- (ii) $pf(M) = -\infty$
- (iii) For every system of parameters \underline{x} on M we have $e(\underline{x}; M) = \ell_A(M/Q_M(\underline{x}))$

Let us list basic facts on $p(M)$ and $pf(M)$ from [3], [16], and [5].

2.3 Lemma ([3] and [5]).

(i) $p(M) = p(M/H_{\mathfrak{m}}^0(M)) = p_{A/\text{Ann}(M)}(M)$

$pf_A(M) = pf_A(M/H_{\mathfrak{m}}^0(M)) = pf_{A/\text{Ann}(M)}(M)$

(ii) $p_A(M) = p_{\hat{A}}(\widehat{M}), pf_A(M) = pf_{\hat{A}}(\widehat{M})$, where \widehat{M} is the m -adic completion of M .

(iii) Let \underline{x} be an system of parameters of M with $\dim(0 : x_1) < d-1$. Then

$$pf(M/x_1M) \leq pf(M) \leq pf(M/x_1M) + 1.$$

2.4 Lemma ([5, (3.4) and (3.5)]).

- (i) $p(M) \leq \dim M - 1$ and if $\dim M = d > 1$ then $pf(M) \leq d - 2$.
- (ii) $pf(M) \leq p(M)$. If $\text{depth}(M) > p(M)$ then $pf(M) = p(M)$.

2.5 Lemma ([5, (3.6)]).

- (i) If $pf(M) = -\infty$ then $H_{\mathfrak{m}}^i(M) = 0$ for all $i = p(M) + 1, \dots, d - 1$.
- (ii) If $pf(M) \leq 0$ then $\ell(H_{\mathfrak{m}}^i(M)) < \infty$ for all $i = p(M) + 1, \dots, d - 1$.

2.6 Proposition Assume that $\dim M = d \geq 1$. Then,

- (i) $p(M) = \max_{0 \leq i \leq d-1} \{N\text{-dim } H_{\mathfrak{m}}^i(M)\}$,

(ii) Suppose that $p = p(M) > 0$. Set $\mathcal{Q} = \bigcup_{i=p}^{d-1} \text{Att}(H_{\mathfrak{m}}^i(M)) \setminus \{\mathfrak{m}\}$. Let x be a parameter element of M such that $x \notin \bigcup_{\mathfrak{p} \in \mathcal{Q}} \mathfrak{p}$. Then $p(M/xM) = p(M) - 1$.

Proof (i). Denote $a_i(M)$ be the annihilator of the i -th local cohomology module $H_{\mathfrak{m}}^i(M)$ of M with respect to the maximal ideal \mathfrak{m} and set $\mathfrak{a}(M) = \mathfrak{a}_0(M) \cdots \mathfrak{a}_{d-1}(M)$. It follows from Lemma 2.3, [3, (3.1)] and Lemma 2.2 that

$$\begin{aligned} p_A(M) &= p_{\widehat{A}}(\widehat{M}) = \dim_{\widehat{A}} \widehat{A}/\mathfrak{a}(\widehat{M}) &= \max_{0 \leq i \leq d-1} \left\{ \dim_{\widehat{A}} H_{\mathfrak{m}}^i(\widehat{M}) \right\} \\ &= \max_{0 \leq i \leq d-1} \left\{ N\text{-dim}_{\widehat{A}}(H_{\mathfrak{m}}^i(\widehat{M})) \right\} &= \max_{0 \leq i \leq d-1} \left\{ N\text{-dim}(H_{\mathfrak{m}}^i(M)) \right\}. \end{aligned}$$

(ii). Let $x \notin \bigcup_{\mathfrak{p} \in \mathcal{Q}} \mathfrak{p}$ be a parameter element of M . Choose $x_2, \dots, x_d \in A$ such that $\underline{x} = (x, x_2, \dots, x_d)$ is an system of parameters of M . For each $\mathfrak{q} \in \text{Ass}(M)$ with $\dim A/\mathfrak{q} \geq p(M)$ we have $\mathfrak{q} \in \text{Att}(H_{\mathfrak{m}}^{\dim A/\mathfrak{q}}(M))$ by Lemma 2.1. Thus $x \notin \mathfrak{q}$ for each $\mathfrak{q} \in \text{Ass}(M)$ with $\dim(A/\mathfrak{q}) \geq p$. This implies that $\dim(0 :_M x) < p < \dim M$. Hence, $e(\underline{x}'; M/xM) = e(\underline{x}; M)$ where $\underline{x}' = (x_2, \dots, x_d)$. By [3] (2.2), we get

$$I_M(\underline{n}, \underline{x}) \leq n_1 I_M((1, n_2, \dots, n_d); \underline{x}) = n_1 I_{M/xM}((n_2, \dots, n_d); \underline{x}'),$$

where $\underline{n} = (n_1, n_2, \dots, n_d)$ is a d -tuple positive integers. Therefore $p(M) \leq p(M/xM) + 1$.

We next show the converse inequality $p(M/xM) + 1 \leq p(M)$. As $p(M) > 0$, we need only to argue for $p(M/xM) > 0$. By the statement (i), there exists $j \in \{0, \dots, d-2\}$ such that $p(M/xM) = N\text{-dim } H_{\mathfrak{m}}^j(M/xM)$. There are only two situations arising.

Case 1: $0 \leq j < p$. In this case

$$p(M/xM) = N\text{-dim}(H_{\mathfrak{m}}^j(M/xM)) \leq j \leq p - 1$$

by Lemma 2.2.

Case 2: $p \leq j \leq d - 2$. As $\dim(0 :_M x) < p \leq j$, we have $H_{\mathfrak{m}}^j(0 :_M x) = H_{\mathfrak{m}}^{j+1}(0 :_M x) = 0$. Therefore, the exact sequences

$$0 \longrightarrow (0 :_M x) \longrightarrow M \longrightarrow M/(0 :_M x) \longrightarrow 0$$

and

$$0 \longrightarrow M/(0 :_M x) \xrightarrow{x} M \longrightarrow M/xM \longrightarrow 0$$

induce an exact sequence of local cohomology modules

$$0 \longrightarrow H_{\mathfrak{m}}^j(M)/xH_{\mathfrak{m}}^j(M) \longrightarrow H_{\mathfrak{m}}^j(M/xM) \longrightarrow (0 :_{H_{\mathfrak{m}}^{j+1}(M)} x) \longrightarrow 0 \quad (1)$$

By our assumption, x is pseudo- $H_{\mathfrak{m}}^j(M)$ -coregular. Consequently, $\text{N-dim}(H_{\mathfrak{m}}^j(M)/xH_{\mathfrak{m}}^j(M)) \leq 0$. Moreover, when $\text{N-dim}(H_{\mathfrak{m}}^{j+1}(M)) > 0$, $x \notin \bigcup_{\mathfrak{p} \in \mathcal{Q}} \mathfrak{p}$ implies that x is a parameter element on $H_{\mathfrak{m}}^{j+1}(M)$ so that

$$\text{N-dim}(0 :_{H_{\mathfrak{m}}^{j+1}(M)} x) = \text{N-dim}(H_{\mathfrak{m}}^{j+1}(M)) - 1.$$

It now follows from the exact sequence (1) that

$$\begin{aligned} 0 < p(M/xM) &= \text{N-dim}(H_{\mathfrak{m}}^j(M/xM)) \\ &= \max \left\{ \text{N-dim}(H_{\mathfrak{m}}^j(M)/xH_{\mathfrak{m}}^j(M)); \text{N-dim}(0 :_{H_{\mathfrak{m}}^{j+1}(M)} x) \right\} \\ &= \text{N-dim}(0 :_{H_{\mathfrak{m}}^{j+1}(M)} x) = \text{N-dim}(H_{\mathfrak{m}}^{j+1}(M)) - 1 \leq p(M) - 1. \end{aligned} \quad \square$$

2.7 Lemma *Let $\underline{x} = (x_1, x_2, \dots, x_d)$ be a system of parameters of M . Put $M_1 = M/x_1M$. For each $\underline{n}' = (n_2, \dots, n_d) \in \mathbb{N}^{d-1}$, set $\underline{x}'(\underline{n}') := (x_2^{n_2}, \dots, x_d^{n_d})$ and $\underline{x}(\underline{n}') = (x_1, x_2^{n_2}, \dots, x_d^{n_d})$. Then, there exists an epimorphism*

$$\varphi_{\underline{n}'} : M_1/Q_{M_1}(\underline{x}'(\underline{n}')) \longrightarrow M/Q_M(\underline{x}(\underline{n}'))$$

defined by $\varphi_{\underline{n}'}(\bar{u} + Q_{M_1}(\underline{x}'(\underline{n}'))) = u + Q_M(\underline{x}(\underline{n}'))$ for each $u \in M$.

Moreover, if $x_1 \notin \bigcup_{\mathfrak{q} \in \text{Att}(H_{\mathfrak{m}}^{d-1}(M)) \setminus \{\mathfrak{m}\}} \mathfrak{q}$, then for all n_2, \dots, n_d enough large,

we have an exact sequence

$$0 \longrightarrow H_{\mathfrak{m}}^{d-1}(M)/x_1H_{\mathfrak{m}}^{d-1}(M) \longrightarrow M_1/Q_{M_1}(\underline{x}'(\underline{n}')) \xrightarrow{\varphi_{\underline{n}'}} M/Q_M(\underline{x}(\underline{n}')) \longrightarrow 0.$$

Proof For d -tuples of positive integers $\underline{n} = (n_1, \dots, n_d)$ and $\underline{m} = (m_1, \dots, m_d)$ we define $\underline{n} \leq \underline{m}$ if $n_i \leq m_i$ for all i . Then the map

$$\delta_{\underline{n}, \underline{m}} : M/Q_M(\underline{x}(\underline{n})) \longrightarrow M/Q_M(\underline{x}(\underline{m})),$$

which is defined by $\delta_{\underline{n}, \underline{m}}(u + Q_M(\underline{x}(\underline{n}))) = \prod_{i=1}^d x_i^{m_i - n_i} u + Q_M(\underline{x}(\underline{m}))$ for each $u \in M$, is injective by [4, (3.1)]. Moreover, $\{\delta_{\underline{n}, \underline{m}}; M/Q_M(\underline{x}(\underline{n}))\}$ forms a direct system and $\varinjlim_{\underline{n}} M/Q_M(\underline{x}(\underline{n})) \cong H_{\mathfrak{m}}^d(M)$.

Similarly, we also have the direct system $\{\bar{\delta}_{\underline{n}', \underline{m}'}; M_1/Q_{M_1}(\underline{x}'(\underline{n}'))\}$ and $\varinjlim_{\underline{n}'} M_1/Q_{M_1}(\underline{x}'(\underline{n}')) \cong H_{\mathfrak{m}}^{d-1}(M_1)$.

For $\underline{n}' \leq \underline{m}'$ with $\underline{n}' = (n_2, \dots, n_d)$ and $\underline{m}' = (m_2, \dots, m_d)$, we have the following commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \text{Ker } \varphi_{\underline{n}'} & \longrightarrow & M_1/Q_{M_1}(\underline{x}'(\underline{n}')) & \xrightarrow{\varphi_{\underline{n}'}} & M/Q_M(\underline{x}(\underline{n}')) & \longrightarrow & 0 \\ & & \downarrow \pi_{\underline{n}', \underline{m}'} & & \downarrow \bar{\delta}_{\underline{n}', \underline{m}'} & & \downarrow \delta_{\underline{n}', \underline{m}'} & & \\ 0 & \longrightarrow & \text{Ker } \varphi_{\underline{m}'} & \longrightarrow & M_1/Q_{M_1}(\underline{x}'(\underline{m}')) & \xrightarrow{\varphi_{\underline{m}'}} & M/Q_M(\underline{x}(\underline{m}')) & \longrightarrow & 0 \end{array}$$

where $\pi_{\underline{n}', \underline{m}'}$ is the reduced homomorphism. Thus, $\{\pi_{\underline{n}', \underline{m}'}; \text{Ker } \varphi_{\underline{n}'}\}$ and $\{\delta_{\underline{n}', \underline{m}'}; M/Q_M(\underline{x}(\underline{n}'))\}$ form direct systems. Therefore, we obtain the following commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \text{Ker } \varphi_{\underline{n}'} & \longrightarrow & M_1/Q_{M_1}(\underline{x}'(\underline{n}')) & \xrightarrow{\varphi_{\underline{n}'}} & M/Q_M(\underline{x}(\underline{n}')) & \longrightarrow & 0 \\ & & \downarrow \pi_{\underline{n}'} & & \downarrow \bar{\delta}_{\underline{n}'} & & \downarrow \delta_{\underline{n}'} & & \\ 0 & \longrightarrow & \varinjlim_{\underline{n}'} \text{Ker } \varphi_{\underline{n}'} & \longrightarrow & H_{\mathfrak{m}}^{d-1}(M_1) & \xrightarrow{u} & \varinjlim_{\underline{n}'} M/Q_M(\underline{x}(\underline{n}')) & \longrightarrow & 0 \end{array}$$

where $\pi_{\underline{n}'}, \bar{\delta}_{\underline{n}'}$ and $\delta_{\underline{n}'}$ are the natural homomorphisms and $u = \varinjlim_{\underline{n}'} \varphi_{\underline{n}'}$. Since the homomorphisms $\delta_{\underline{n}', \underline{m}'}, \bar{\delta}_{\underline{n}', \underline{m}'}$ and $\pi_{\underline{n}', \underline{m}'}$ are injective, we have $\delta_{\underline{n}'}, \bar{\delta}_{\underline{n}'}$ and $\pi_{\underline{n}'}$ are injective.

By Lemma 2.1, $x_1 \notin \mathfrak{q}$ for all $\mathfrak{q} \in \text{Ass}(M)$ with $\dim A/\mathfrak{q} \geq d-1$. Thus $\dim(0 :_M x_1) < d-1$ and then $H_{\mathfrak{m}}^i(M/(0 :_M x_1)) \cong H_{\mathfrak{m}}^i(M)$ for $i \geq d-1$. Therefore, from the exact sequence

$$0 \longrightarrow M/(0 :_M x_1) \xrightarrow{x_1} M \longrightarrow M_1 \longrightarrow 0,$$

we have an exact sequence of local cohomology modules

$$0 \longrightarrow H_{\mathfrak{m}}^{d-1}(M)/x_1 H_{\mathfrak{m}}^{d-1}(M) \longrightarrow H_{\mathfrak{m}}^{d-1}(M_1) \xrightarrow{\Delta} H_{\mathfrak{m}}^d(M),$$

where Δ is the connecting homomorphism.

Further, we can also show a monomorphism $j : \varinjlim_{\underline{n}'} M/Q_M(\underline{x}(\underline{n}')) \longrightarrow H_{\mathfrak{m}}^d(M)$ such that the following diagram is commutative

$$\begin{array}{ccc}
 H_{\mathfrak{m}}^{d-1}(M_1) & \xrightarrow{u} & \varinjlim_{\underline{n}'} M/Q_M(\underline{x}(\underline{n}')) \\
 \searrow \Delta & & \swarrow j \\
 & & H_{\mathfrak{m}}^d(M)
 \end{array}$$

Hence $\varinjlim_{\underline{n}'} \text{Ker } \varphi_{\underline{n}'} \cong \text{Ker } u \cong \text{Ker } \Delta \cong H_{\mathfrak{m}}^{d-1}(M)/x_1 H_{\mathfrak{m}}^{d-1}(M)$. Since $H_{\mathfrak{m}}^{d-1}(M)/x_1 H_{\mathfrak{m}}^{d-1}(M)$ has finite length by the choice of x_1 , $\pi_{\underline{n}'}$ is an isomorphism for enough large \underline{n}' ($\underline{n}' \gg 0$ for short). So we get

$$\text{Ker } \varphi_{\underline{n}'} \cong H_{\mathfrak{m}}^{d-1}(M)/x_1 H_{\mathfrak{m}}^{d-1}(M)$$

for $\underline{n}' \gg 0$ as required. \square

2.8 Corollary *Let M be a pseudo Cohen-Macaulay module with $p := p(M) > 0$. Let x_1 be a parameter element with $\dim(0 :_M x_1) < d - 1$. Then x_1 is a $H_{\mathfrak{m}}^{d-1}(M)$ -coregular element.*

Proof With the same notations and using the same argument in the proof of Lemma 2.7 we have $\varinjlim_{\underline{n}'} \text{Ker } \varphi_{\underline{n}'} \cong H_{\mathfrak{m}}^{d-1}(M)/x_1 H_{\mathfrak{m}}^{d-1}(M)$. On the other hand, by virtue of Lemma 2.3, M_1 is a pseudo Cohen-Macaulay module. Thus,

$$\ell_A(M/Q_M(\underline{x}(\underline{n}'))) = e(\underline{x}(\underline{n}'); M) = e(\underline{x}'(\underline{n}'); M_1) = \ell_A(M_1/Q_{M_1}(\underline{x}'(\underline{n}'))).$$

Therefore, the epimorphism $\varphi_{\underline{n}'} : M_1/Q_{M_1}(\underline{x}'(\underline{n})) \rightarrow M/Q_M(\underline{x}(\underline{n}))$ defined in Lemma 2.7 must be an isomorphism. This implies that $\text{Ker } \varphi_{\underline{n}'} = 0$ for all $\underline{n}' \in \mathbb{N}^{d-1}$. Hence $\varinjlim_{\underline{n}'} \text{Ker } \varphi_{\underline{n}'} = 0$ and so $H_{\mathfrak{m}}^{d-1}(M) = x_1 H_{\mathfrak{m}}^{d-1}(M)$, as required. \square

3 Parametric characterizations for pseudo Cohen-Macaulay modules

Following [3], a subsequence (x_1, \dots, x_j) of a system of parameters of M is called a reducing sequence if $x_i \notin \mathfrak{p}$ for all $\mathfrak{p} \in \text{Ass}(M/(x_1, \dots, x_{i-1})M)$ with $\dim A/\mathfrak{p} \geq d - i$, ($i = 1, \dots, j$). Note that if $\underline{x} = (x_1, \dots, x_d)$ is a system of parameters on M and x_1, \dots, x_{d-1} form a reducing sequence, then \underline{x} is just a reducing system of parameters as introduced in [1]. It should be mentioned that every A -module admits a reducing parameter system of parameters.

3.1 Definition Let $\underline{x} = (x_1, \dots, x_t)$ be a sequence of elements in \mathfrak{m} . We set $M_i := M/(x_1, \dots, x_i)M$ for all $i = 0, \dots, t$. The sequence \underline{x} is called *pseudo regular* for M if x_i is an $H_{\mathfrak{m}}^{d-i}(M_{i-1})$ -coregular element for all $i = 1, \dots, t$. If $\underline{x} = (x_1, \dots, x_d)$ is an system of parameters on M and (x_1, \dots, x_{d-1}) forms a pseudo regular sequence, then it is called a *pseudo regular system of parameters*.

3.2 Remark

(i) An arbitrary system of parameters of a Cohen-Macaulay module M is a pseudo regular system of parameters

(ii) $\underline{x} = (x_1, \dots, x_t)$ is a pseudo regular sequence of M if and only if (x_1, \dots, x_{j-1}) is pseudo regular sequence of M and (x_j, \dots, x_t) is a pseudo regular sequence of M_{j-1} for each $j = 2, \dots, t$.

(iii) By Lemma 2.1, every pseudo regular system of parameters for M is a reducing system of parameters of M .

3.3 Theorem Assume that $\dim M = d > 1$. Then the following statements are equivalent:

- (i) M is pseudo Cohen-Macaulay;
- (ii) Any reducing system of parameters of M is pseudo regular system of parameters;
- (iii) M admits a reducing system of parameters which is pseudo regular system of parameters;
- (iv) M admits a pseudo regular system of parameters.

Proof It suffices to prove that (i) \implies (ii) and (iv) \implies (i).

(i) \implies (ii). We prove by induction on d . Let $d = 2$ and assume that $\underline{x} = (x_1, x_2)$ is a reducing system of parameters of M . By Corollary 2.8, x_1 is $H_{\mathfrak{m}}^1(M)$ -coregular and then \underline{x} is pseudo regular system of parameters of M . Suppose that $d > 2$ and that our assertion is true for all pseudo Cohen-Macaulay A -modules of smaller dimension.

Let $\underline{x} = (x_1, \dots, x_d)$ be a reducing system of parameters of M . As $\dim(0 : x_1) < d - 1$, then $M_1 := M/x_1M$ is pseudo Cohen-Macaulay by virtue of Lemma 2.3 (iii). The inductive hypothesis implies that (x_2, \dots, x_d) is a pseudo regular system of parameters of M_1 .

The induction is finished now by Corollary 2.8 and Remark 3.2 (ii).

(iv) \implies (i). Again we use induction on d . Let $d = 2$, and assume that M has a pseudo regular system of parameters, say $\underline{x} = (x_1, x_2)$. Let $\underline{n} = (n_1, n_2) \in \mathbb{N}^2$. By [22, (3.2)], $J_{M, \underline{x}}(\underline{n}) = \text{Rl}(H_{\mathfrak{m}}^1(M))$ for all $n_1, n_2 \gg 0$. As x_1 is $H_{\mathfrak{m}}^1(M)$ -coregular, it follows that $\text{Rl}(H_{\mathfrak{m}}^1(M)) = 0$ and so M is pseudo Cohen-Macaulay.

Assume that $d > 2$ and $\underline{x} = (x_1, x_2, \dots, x_d)$ is a pseudo regular system of parameters for M . Then $\underline{x}' = (x_2, \dots, x_d)$ is a pseudo regular system of parameters

for $M_1 = M/x_1M$. For all $n_2, \dots, n_d \geq 1$, set $\underline{x}(\underline{n}') = (x_1, x_2^{n_2}, \dots, x_d^{n_d})$, $\underline{x}'(\underline{n}') = (x_2^{n_2}, \dots, x_d^{n_d})$. The inductive hypothesis gives

$$\underline{x}'(\underline{n}'); M_1 = \ell(M_1/Q_{M_1}(\underline{x}'(\underline{n}'))). \tag{2}$$

On the other hand, Lemma 2.1 shows that $x_1 \notin \mathfrak{p}$, for all $\mathfrak{p} \in \text{Ass}(M)$ with $\dim A/\mathfrak{p} \geq d - 1$. We thus have $\dim(0 : x_1) < d - 1$ and hence

$$e(\underline{x}'(\underline{n}'); M_1) = e(\underline{x}(\underline{n}'); M). \tag{3}$$

Take n_2, \dots, n_d large enough to obtain the exact sequence defined in Lemma 2.7,

$$0 \longrightarrow H_{\mathfrak{m}}^{d-1}(M)/x_1H_{\mathfrak{m}}^{d-1}(M) \longrightarrow M_1/Q_{M_1}(\underline{x}'(\underline{n}')) \longrightarrow M/Q_M(\underline{x}(\underline{n}')) \longrightarrow 0.$$

As x_1 is $H_{\mathfrak{m}}^{d-1}(M)$ -coregular, we obtain from above exact sequence that

$$\ell_A(M_1/Q_{M_1}(\underline{x}'(\underline{n}'))) = \ell_A(M/Q_M(\underline{x}(\underline{n}))). \tag{4}$$

Combining (2), (3) and (4), for all $n_2, \dots, n_d \gg 0$, we get

$$e(\underline{x}(\underline{n}'); M) = \ell_A(M/Q_M(\underline{x}(\underline{n})))$$

and this finishes our proof. □

According to [9], M is called an f -module if every system of parameters $\underline{x} = (x_1, \dots, x_d)$ is a M -filter regular sequence, i.e. $x_i \notin \mathfrak{q}$ for all $\mathfrak{q} \in \text{Ass}(M/(x_1, \dots, x_{i-1})M) \setminus \{\mathfrak{m}\}$; ($i = 1, \dots, d$).

We next combine Theorem 3.3 with [9, (2.5) and (2.11)] to obtain

3.4 Corollary *M is both f -module and pseudo Cohen-Macaulay if and only if every system of parameters for M is pseudo regular system of parameters.*

3.5 Theorem *Suppose that $p = p(M) > 0$. Then M is pseudo Cohen-Macaulay if and only if $H_{\mathfrak{m}}^i(M) = 0$ for all $i = p+1, \dots, d-1$ and there exists a subsystem of parameters (x_1, \dots, x_p) on M such that x_i is an $H_{\mathfrak{m}}^{p-i+1}(M_{i-1})$ -coregular element for all $i = 1, \dots, p$.*

Proof Assume that M is pseudo Cohen-Macaulay with $p(M) > 0$. Then $H_{\mathfrak{m}}^i(M) = 0$ for all $i = p+1, \dots, d-1$ by Lemma 2.5. It follows from Corollary 2.8 that $\text{Width}(H_{\mathfrak{m}}^{d-1}(M)) > 0$ and that $m \notin \text{Att}(H_{\mathfrak{m}}^p(M))$. Set

$$\mathfrak{P} = \{\mathfrak{q} \in \text{Ass}(M) \mid \dim A/\mathfrak{q} = d\} \cup \text{Att}(H_{\mathfrak{m}}^p(M))$$

and choose $x_1 \notin \bigcup_{\mathfrak{q} \in \mathfrak{p}} \mathfrak{q}$. Obviously, x_1 is a parameter element of M and also a $H_{\mathfrak{m}}^p(M)$ -coregular element. Observe that $p(M/x_1M) = p(M) - 1$ by Proposition 2.6. Now the existence of the required subsystem of parameters (x_1, \dots, x_p) follows by induction on p .

Conversely, assume that $H_{\mathfrak{m}}^i(M) = 0$ for all $i = p + 1, \dots, d - 1$ and that M admits a subsystem of parameters (x_1, \dots, x_p) such that x_i is an $H_{\mathfrak{m}}^{p-i+1}(M)$ -coregular element for all $i = 1, \dots, p$. Take x_{p+1}, \dots, x_d such that $\underline{x} = (x_1, \dots, x_p, x_{p+1}, \dots, x_d)$ becomes a system of parameters on M . We will prove by induction on p that $J_{M, \underline{x}}(\underline{n}) = 0$ for all $\underline{n} \gg 0$.

The case $p = 1$ was proved in [5, (4.4)].

Assume that $p > 1$ and that our claim is true for all modules with polynomial type less than p . Set $M_1 = M/x_1M$. Because $H_{\mathfrak{m}}^i(M) = 0$ for all $i = p + 1, \dots, d - 1$ and x_1 is $H_{\mathfrak{m}}^p(M)$ -coregular, Lemma 2.1 shows that $x_1 \notin \mathfrak{q}$ for all $\mathfrak{q} \in \text{Ass}(M)$ with $\dim A/\mathfrak{q} \geq p$. Therefore $\dim(0 :_M x_1) < p \leq d - 1$, $e(x_1, \dots, x_d; M) = e(x_2, \dots, x_d; M_1)$ and (by Proposition 2.6) $p(M_1) = p - 1 > 0$. Furthermore, for all $n_2, \dots, n_d \gg 0$, Lemma 2.7 gives us $M_1/Q_{M_1}(\underline{x}'(\underline{n}')) \cong M/Q_M(\underline{x}(\underline{n}'))$, where $\underline{x}'(\underline{n}')) = (x_2^{n_2}, \dots, x_d^{n_d})$ and $\underline{x}(\underline{n}')) = (x_1, x_2^{n_2}, \dots, x_d^{n_d})$. Hence

$$J_{M, \underline{x}}(\underline{n}) = J_{M_1, \underline{x}'}(\underline{n}'), \forall n_2, \dots, n_d \gg 0. \quad (5)$$

On the other hand, since $\dim(0 :_M x_1) < p$ for each $i \in \{p, \dots, d - 1\}$, we have an exact sequence

$$0 \longrightarrow H_{\mathfrak{m}}^i(M)/x_1 H_{\mathfrak{m}}^i(M) \longrightarrow H_{\mathfrak{m}}^i(M_1) \longrightarrow (0 :_{H_{\mathfrak{m}}^{i+1}(M)} x_1) \longrightarrow 0 \quad (6)$$

Since $H_{\mathfrak{m}}^{i+1}(M) = 0$ for all $i = p, \dots, d - 2$ and $(H_{\mathfrak{m}}^p(M)/x_1 H_{\mathfrak{m}}^p(M)) = 0$, the exact sequence (6) implies $H_{\mathfrak{m}}^i(M_1) = 0$ for all $i = p, \dots, d - 2$. The induction is complete by applying the inductive hypothesis to M_1 and using the equality (5). \square

The next result is an immediate consequence of Theorem 3.5 and Proposition 2.6 (ii).

3.6 Corollary *Let M be pseudo Cohen-Macaulay with $p = p(M) > 0$. Then M admits a subsystem of parameters (x_1, \dots, x_p) such that*

$$\text{N-dim}(H_{\mathfrak{m}}^{p-i+1}(M_{i-1})) = p - i + 1$$

and

$$\text{Width}(H_{\mathfrak{m}}^{p-i+1}(M_{i-1})) \geq \min\{2, p - i + 1\}.$$

for all $i = 1, \dots, p$.

The rest of this section is devoted to results on sequentially Cohen-Macaulay modules. These modules was first introduced by P. Stanley in [24] (Chapter III, 2.9) in the graded case. We recall here a definition for the local case from [8].

3.7 Definition ([8, (4.1)]. A filtration $0 = N_0 \subset N_1 \subset \dots \subset N_t = M$ of submodules of M is said to be a *Cohen-Macaulay filtration* if

- (a) Each quotient N_i/N_{i-1} is Cohen-Macaulay.
- (b) $\dim N_1/N_0 < \dim N_2/N_1 < \dots < \dim N_t/N_{t-1}$.

We say that M is *sequentially Cohen-Macaulay* if it admits a Cohen-Macaulay filtration.

3.8 Lemma *Let M be a sequentially Cohen-Macaulay A -module. Then, for all each $i = 0, \dots, d$, the local cohomology module $H_{\mathfrak{m}}^i(M)$ vanishes or is a co-Cohen-Macaulay module of Noetherian dimension i .*

Proof Let $\{M_i\}_{0 \leq i \leq d}$ be a Cohen-Macaulay filtration of M . Set $\mathcal{M}_i = M_i/M_{i-1}$ for all $i = 1, \dots, d$ and $\mathcal{M}_0 = M_0$. If \mathcal{M}_i does not vanish, then it is Cohen-Macaulay module of dimension i . It follows from [17] and [7, (3.5)] that

$$\text{Width}(H_{\mathfrak{m}}^i(\mathcal{M}_i)) = i = N - \dim(H_{\mathfrak{m}}^i(\mathcal{M}_i)).$$

Since $H_{\mathfrak{m}}^i(M) \cong H_{\mathfrak{m}}^i(\mathcal{M}_i)$, $\forall i \geq 0$ by [20] (5.4), this equality shows that $H_{\mathfrak{m}}^i(M)$ is a co-Cohen-Macaulay module. \square

3.9 Theorem *Suppose that $d \geq 1$. Then the following conditions are equivalent:*

- (i) M is a sequentially Cohen-Macaulay module;
- (ii) If $\underline{x} = (x_1, \dots, x_d)$ is an arbitrary filter-regular system of parameters of M then x_i is a coregular element on $H_{\mathfrak{m}}^j(M/(x_1, \dots, x_{i-1})M)$ for all $j = 1, \dots, d-i$ and all $i = 1, \dots, d-1$;
- (iii) There exists a filter-regular system of parameters $\underline{x} = (x_1, \dots, x_d)$ of M such that x_i is a coregular element on $H_{\mathfrak{m}}^j(M/(x_1, \dots, x_{i-1})M)$ for all $j = 1, \dots, d-i$ and all $i = 1, \dots, d-1$;
- (iv) There exists a system of parameters $\underline{x} = (x_1, \dots, x_d)$ of M such that x_i is a coregular element on $H_{\mathfrak{m}}^j(M/(x_1, \dots, x_{i-1})M)$ for all $j = 1, \dots, d-i$ and all $i = 1, \dots, d-1$.

Proof It is enough to prove (i) \implies (ii) and (iv) \implies (i).

(i) \implies (ii). We make induction on d . It is clearly true for $d = 1$. Suppose that $d \geq 2$ and that statement (ii) is true for all modules of dimension $< d$.

Let $\underline{x} = (x_1, \dots, x_d)$ be a filter regular system of parameters of M . Let $i \in \{0, \dots, d\}$. Since M is a sequentially Cohen-Macaulay, $H_{\mathfrak{m}}^i(M)$ is zero or a co-Cohen-Macaulay of Noetherian dimension i by Lemma 3.8. By Lemma 2.1 this implies that x_1 is a $H_{\mathfrak{m}}^i(M)$ -coregular element. Thus $H_{\mathfrak{m}}^i(M)/x_1 H_{\mathfrak{m}}^i(M) = 0$ and $(0 :_{H_{\mathfrak{m}}^i(M)} x_1)$ is zero or a co-Cohen-Macaulay module.

On the other hand, since x_1 is a filter-regular element of M , we have $\dim(0 :_M x_1) = 0$. This yields the exact sequence

$$0 \longrightarrow H_{\mathfrak{m}}^j(M)/x_1 H_{\mathfrak{m}}^j(M) \longrightarrow H_{\mathfrak{m}}^j(M/x_1 M) \longrightarrow (0 :_{H_{\mathfrak{m}}^{j+1}(M)} x_1) \longrightarrow 0$$

for all $j = 1, \dots, d-2$. Thus, $H_{\mathfrak{m}}^j(M/x_1 M) \cong (0 :_{H_{\mathfrak{m}}^{j+1}(M)} x_1)$ for all $j = 1, \dots, d-2$. Therefore, for each $j \in \{1, \dots, d\}$, $H_{\mathfrak{m}}^j(M/x_1 M)$ vanishes or is a co-Cohen-Macaulay. Observe that $\underline{x}' = (x_2, \dots, x_d)$ is a filter regular sequence of $M/x_1 M$. So we get claim (ii) by induction on d .

(iv) \implies (i). We use induction on d . Clearly it is true for $d = 1$. Suppose that $d \geq 2$ and that statement (i) is proved for all modules of dimension $< d$. It is easy to see, that x_1 is a filter-regular element of M . Similarly as above, we obtain $H_{\mathfrak{m}}^j(M/x_1 M) \cong (0 :_{H_{\mathfrak{m}}^{j+1}(M)} x_1)$ for all $j = 1, \dots, d-2$. For each $j = 1, \dots, d-3$, this isomorphism and the inductive hypothesis give us that $(0 :_{H_{\mathfrak{m}}^{j+1}(M)} x_1)$ is either zero or a co-Cohen-Macaulay module of Noetherian dimension j .

If $(0 :_{H_{\mathfrak{m}}^{j+1}(M)} x_1) = 0$, then $H_{\mathfrak{m}}^{j+1}(M) = 0$ by the Nakayama Lemma for Artinian modules (see [12]). If $(0 :_{H_{\mathfrak{m}}^{j+1}(M)} x_1)$ is a co-Cohen-Macaulay module of Noetherian dimension j , then $H_{\mathfrak{m}}^{j+1}(M)$ is a Cohen-Macaulay module of dimension $j+1$.

As x_1 is a $H_{\mathfrak{m}}^1(M)$ -coregular element, $H_{\mathfrak{m}}^1(M)$ is either zero or a co-Cohen-Macaulay module of Noetherian dimension 1. By [20, (5.5)], M is a sequentially Cohen-Macaulay module. The proof is now complete. \square

3.10 Corollary *Any sequentially Cohen-Macaulay A -module is pseudo Cohen-Macaulay module.*

References

- [1] Auslander, M. and D. A. Buchsbaum, *Codimension and multiplicity*, Ann. of Math. **68**(1958), 625-657.
- [2] Brodmann, M. P. and R. Y. Sharp, "Local cohomology: an algebraic introduction with geometric applications", Cambridge University Press 1998.

- [3] Cuong, N.T., *On the least degree of polynomials bounding above the differences between lengths and multiplications of certain systems of parameters in local rings*, Nagoya Math. J. **125**(1992), 105-114.
- [4] Cuong, N. T.; N. T. Hoa and N. T. H. Loan, *On certain length functions associated to a system of parameters in local rings*, Vietnam J. Math. **27(3)** (1999), 259-272.
- [5] Cuong N. T. and N.D. Minh, *Lengths of generalized fractions of modules having small polynomial type*, Math. Proc. Camb. Phil. Soc. **128**(2000), 269-282.
- [6] Cuong N.T., Morales M. and L. T. Nhan (2000), *On the length of generalized fractions*, prépublication de l'Institut Fourier n^o 539.
- [7] Cuong N.T. and L. T. Nhan, *On the Noetherian dimension of Artinian modules*, to appear in Vietnam J. Math.
- [8] Cuong N. T. and L. T. Nhan, *On pseudo Cohen-Macaulay modules and pseudo generalized Cohen-Macaulay modules*, in preparation.
- [9] N. T. Cuong, P. Schenzel and N. V. Trung, Verallgemeinerte Cohen - Macaulay Moduln, *Math. Nachr.* **85** (1978), 57-75.
- [10] Hartshorne R., "Residues and Duality", Lecture Notes in Mathematics, **20**, Springer, 1966.
- [11] Herzog J. and Sbarra E., *Sequentially Cohen-Macaulay modules and local cohomology*, in preparation.
- [12] Kirby, D., *Artinian modules and Hilbert polynomials*, Quart. J. Math. Oxford **(2) 24** (1973), 47-57.
- [13] Kirby, D., *Dimension and length of Artinian modules*, Quart. J. Math. Oxford **(2) 41** (1990), 419-429.
- [14] Matsumura, H., "Commutative ring theory", *Cambridge University Press*, 1986.
- [15] MacDonald, I. G., *Secondary representation of modules over a commutative ring*, Sympos. Math. **11** (1973), 23-43.
- [16] Minh, N.D., *On the least degree of polynomials bounding above the differences between multiplicities and length of generalized fractions*, Acta Math. Vietnam. **20(1)**(1995), 115- 128.
- [17] Ooishi, A., *Matlis duality and the width of a module*, Hiroshima Math. J. **6**(1976), 573-587.
- [18] Roberts, R.N., *Krull dimension of Artinian of modules over quasi-local ring*, Quart. J. Math. Oxford **(3) 26** (1975), 269-273.
- [19] Schenzel, P., "Dualisierende Komplexe in der lokalen Algebra und Buchsbaum Ringe", Lecture Notes in Mathematics **907**, Springer-Verlag, Berlin-Heidelberg-New York, 1982.

- [20] Schenzel, P., *On the dimension filtration and Cohen-Macaulay filtered modules*, Commutative algebra and algebraic geometry, (Ferrara), 245-264, Lecture Notes in Pure and Appl. Math., 206, Dekker, New York, 1999.
- [21] Sharp, R. Y., *Some results on the vanishing of local cohomology modules*, Proc. London Math. Soc. **(3) 30**(1975), 177-195.
- [22] Sharp, R. Y. and M. A. Hamieh, *Lengths of certain generalized fractions*, J. Pure Appl. Algebra **38** (1985), 323-336.
- [23] Sharp, R. Y. and H. Zakeri, *Local cohomology and modules of generalized fractions*, Mathematika **29**(1982), 296-306.
- [24] Stanley R.P., "Combinations and commutative algebra", Second edition, Progress in Math., Vol. 41, *Birkhäuser Boston*, 1996.
- [25] Strooker J.R., "Homological Questions in Local Algebra ", *LMS Lecture Note Series*, 145.
- [26] Tang, Z. and H. Zakeri, *Co-Cohen- Macaulay modules and modules of generalized fractions*, Comm. Algebra **(6) 22**(1994), 2173-2204.