

FOLIATIONS FORMED BY K-ORBITS OF MAXIMAL DIMENSION OF SOME MD5-GROUPS

Le Anh Vu

*Department of Mathematics and Informatics
University of Pedagogy, Ho Chi Minh City, Vietnam
e-mail: leanhvu@hcmup.edu.vn*

Abstract

In this paper we consider some MD5-groups and MD5-algebras, i.e., five-dimensional solvable Lie algebras and groups such that their orbits in the co-adjoint representation (K-orbits) are orbits of dimension zero or maximal dimension. We describe the geometry of K-orbits of MD5-groups. The foliations formed by K-orbits of maximal dimension of these MD5-groups and their measurability are also presented in the paper.

Introduction

Let G be an n -dimensional Lie group. It is called an MD n -group (see [4]), iff its orbits in the co-adjoint representation (K-orbits) are orbits of dimension zero or maximal dimension. The corresponding Lie algebras are called MD n -algebras. All MD4-algebras were first listed by D.V Tra in 1984 (see [5]) and then classified up to an isomorphism by the author in 1990 (see [8], [9]). The description of the geometry of K-orbits of all indecomposable MD4-groups, the topological classification of foliations formed by K-orbits of maximal dimension and the characterization of C^* -algebras associated to these foliations by the method of K-functors were also given by the author in 1990 (see [6], [7], [8], [9]). Until now, no complete classification of MD n -algebras with $n \geq 5$ is known. In this paper we concern with a similar problem for MD5-groups and MD5-algebras.

Key words: MD5-group, MD5-algebra, foliation, K-orbit, C^* -algebra, co-adjoint representation, Lie algebra, measurable foliation, transverse measure, X-invariant
2000 AMS Mathematics Subject Classification: Primary 22E45, Secondary 46E25, 20C20

We begin our discussion in Section 2 by giving some interesting examples of MD5-algebras. Section 3 is devoted to the geometric description of K-orbits of MD5-groups corresponding to these MD5-algebras and a discussion of the foliations formed by their maximal dimensional K-orbits. At first, we recall some concepts and notations which will be used later.

1. K-Orbits of a Lie group and measurable foliations

1.1 The Co-adjoint Representation and K-orbits of a Lie Group

1. Let G be a Lie group. We denote by \mathcal{G} the Lie algebra of G and by \mathcal{G}^* the dual space of \mathcal{G} . To each element g of G we associate an automorphism

$$A_{(g)} : G \longrightarrow G$$

$$x \longmapsto A_{(g)}(x) := gxg^{-1}$$

(which is called the internal automorphism associated to g). $A_{(g)}$ induces the tangent map

$$A_{(g)*} : \mathcal{G} \longrightarrow \mathcal{G}$$

$$X \longmapsto A_{(g)*}(X) := \left. \frac{d}{dt} [g \cdot \exp(tX) g^{-1}] \right|_{t=0}.$$

2. The action

$$Ad : G \longrightarrow Aut(\mathcal{G})$$

$$g \longmapsto Ad(g) := A_{(g)*}$$

is called the *adjoint representation* of G in \mathcal{G} .

3. The action

$$K : G \longrightarrow Aut(\mathcal{G}^*)$$

$$g \longmapsto K_{(g)}$$

such that

$$\langle K_{(g)}F, X \rangle := \langle F, Ad(g^{-1})X \rangle; \quad (F \in \mathcal{G}^*, X \in \mathcal{G})$$

is called the *co-adjoint representation* of G in \mathcal{G}^* .

4. Each orbit of the co-adjoint representation of G is called a *K-orbit*. The dimension of a K-orbit of G is always even (see [2]).

1.2 Measurable Foliations

1. Let V be a smooth manifold. We denote by TV its tangent bundle, so that for each $x \in V$, T_xV is the tangent space of V at x . A smooth subbundle

\mathcal{F} of TV is called *integrable* iff the following condition is satisfied: every x from V is contained in a submanifold W of V such that $T_p(W) = \mathcal{F}_p (\forall p \in W)$.

2. A *foliation* (V, \mathcal{F}) is given by a smooth manifold V and an integrable subbundle \mathcal{F} of TV. Then, V is called the *foliated manifold* and \mathcal{F} is called *the subbundle defining the foliation*.

3. The *leaves* of the foliation (V, \mathcal{F}) are the maximal connected submanifolds L of V with $T_x(L) = \mathcal{F}_x (\forall x \in L)$.

The set of leaves with the quotient topology is denoted by V/\mathcal{F} and will be called the *space of leaves* of (V, \mathcal{F}) . It is a fairly untractable topological space.

4. The partition of V in leaves $V = \bigcup_{\alpha \in V/\mathcal{F}} L_\alpha$ is characterized geometrically by the following local triviality: Every $x \in V$ has a system of local coordinates $\{U; x^1, x^2, \dots, x^n\} (x \in U; n = \dim \mathcal{F})$ such that for any leaf L with $L \cap U \neq \emptyset$, each connected component of $L \cap U$ (which is called a *plaque* of the leaf L) is given by the equations

$$x^{k+1} = c^1, x^{k+2} = c^2, \dots, x^n = c^{n-k}; k = \dim \mathcal{F},$$

where c^1, c^2, \dots, c^{n-k} are constants (depending on each plaque). Such a system $\{U, x^1, x^2, \dots, x^n\}$ is called a *foliation chart*.

A foliation can be given by a partition of V in a family \mathcal{C} of its submanifolds such that each $L \in \mathcal{C}$ is a maximal connected integral submanifold of some integrable subbundle \mathcal{F} of TV. Then \mathcal{C} is the family of leaves of the foliation (V, \mathcal{F}) . Sometimes \mathcal{C} is identified with \mathcal{F} and we will say that (V, \mathcal{F}) is formed by \mathcal{C} .

5. A submanifold N of the foliated manifold V is called a *transversal* iff $T_x V = T_x N \oplus \mathcal{F}_x (\forall x \in N)$. Thus, $\dim N = n - \dim \mathcal{F} = \text{codim} \mathcal{F}$.

A Borel subset B of V such that $B \cap L$ is countable for any leaf L is called a *Borel transversal* to (V, \mathcal{F}) .

A *transverse measure* Λ for the foliation (V, \mathcal{F}) is σ -additive map $B \mapsto \Lambda(B)$ from the set of all Borel transversals to $[0, +\infty]$ such that the following conditions are satisfied:

(i) If $\psi : B_1 \rightarrow B_2$ is a Borel bijection and $\psi(x)$ is on the leaf of any $x \in B_1$, then $\Lambda(B_1) = \Lambda(B_2)$.

(ii) $\Lambda(K) < +\infty$ if K is any compact subset of a smooth transversal submanifold of V , then (V, \mathcal{F}) is called a *measurable foliation*, following A. Connes.

6. Let (V, \mathcal{F}) be a foliation with \mathcal{F} is oriented. Then the complement of zero section of the bundle $\Lambda^k(\mathcal{F})$ ($k = \dim \mathcal{F}$) has two components $\Lambda^k(\mathcal{F})^-$ and $\Lambda^k(\mathcal{F})^+$.

Let μ be a measure on V and $\{U, x^1, x^2, \dots, x^n\}$ be a foliation chart of (V, \mathcal{F}) . Then U can be identified with the direct product $N \times \Pi$ of some smooth transversal submanifold N of V and a some plaque Π . The restriction of μ on $U \equiv N \times \Pi$ becomes the product $\mu_N \times \mu_\Pi$ of measures μ_N and μ_Π respectively.

Let $X \in C^\infty(\Lambda^k(\mathcal{F}))^+$ be a smooth k -vector field and μ_X be the measure on each leaf L determined by the volume element X .

The measure μ is called *X-invariant* iff μ_X is proportional to μ_Π for an arbitrary foliation chart $\{U, x^1, x^2, \dots, x^n\}$.

7. Let $(X, \mu), (Y, \nu)$ be two pairs where $X, Y \in C^\infty(\Lambda^k(\mathcal{F}))^+$ and μ, ν are measures on V such that μ is X -invariant, ν is Y -invariant. $(X, \mu), (Y, \nu)$ are *equivalent* iff $Y = \varphi X$ and $\mu = \varphi\nu$ for some $\varphi \in C^\infty(V)$.

There is a bijective map between the set of transverse measures for (V, \mathcal{F}) and the one of equivalence classes of pairs (X, μ) , where $X \in C^\infty(\Lambda^k(\mathcal{F}))^+$ and μ is a X -invariant measure on V .

Thus, to prove that (V, \mathcal{F}) is measurable, we only need to choose some suitable pair (X, μ) on V (see [1]).

2. Some examples of MD5-algebras and MD5-groups

From now on, G will denote a connected solvable Lie group of dimension 5. The Lie algebra of G is denoted by \mathcal{G} . We always choose a fixed basis (S, T, X, Y, Z) in \mathcal{G} . Then its Lie algebra is isomorphic to \mathbb{R}^5 as a real vector space. The notation \mathcal{G}^* will mean the dual space of \mathcal{G} . Clearly \mathcal{G}^* can be identified with \mathbb{R}^5 by fixing in it the basis $(S^*, T^*, X^*, Y^*, Z^*)$ which is dual to the basis (S, T, X, Y, Z) .

Recall that a group G is called an *MD5-group* if its K-orbits are orbits of dimension zero or maximal dimension. Then its Lie algebra is called an *MD5-algebra*. Note that for any MD4-algebra \mathcal{G}_0 , the direct product $\mathcal{G} = \mathcal{G}_0 \times \mathbb{R}$ with the commutative Lie \mathbb{R} is a MD5-algebra. It is called a *decomposable* MD5-algebra the study of which can be directly reduced to the case of MD4-algebras. Therefore, we will restrict in the case of *indecomposable* MD5-algebras.

2.1 Some Examples of Indecomposable MD5-algebras and MD5-groups

1. Denote by \mathcal{G}_1 the real algebra of dimension 5 with the basis (S, T, X, Y, Z) such that

$$\mathcal{G}_1^1 = [\mathcal{G}_1, \mathcal{G}_1] = \langle Y, Z \rangle \cong \mathbb{R}^2; \text{End}(\mathcal{G}_1^1) \cong \text{Mat}(2, \mathbb{R});$$

$$[S, T] = [T, X] = [X, S] = 0; ad_X = ad_T = 0; ad_S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

The simply connected Lie group associated to \mathcal{G}_1 is denoted by G_1 .

2. Let \mathcal{G}_2 be the real algebra of dimension 5 with the basis (S, T, X, Y, Z) such that

$$\mathcal{G}_2^1 = [\mathcal{G}_2, \mathcal{G}_2] = \langle T, X, Y, Z \rangle \cong \mathbb{R}^4;$$

$$ad_S = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \in Aut(\mathcal{G}_2^1) \cong GL(4, \mathbb{R}).$$

The simply connected Lie group associated to \mathcal{G}_2 is denoted by G_2 .

3. Let \mathcal{G}_3 be the real algebra of dimension 5 with the basis (S, T, X, Y, Z) , such that

$$\mathcal{G}_3^1 = [\mathcal{G}_3, \mathcal{G}_3] = \langle T, X, Y, Z \rangle \cong \mathbb{R}^4;$$

$$ad_S = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \in Aut(\mathcal{G}_3^1) \cong GL(4, \mathbb{R}).$$

The simply connected Lie group associated to \mathcal{G}_3 is denoted by G_3 .

2.2 Remarks

1. The Lie algebras $\mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3$ are the semi-direct products of the form $\mathbb{R} \times_{\varphi} \mathcal{A}$ of the Lie abelian algebra $\mathcal{A} = \langle T, X, Y, Z \rangle \cong \mathbb{R}^4$ with $\mathcal{B} = \langle S \rangle \cong \mathbb{R}$ by the corresponding actions $\varphi = ad_S$.

2. In the next section we shall prove that $\mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3$ are indecomposable MD5-algebras (see Section 3, Corollary 2). Hence, G_1, G_2, G_3 are also MD5-groups.

3. The main results

3.1 The Geometry of K-orbits of G_1, G_2, G_3

Throughout this section, G will denote one of the groups $G_1, G_2, G_3, \mathcal{G}$ for its Lie algebra, $\mathcal{G}^* = \langle S^*, T^*, X^*, Y^*, Z^* \rangle \cong \mathbb{R}^5$ for the dual space of \mathcal{G} , and $F = s_F S^* + t_F T^* + x_F X^* + y_F Y^* + z_F Z^* \equiv (s_F, t_F, x_F, y_F, z_F)$ an arbitrary element of \mathcal{G}^* , and finally Ω_F for the K-orbit of G which contains F .

Theorem 1

$G = G_1$:

1. If $x_F = y_F = 0$ then Ω_F is a K-orbit of dimension zero, i.e.,

$$\Omega_F = \{F(s_F, t_F, 0, 0, z_F)\}.$$

2. If $x_F^2 + y_F^2 \neq 0$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, t_F, x, y, z_F)/x^2 + y^2 = x_F^2 + y_F^2\}$$

(a cylinder of revolution).

$G = G_2$:

1. If $t_F = x_F = y_F = z_F = 0$ then Ω_F is a K-orbit of dimension zero, i.e.,

$$\Omega_F = \{F(s_F, 0, 0, 0, 0)\}.$$

2. If $t_F \neq 0 = x_F = y_F = z_F$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, t, 0, 0, 0)/t_F t > 0\}$$

(a coordinate half - plane).

3. If $x_F \neq 0 = t_F = y_F = z_F$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, 0, x, 0, 0)/x_F x > 0\}$$

(a coordinate half - plane).

4. If $y_F \neq 0 = t_F = x_F = z_F$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, 0, 0, y, 0)/y_F y > 0\}$$

(a coordinate half - plane).

5. If $z_F \neq 0 = t_F = x_F = y_F$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, 0, 0, 0, z)/z_F z > 0\}$$

(a coordinate half - plane).

6. If $t_F x_F \neq 0 = y_F z_F$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, t, x, 0, 0)/x_F t - t_F x = 0, t_F t > 0, x_F x > 0\}$$

(a part of plane).

7. If $t_F y_F \neq 0 = x_F z_F$ then Ω_F is a K -orbit of dimension two as follows

$$\Omega_F = \{(s, t, 0, y, 0)/y_F t - t_F y = 0, t_F t > 0, y_F y > 0\}$$

(a part of plane).

8. If $t_F z_F \neq 0 = x_F y_F$ then Ω_F is a K -orbit of dimension two as follows

$$\Omega_F = \{(s, t, 0, 0, z)/z_F t - t_F z = 0, t_F t > 0, z_F z > 0\}$$

(a part of plane).

9. If $x_F y_F \neq 0 = t_F z_F$ then Ω_F is a K -orbit of dimension two as follows

$$\Omega_F = \{(s, 0, x, y, 0)/y_F x - x_F y = 0, x_F x > 0, y_F y > 0\}$$

(a part of plane).

10. If $x_F z_F \neq 0 = t_F y_F$ then Ω_F is a K -orbit of dimension two as follows

$$\Omega_F = \{(s, 0, x, 0, z)/z_F x - x_F z = 0, x_F x > 0, z_F z > 0\}$$

(a part of plane).

11. If $y_F z_F \neq 0 = t_F x_F$ then Ω_F is a K -orbit of dimension two as follows

$$\Omega_F = \{(s, 0, 0, y, z)/z_F y - y_F z = 0, y_F y > 0, z_F z > 0\}$$

(a part of plane).

12. If $t_F x_F y_F \neq 0 = z_F$ then Ω_F is a K -orbit of dimension two as follows

$$\Omega_F = \{(s, t, x, y, 0)/t_F x - x_F t = 0, t_F y - y_F t = 0, t_F t > 0, x_F x > 0, y_F y > 0\}$$

(a part of plane).

13. If $t_F x_F z_F \neq 0 = y_F$ then Ω_F is a K -orbit of dimension two as follows

$$\Omega_F = \{(s, t, x, 0, z)/t_F x - x_F t = 0, t_F z - z_F t = 0, t_F t > 0, x_F x > 0, z_F z > 0\}$$

(a part of plane).

14. If $t_F y_F z_F \neq 0 = x_F$ then Ω_F is a K -orbit of dimension two as follows

$$\Omega_F = \{(s, t, 0, y, z)/t_F y - y_F t = 0, t_F z - z_F t = 0, t_F t > 0, y_F y > 0, z_F z > 0\}$$

(a part of plane).

15. If $x_F y_F z_F \neq 0 = t_F$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, 0, x, y, z) / x_F y - y_F x = 0, x_F z - z_F x = 0, x_F x > 0, y_F y > 0, z_F z > 0\}$$

(a part of plane).

16. If $t_F x_F y_F z_F \neq 0$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, t, x, y, z) / t_F x - x_F t = 0, \quad t_F y - y_F t = t_F z - z_F t = 0, \\ t_F t > 0, x_F x > 0, y_F y > 0, z_F z > 0\}$$

(a part of plane).

$G = G_3$:

1. If $t_F = x_F = y_F = z_F = 0$ then Ω_F is a K-orbit of dimension zero, i.e.,

$$\Omega_F = \{F(s_F, 0, 0, 0, 0)\}.$$

2. If $x_F = y_F = 0 \neq t_F^2 + z_F^2$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, t, 0, 0, z) / t^2 + z^2 = t_F^2 + z_F^2\}$$

(a cylinder of revolution).

3. If $t_F = z_F = 0 \neq x_F^2 + y_F^2$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, 0, x, y, 0) / x^2 + y^2 = x_F^2 + y_F^2\}$$

(a cylinder of revolution).

4. If $x_F^2 + y_F^2 \neq 0 \neq t_F^2 + z_F^2$ then Ω_F is a K-orbit of dimension two as follows

$$\Omega_F = \{(s, t, x, y, z) / t^2 + z^2 = t_F^2 + z_F^2; x^2 + y^2 = x_F^2 + y_F^2; tx - yz = t_F x_F - y_F z_F\}.$$

Corollary 2

1. G_1, G_2, G_3 are indecomposable MD5-groups.
2. $\mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3$ are indecomposable MD5-algebras.

3. Sketch the proof of Theorem 1

1. For each G , we denote the set $\{F_U \in \mathcal{G}^*/U \in \mathcal{G}\}$ by $\Omega_F(\mathcal{G})$, where F_U is the linear form on the Lie algebra \mathcal{G} of G defined by

$$\langle F_U, A \rangle = \langle F, \exp(ad_U)(A) \rangle, A, U \in \mathcal{G}.$$

At first, we have to compute $\exp(ad_U)$ and define F_U . After that, $\Omega_F(\mathcal{G})$ is described by the same method presented in [6], [8].

2. Note that for $G = G_2$, the map $\exp: \mathcal{G} \rightarrow G$ is surjective (see [3]). Hence, $\Omega_F = \Omega_F(\mathcal{G})$.
3. For $G \in \{G_1, G_3\}$, the equation $\Omega_F = \Omega_F(\mathcal{G})$ is verified by using [10, Lemma II.1.5].

3.2 MD5-foliations associated to G_1, G_2, G_3

Theorem 3 *Let $G \in \{G_1, G_2, G_3\}$, \mathcal{F}_G be the family of all its K -orbits of maximal dimension and $V_G = \bigcup \{\Omega/\Omega \in \mathcal{F}_G\}$. Then (V_G, \mathcal{F}_G) is a measurable foliation in the sense of Connes. We call it MD5-foliation associated to MD5-group G .*

Sketch of the proof of Theorem 3

The proof is analogous to the case of MD4-groups in [6], [8], [10]. First, we need to define a smooth tangent 2-vector field on the manifold V_G such that each K -orbit Ω from \mathcal{F}_G is a maximal connected integrable submanifold corresponding to it. As the next step, we have to show that the Lebesgue measure is invariant for that 2-vector field. The last step is a simple matter and can be verified by direct computations. Now we introduce the smooth tangent 2-vector fields corresponding to each of G from $\{G_1, G_2, G_3\}$.

- $\mathcal{S}_{G_1} = \mathcal{X}_1 \wedge \mathcal{X}_2$ on the foliated manifold $V_G \equiv \mathbb{R}^2 \times (\mathbb{R}^2 \setminus \{O(0,0)\}) \times \mathbb{R}$, where
 1. $\mathcal{X}_1(s, t, x, y, z) = (0, 0, y, -x, 0); \forall (s, t, x, y, z) \in V_G;$
 2. $\mathcal{X}_2(s, t, x, y, z) = (1, 0, 0, 0, 0); \forall (s, t, x, y, z) \in V_G.$
- $\mathcal{S}_{G_2} = \mathcal{X}_1 \wedge \mathcal{X}_2$ on the foliated manifold $V_G \equiv \mathbb{R} \times (\mathbb{R}^4 \setminus \{O(0,0,0,0)\})$, where
 1. $\mathcal{X}_1(s, t, x, y, z) = (0, t, x, y, z); \forall (s, t, x, y, z) \in V_G;$
 2. $\mathcal{X}_2(s, t, x, y, z) = (1, 0, 0, 0, 0); \forall (s, t, x, y, z) \in V_G.$
- $\mathcal{S}_{G_3} = \mathcal{X}_1 \wedge \mathcal{X}_2$ on the foliated manifold $V_G \equiv \mathbb{R} \times (\mathbb{R}^4 \setminus \{O(0,0,0,0)\})$, where

1. $\mathcal{X}_1(s, t, x, y, z) = (0, -z, y, -x, t); \forall (s, t, x, y, z) \in V_G;$
2. $\mathcal{X}_2(s, t, x, y, z) = (1, 0, 0, 0, 0); \forall (s, t, x, y, z) \in V_G.$

Acknowledgement The author would like take this opportunity to thank his teacher, Prof. DSc. Do Ngoc Diep for his excellent advice and support. Thanks are due also to the author's colleagues Nguyen Van Sanh, Tran Dao Dong and Doan The Hieu for their encouragement.

References

- [1] A. Connes, *A Survey of Foliations and Operator Algebras*, Proc. Symp. Pure Math., 38(1982), 521 - 628, Part I.
- [2] A. A. Kirillov, "Elements of the Theory of Representations", Springer - Verlag, Berlin - Heidenberg - New York, 1976.
- [3] M. Saito, *Sur Certains Groupes de Lie Resolubles*, Sci. Papers of the College of General Education, Uni. of Tokyo, 7(1957), 1 -11, 157 - 168.
- [4] V. M. Son et H. H. Viet, *Sur la Structure des C^* -algebres d' une Classe de Groupes de Lie*, J. Operator Theory, 11(1984), 77 - 90.
- [5] D. V. Tra, *On the Lie Algebras of low dimension*, Sci. Papers of the 12th College of Institute of Math. Vietnam, Hanoi 1984 (in Vietnamese).
- [6] Le Anh Vu, *The Foliation Formed by the K -orbits of Maximal Dimension of the Real Diamond Group*, Vietnam J. Math., XV(1987), N^o 3, 7 - 10 (in Vietnamese).
- [7] Le Anh Vu, *On the Structure of the C^* -algebra of the Foliation Formed by the K -orbits of Maximal Dimension of the Real Diamond Group*, J. Operator Theory, 24(1990), 227 - 238.
- [8] Le Anh Vu, *On the Foliations Formed by the Generic K -orbits of the MD_4 -Groups*, Acta Math. Vietnam, N^o 2(1990), 39 - 55.
- [9] Le Anh Vu, *Foliations Formed by Orbits of Maximal Dimension in the Co-adjoint Representation of a Class of Solvable Lie Groups*, Vest. Moscow Uni., Math. Bulletin, Vol. 48(1993), N^o 3, 24 - 27.
- [10] Le Anh Vu, *Foliations Formed by K -orbits of Maximal Dimension of the Class of Lie Groups MD_4* , Ph. D. Dissertation, Institute of Math. Vietnam (1990) (in Vietnamese).