

INVOLUTIVE WEAK GLOBULAR ω -CATEGORIES

Paratat Bejrakarbun^a and Paolo Bertozzini^b

*Department of Mathematics and Statistics,
Faculty of Science and Technology
Thammasat University, Pathumthani 12121, Thailand
email: ^a paratat.bejr@dome.tu.ac.th ^b paolo.th@gmail.com*

Dedicated to the memory of Francis William Lawvere

Abstract

We investigate the notion of *involutive* weak globular ω -categories making use of T.Leinster’s approach: as algebras for the initial contracted globular operad in the bicategory of globular collections induced by the Cartesian monad of the free *involutive* strict ω -category functor on globular ω -sets. An apparently more restrictive notion of involutive weak globular ω -categories as algebras for the initial *operadic-contraction* in the bicategory of globular *contracted-collections* induced by the previous Cartesian monad (where here the operadic multiplications and units satisfy further compatibility axioms with the contractions) is also considered.

Contents

1	Introduction and Motivation	2
2	Preliminaries	5
2.1	Strict Globular ω -categories	5
2.2	Monads in Bicategories and Cartesian Monads	8
2.2.1	Bicategories	8
2.2.2	Monads in a Bicategory	10
2.3	Leinster Weak Globular ω -categories	14

Key words: Higher Category, Involutive Category, Operad, Monad.
2010 AMS Classification: 18N65, 18N70, 18M40, 18N30, 18N99.

2.3.1	Generalized Multicategories and Generalized Operads	14
2.3.2	Contractions and Globular Collections	19
2.3.3	Weak ω -categories	22
2.3.4	Examples	24
3	Involutive Leinster Weak Globular ω-categories	25
3.1	Involutive Strict Globular ω -categories	25
3.2	Involutive Weak Globular ω -categories	31
3.3	Examples	51
4	Outlook	52

1 Introduction and Motivation

Category theory, since its inception in [Eilenberg Mac Lane 1945], has always been evolving in very close connection with algebraic topology. The interlink between these two subjects became even more substantial with the development of higher category theory (among the several resources available, see [Cheng Lauda 2004], [Leinster 2004, pages 19-30] for an introductory discussion, [Baez 1997, Baez Dolan 1995] for original motivations also from physics and the wiki-site <http://ncatlab.org/nlab> for further details).

Attempts to test the architecture of higher category theory within non-commutative topology are still in their prenatal stage (see for example [Bertozzini Conti Lewkeeratiyutkul Suthichitranont 2020] where only some “non-commutative” variants of strict n -categories have been considered).

Non-commutative topology is notoriously dominated by the central role of C^* -algebras as an arena generalizing the well-known Gel’fand-Naïmark duality between commutative unital C^* -algebras and compact Hausdorff topologies. As a first minimal attempt to elaborate categorical environments capable of supporting non-commutative homotopy/cobordism and define weak notions of (higher) C^* -categories, one would like to axiomatize the existence of (higher) involutions, vertically categorifying, in a weak environment, several already known notions of involutive categories (see [Yau 2020] and [Bertozzini Conti Lewkeeratiyutkul Suthichitranont 2020, section 4] for further references).

In our previous joint paper [Bejrakarbun Bertozzini 2017] we have been providing a definition of weak involutive ω -category, as an algebra for the free involutive ω -category monad, in the spirit of J.Penon’s algebraic definition of weak ω -categories [Penon 1999]. Our ideological point of view is to consider involutions not (only) as symmetries of a (higher) categorical structure, but as unary operations on the very same footing of the binary compositions and nullary identities present in ordinary categories.

↪ *The purpose of this work is to produce an algebraic definition of involutive weak globular ω -categories following an operadic definition in the style of [Leinster 2004].*

For now (for simplicity) we limited ourselves to the usual axiomatic setting of (weak) higher categories, although we plan to further develop our work in the direction of non-commutative exchange [Bertozzini Conti Lewkeeratiyutkul Suthichitranont 2020, section 3.3] in view of application to operator algebraic structures.

In the original treatment from T.Leinster, the operadic and contraction structures introduced onto a given globular T -collection are essentially independent; on the other side, terminal objects in the category of globular T -collections seem to be naturally contracted T -operads that further satisfy compatibility axioms between operadic multiplication/unit and contraction.

↪ *We put forward a more restrictive notion of weak (involutive) globular ω -category as an algebra for an initial “operadic contraction”, that is universal among those contracted operads whose multiplication and unit satisfy additional compatibility conditions with the contraction.*

As a first-aid motivation for readers that might not be familiar with the intricacies of operadic definitions of weak ω -categories, we provide here below a brief synopsis of the construction:

- ▶ one first introduces strict (involutive) ω -categories and constructs the Cartesian monad \hat{T} (respectively \hat{T}^* in our involutive case) induced by the free (involutive) ω -category functor,
- ▶ the monad \hat{T} (respectively \hat{T}^*) applied to the the terminal globular ω -set \bullet specifies the input-type “arity” of general operations to be axiomatized via operads,
- ▶ to the Cartesian monad \hat{T} (respectively \hat{T}^*) a bicategory $\mathcal{E}_{\hat{T}}$ (respectively $\mathcal{E}_{\hat{T}^*}$) is associated whose 1-cells $E \xleftarrow{t_M} M \xrightarrow{s_M} \hat{T}(E)$ (respectively $E \xleftarrow{t_M} M \xrightarrow{s_M} \hat{T}^*(E)$) represent systems “labeling the multi-input one-target operations” with source parametrized by $\hat{T}(E)$ (respectively by $\hat{T}^*(E)$) and target in E ,
- ▶ generalized \hat{T} -multicategories (respectively \hat{T}^* -multicategories) are defined as monads in the previous bicategory and generalized \hat{T} -operads are just generalized \hat{T} -multicategories whose labeling is provided by $\hat{T}(\bullet)$ (respectively by $\hat{T}^*(\bullet)$),
- ▶ the actual unbiased description of the evaluation of all the operations involved into the definition of a weak globular ω -category and of their coherence structure are together uniquely specified by a choice of contraction on a \hat{T} -operad (respectively on a \hat{T}^* -operad),

- ▶ the monad L (respectively L^*) that is the initial/universal contracted \hat{T} -operad (respectively contracted \hat{T}^* -operad) is supposed to specify the labeling of operations, in a weak (involutive) globular ω -category, with certain $\hat{T}(\bullet)$ (respectively $\hat{T}^*(\bullet)$) inputs and describe their formal compositions and identities,
- ▶ algebras for the initial contracted \hat{T} -operad L (respectively for initial contracted \hat{T}^* -operad L^*) are the actual weak (involutive) ω -categories,
- ▶ a contracted \hat{T} -operad (respectively contracted \hat{T}^* -operad) P produces a strict functor from algebras over P to algebras over L (respectively over L^*), that by definition are the weak (involutive) ω -categories: to give an example of weak (involutive) ω -category is equivalent to provide an algebra for a contracted \hat{T} -operad (respectively contracted \hat{T}^* -operad) P .

Notice that in our involutive case: the involution is used to specify the input type monad \hat{T}^* , it is not used to compose formal labeled operations and hence (apart from the labeling input $\hat{T}^*(\bullet)$) it does not modify the definition of the monad underlying the definition of initial operad L^* : there is usually no involution on the collection of operations making up L^* (the operad only takes care of the nesting of operations); involutions and their evaluations are instead hidden in the choice of contraction that explicitly depends on the base labeling via $\hat{T}^*(\bullet)$ in place of $\hat{T}(\bullet)$.

The content of the paper consists of this brief motivational introduction section 1 followed by a section 2 of preliminaries, where we recall (in a notation compatible with our previous work) already available material on strict ω -categories, monads in bicategories and T.Leinster's construction of weak ω -categories as algebras for a certain generalized operad.

Section 3 of the paper opens recalling our previously developed definition of *involutive* strict ω -category and continues exposing the new material on an operadic definition of *involutive weak ω -categories* as algebras for a generalized initial \hat{T}^* -operad in the bicategory of \hat{T}^* -collections, where \hat{T}^* is the monad of the free involutive strict ω -category construction presented in [Bejrakarbum Bertozzini 2017, propositions 3.1 and 3.2].

Our main existence theorem 3.12 is obtained from a direct procedure, detailed in theorem 3.16, explicitly constructing by recursion a *free contracted \hat{T}^* -operadic magma* and quotienting it in order to obtain a *free contracted \hat{T}^* -operad* over a \hat{T}^* -collection.

The more restrictive notion of *operadic contraction* (in place of the more general contracted-operads considered in [Leinster 2004, definition 9.2.1]) is introduced in remark 2.22, where we also mention the possibility to utilize them to define a tighter variant of Leinster's algebraic notion of weak globular ω -categories. A parallel treatment of this issue in the involutive case is described in remarks 3.10 and 3.19; the terminal operadic-contraction $\hat{T}^*(\bullet)$ is examined in detail in remark 3.11.

We close the work in section 4 with some outlook on possible further work in the direction of weak higher C^* -categories and higher categorical non-commutative geometry.

2 Preliminaries

This section is dedicated to a description of all the long background material necessary to formulate algebraic operadic notions of weak ω -categories; most of the material is directly inspired by [Leinster 2004].

Before starting, a foundational disclaimer: although no set-theoretical contradiction will emerge in this work, formally (especially in section 2.3.1) we will use monads internal to a bicategory of non-small categories ¹

2.1 Strict Globular ω -categories

We recall the formalism and definition of strict globular ω -categories as used in [Bejrakarbum Bertozzini 2017].

Definition 2.1. An ω -**quiver** $Q^0 \xrightleftharpoons[t^0]{s^0} Q^1 \xrightleftharpoons[t^1]{s^1} \dots \xrightleftharpoons[t^{n-1}]{s^{n-1}} Q^n \xrightleftharpoons[t^n]{s^n} \dots$ consists of a sequence of sets $(Q^k)_{k \in \mathbb{N}}$ equipped with sequences of **source** $(s^k)_{k \in \mathbb{N}}$ and **target** $(t^k)_{k \in \mathbb{N}}$ maps.

A **globular ω -set** is an ω -quiver that satisfies the **globularity conditions**:

$$s^k \circ s^{k+1} = s^k \circ t^{k+1}, \quad t^k \circ s^{k+1} = t^k \circ t^{k+1}, \quad \forall k \in \mathbb{N}.$$

For any $k \in \mathbb{N}$, an element $x \in Q^k$ is called a **globular k -cell** of the globular ω -set.

A globular ω -set is **reflexive** if it is equipped with a sequence $(t^k)_{k \in \mathbb{N}}$ of maps

$$Q^0 \xrightarrow{t^0} Q^1 \xrightarrow{t^1} \dots \xrightarrow{t^{n-1}} Q^n \xrightarrow{t^n} \dots$$

such that $s^k \circ t^k = \text{Id}_{Q^k} = t^k \circ t^k$ for every $k \in \mathbb{N}$.

A (**reflexive**) **globular ω -magma** is a (reflexive) globular ω -set equipped with a family of **compositions**

$$\circ_p^m : Q^m \times_{Q^p} Q^m \rightarrow Q^m, \quad (x', x) \mapsto x' \circ_p^m x, \quad \forall m \in \mathbb{N}_0, \quad 0 \leq p < m,$$

¹A simple solution would be to consider a set theory based on classes of at least “3 types” (2-classes consisting of elements called 1-classes, whose elements are called 0-classes and identified as sets) suitably formulating the axiom of “class formation” in such a way that, for $k \in \{2, 1, 0\}$, “proper classes” of level k cannot be elements of classes of level strictly less than k .

where $\mathcal{Q}^m \times_{\mathcal{Q}^p} \mathcal{Q}^m := \{(x', x) \in \mathcal{Q}^m \times \mathcal{Q}^m \mid t^p \circ t^{p+1} \circ \dots \circ t^{m-1}(x) = s^p \circ s^{p+1} \circ \dots \circ s^{m-1}(x')\}$, such that the following conditions hold: if $m \in \mathbb{N}_0$, $0 \leq p < m$ and $(x', x) \in \mathcal{Q}^m \times_{\mathcal{Q}^p} \mathcal{Q}^m$,

$$\begin{aligned} & (s^q \circ s^{q+1} \circ \dots \circ s^{m-1})(x' \circ_p^m x) \\ &= \begin{cases} (s^q \circ s^{q+1} \circ \dots \circ s^{m-1})(x') \circ_p^q (s^q \circ s^{q+1} \circ \dots \circ s^{m-1})(x), & q > p; \\ (s^q \circ s^{q+1} \circ \dots \circ s^{m-1})(x'), & q \leq p. \end{cases} \end{aligned}$$

$$\begin{aligned} & (t^q \circ t^{q+1} \circ \dots \circ t^{m-1})(x' \circ_p^m x) \\ &= \begin{cases} (t^q \circ t^{q+1} \circ \dots \circ t^{m-1})(x') \circ_p^q (t^q \circ t^{q+1} \circ \dots \circ t^{m-1})(x), & q > p; \\ (t^q \circ t^{q+1} \circ \dots \circ t^{m-1})(x), & q \leq p. \end{cases} \end{aligned}$$

A **strict globular ω -category** $(\mathcal{C}, s, t, \iota, \circ)$ is a reflexive globular ω -magma

$$\begin{array}{c} \mathcal{C}^0 \begin{array}{c} \xleftarrow{s^0} \\ \xrightarrow{t^0} \end{array} \mathcal{C}^1 \begin{array}{c} \xleftarrow{s^1} \\ \xrightarrow{t^1} \end{array} \mathcal{C}^1 \cdots \mathcal{C}^n \begin{array}{c} \xleftarrow{s^n} \\ \xrightarrow{t^n} \end{array} \mathcal{C}^{n+1} \cdots, \\ \mathcal{C}^n \times_{\mathcal{C}^p} \mathcal{C}^n \xrightarrow{\circ_p^n} \mathcal{C}^n, \end{array}$$

that satisfies the following list of algebraic axioms:

- **(associativity)** for all $p, m \in \mathbb{N}$, such that $0 \leq p < m$, and all $x, y, z \in \mathcal{C}^m$ with $(z, y), (y, x) \in \mathcal{C}^m \times_{\mathcal{C}^p} \mathcal{C}^m$:

$$(z \circ_p^m y) \circ_p^m x = z \circ_p^m (y \circ_p^m x),$$

- **(unitality)** for all $p, m \in \mathbb{N}$, such that $0 \leq p < m$, and all $x \in \mathcal{C}^m$:

$$(\iota^{m-1} \circ \dots \circ \iota^p \circ t^p \circ \dots \circ t^{m-1})(x) \circ_p^m x = x = x \circ_p^m (\iota^{m-1} \circ \dots \circ \iota^p \circ s^p \circ \dots \circ s^{m-1})(x),$$

- **(functoriality of identities)** for all $q, p \in \mathbb{N}$, such that $0 \leq q < p$, and all $(x', x) \in \mathcal{C}^p \times_{\mathcal{C}^q} \mathcal{C}^p$:

$$\iota^p(x') \circ_q^{p+1} \iota^p(x) = \iota^p(x' \circ_q^p x),$$

- **(binary exchange)** for all $q, p, m \in \mathbb{N}$ such that $0 \leq q < p < m$ and all $x, x', y, y' \in \mathcal{C}^m$ with $(y', y), (x', x) \in \mathcal{C}^m \times_{\mathcal{C}^p} \mathcal{C}^m$ and $(y', x'), (y, x) \in \mathcal{C}^m \times_{\mathcal{C}^q} \mathcal{C}^m$:

$$(y' \circ_p^m y) \circ_q^m (x' \circ_p^m x) = (y' \circ_q^m x') \circ_p^m (y \circ_q^m x), \quad \bullet \begin{array}{c} \xrightarrow{\Downarrow x} \\ \xrightarrow{\Downarrow x'} \end{array} \bullet \begin{array}{c} \xrightarrow{\Downarrow y} \\ \xrightarrow{\Downarrow y'} \end{array} \bullet.$$

A **morphism of ω -quivers** $Q \xrightarrow{\phi} \hat{Q}$ (respectively, of globular ω -sets) is a sequence $(\phi^k)_{k \in \mathbb{N}}$ of maps $Q^k \xrightarrow{\phi^k} \hat{Q}^k$ that, for any $q \in \mathbb{N}$, satisfies any one of the following two alternative properties:

$$q\text{-covariance} : \quad \hat{s}^q \circ \phi^{q+1} = \phi^q \circ s^q, \quad \hat{t}^q \circ \phi^{q+1} = \phi^q \circ t^q, \quad (2.1)$$

$$q\text{-contravariance} : \quad \hat{t}^q \circ \phi^{q+1} = \phi^q \circ s^q, \quad \hat{s}^q \circ \phi^{q+1} = \phi^q \circ t^q. \quad (2.2)$$

An index $q \in \mathbb{N}$ satisfying (2.1) (respectively (2.2)) is a **ϕ -covariance** (respectively **ϕ -contravariance**) index.

A **morphism of reflexive ω -quivers** (respectively of reflexive globular ω -sets) is also required to satisfy:

$$\forall k \in \mathbb{N} : \hat{t}^k \circ \phi^k = \phi^{k+1} \circ t^k.$$

A **morphism of (reflexive) globular ω -magmas** is a morphism of (reflexive) globular ω -sets that, for all $k, q \in \mathbb{N}$ such that $0 \leq q < k$, further satisfies:

whenever q is ϕ -covariance index:

$$\phi^k(x \circ_q^k x') = \phi^k(x) \hat{\delta}_q^k \phi^k(x'), \quad \forall (x, x') \in Q^k \times_{Q^q} Q^k,$$

whenever q is ϕ -contravariance index:

$$\phi^k(x \circ_q^k x') = \phi^k(x') \hat{\delta}_q^k \phi^k(x), \quad \forall (x, x') \in Q^k \times_{Q^q} Q^k.$$

An **ω -functor** between two strict globular ω -categories is a morphism of their reflexive globular ω -magmas.

Remark 2.2. Each one of the previous notions of “morphism” provides a strict 1-category where, given a “composable pair of morphisms” $Q \xrightarrow{\psi} \hat{Q} \xrightarrow{\phi} \tilde{Q}$, their composition $Q \xrightarrow{\phi \circ \psi} \tilde{Q}$ is defined componentwise: $(\phi^k)_{k \in \mathbb{N}} \circ (\psi^k)_{k \in \mathbb{N}} := (\phi^k \circ \psi^k)_{k \in \mathbb{N}}$; and, for any object $Q := (Q^k)_{k \in \mathbb{N}}$, its “identity morphism” is defined by $\iota(Q) := (\text{Id}_{Q^k})_{k \in \mathbb{N}}$. \square

The following result is well-known, see for example [Penon 1999] or [Leinster 2004, appendix F].

Proposition 2.3. Let \mathcal{Q} denote the strict 1-category of covariant morphisms between globular ω -sets and let \mathcal{C} be the strict 1-category of covariant ω -functors between strict globular ω -categories.

For any globular ω -set Q in \mathcal{Q} , a **free strict globular ω -category over Q** is a morphism of globular ω -sets $Q \xrightarrow{\eta_Q} \mathfrak{U}(\mathcal{C})$, into the underlying globular ω -set $\mathfrak{U}(\mathcal{C})$ of a strict globular ω -category \mathcal{C} , satisfying the following universal factorization property: for any morphism of globular ω -sets $Q \xrightarrow{\phi} \mathfrak{U}(\hat{\mathcal{C}})$, into the underlying globular ω -set $\mathfrak{U}(\hat{\mathcal{C}})$

of a strict globular ω -category $\hat{\mathcal{C}}$, there exists a unique ω -functor $\mathcal{C} \xrightarrow{\hat{\phi}} \hat{\mathcal{C}}$ such that $\phi = \hat{\phi} \circ \eta_Q$.

The **forgetful functor** $\mathcal{C} \xrightarrow{\mathfrak{U}} \mathcal{Q}$ (forgetting compositions and identities of objects in \mathcal{C}) admits a left-adjoint $\mathfrak{F} \dashv \mathfrak{U}$ **free strict globular ω -category functor** $\mathcal{C} \xleftarrow{\mathfrak{F}} \mathcal{Q}$ that is uniquely determined via a specific construction of free strict globular ω -category $Q \xrightarrow{\eta} \mathfrak{U}(\mathcal{C})$ over the globular ω -set Q above.

2.2 Monads in Bicategories and Cartesian Monads

Algebraic definitions of weak ω -categories in the several approaches available [Penon 1999, Batanin 1998, Leinster 1998, Batanin 2022] make use of algebras/modules over certain (generalized) monads.

In this subsection we review the basic preliminaries on bicategories, introducing monads (respectively algebras over them) as internal monoids in a bicategory (respectively modules over such monoids). For completeness, categorical adjunctions and some of their well-known relations to monads in the bicategory of categories are also recalled, following [Riehl 2016]. Finally Leinster's definition of Cartesian monad is presented.

2.2.1 Bicategories

Bicategories [Bénabou 1967], [Borceux 1994, section I.7.7], [Leinster 1998], [Leinster 2004, section I.1.5], are a horizontal categorification of the well-known notion of weak monoidal category (where a monoidal category is just a strict 2-category with one object). There are alternative possible equivalent definitions of this structure, we present here a version that is adapted to our notation for globular ω -set in definition 2.1.

Definition 2.4. A *bicategory* $(\mathcal{B}, \circ, \iota, \alpha, \lambda, \rho)$ is a reflexive globular 2-magma

$$\mathcal{B}^0 \Leftarrow \mathcal{B}^1 \Leftarrow \mathcal{B}^2$$

such that:

- ▶ the 1-magma $\mathcal{B}^1 \Leftarrow \mathcal{B}^2$ is a strict 1-category with the vertical composition \circ_1^2 and vertical identity ι^1 ; ²
- ▶ $(\circ_0^2, \circ_0^1) : \mathcal{B}^2 \times_{\mathcal{B}^0} \mathcal{B}^2 \rightarrow \mathcal{B}^2$ is a covariant 1-functor; ³

that is further equipped with:

²This condition implies that \mathcal{B}^2 , and $\mathcal{B}^2 \times_{\mathcal{B}^0} \mathcal{B}^2$ are both bundles of 1-categories over the product $\mathcal{B}^0 \times \mathcal{B}^0$ of discrete categories (with projections that are 1-functors) and that $(\iota^1 \circ \iota^0) : \mathcal{B}^0 \rightarrow \mathcal{B}^2$ is also a functor, where \mathcal{B}^0 is considered as a discrete category.

³This condition is equivalent to the strict axioms of binary exchange and functoriality of identities.

- *an associator natural isomorphism* $((-\circ_0^2 -)\circ_0^2 -) \xRightarrow{\alpha} (-\circ_0^2 (-\circ_0^2 -))$ *between the functors*

$$(\mathcal{B}^2 \times_{\mathcal{B}^0} \mathcal{B}^2) \times_{\mathcal{B}^0} \mathcal{B}^2 \xrightarrow{(-\circ_0^2 -)\circ_0^2 -} \mathcal{B}^2, \quad \mathcal{B}^2 \times_{\mathcal{B}^0} (\mathcal{B}^2 \times_{\mathcal{B}^0} \mathcal{B}^2) \xrightarrow{-\circ_0^2 (-\circ_0^2 -)} \mathcal{B}^2,$$

over the naturally isomorphic 1-categories

$$(\mathcal{B}^2 \times_{\mathcal{B}^0} \mathcal{B}^2) \times_{\mathcal{B}^0} \mathcal{B}^2 \xrightarrow{\alpha} \mathcal{B}^2 \times_{\mathcal{B}^0} (\mathcal{B}^2 \times_{\mathcal{B}^0} \mathcal{B}^2);$$

- *a right unitor natural isomorphism* $(-\circ_0^2 -) \xRightarrow{\rho} \mathfrak{I}_{\mathcal{B}^2}$ *between the pair of 1-functors*⁴

$$\mathcal{B}^2 \times_{\mathcal{B}^0} \iota(\mathcal{B}^1) \xrightarrow{-\circ_0^2 -} \mathcal{B}^2, \quad \mathcal{B}^2 \xrightarrow{\mathfrak{I}_{\mathcal{B}^2}} \mathcal{B}^2,$$

over the naturally isomorphic 1-categories $\mathcal{B}^2 \times_{\mathcal{B}^0} \iota(\mathcal{B}^1) \xrightarrow{\rho} \mathcal{B}^2$;⁵

- *a left unitor natural isomorphism* $(-\circ_0^2 -) \xRightarrow{\lambda} \mathfrak{I}_{\mathcal{B}^2}$ *between the two 1-functors*

$$\iota(\mathcal{B}^1) \times_{\mathcal{B}^0} \mathcal{B}^2 \xrightarrow{-\circ_0^2 -} \mathcal{B}^2, \quad \mathcal{B}^2 \xrightarrow{\mathfrak{I}_{\mathcal{B}^2}} \mathcal{B}^2,$$

over the naturally isomorphic 1-categories $\iota(\mathcal{B}^1) \times_{\mathcal{B}^0} \mathcal{B}^2 \xrightarrow{\lambda} \mathcal{B}^2$;

such that any possible diagram involving (iterated) applications of the \circ_0^2 composition functor to the associator isomorphism α , the left/right unitor isomorphisms λ/ρ and their inverses, commutes.

Remark 2.5. Coherence theorems for weak monoidal categories and bicategories assure that the condition on the commuting diagrams in the previous definition is satisfied as long as the following pentagonal and triangular diagrams commute under

⁴Here $\mathfrak{I}_{\mathcal{B}}$ denotes the identity functor of the category \mathcal{B} .

⁵Here $\iota(\mathcal{B}^1)$ is a bundle over $\mathcal{B}^0 \times \mathcal{B}^0$ of discrete categories (consisting only of identity morphisms in \mathcal{B}^2); it is a distinguished terminal object in the category of bundles over $\mathcal{B}^0 \times \mathcal{B}^0$ of 1-categories with fiberwise 1-functors as morphisms.

\circ_1^2 -composition:

associator coherence: for all $A \xrightarrow{z} B \xrightarrow{y} C \xrightarrow{x} D \xrightarrow{w} E$ in \mathcal{B}^1

$$\begin{array}{ccc} ((w \circ_0^1 x) \circ_0^1 y) \circ_0^1 z & \xrightarrow{\alpha_{w,x,y} \circ_0^2 \iota^1(z)} & (w \circ_0^1 (x \circ_0^1 y)) \circ_0^1 z \xrightarrow{\alpha_{w,(x \circ_0^1 y),z}} w \circ_0^1 ((x \circ_0^1 y) \circ_0^1 z) \\ \downarrow \alpha_{(w \circ_0^1 x),y,z} & & \downarrow \iota^1(w) \circ_0^2 \alpha_{x,y,z} \\ (w \circ_0^1 x) \circ_0^1 (y \circ_0^1 z) & \xrightarrow{\alpha_{w,x,(y \circ_0^1 z)}} & w \circ_0^1 (x \circ_0^1 (y \circ_0^1 z)) \end{array}$$

unitors coherence: for all $A \xrightarrow{g} B \xrightarrow{f} C$ in \mathcal{B}^1

$$\begin{array}{ccc} (f \circ_0^1 \iota^0(B)) \circ_0^1 g & \xrightarrow{\alpha_{f, \iota^0(B), g}} & f \circ_0^1 (\iota^0(B) \circ_0^1 g) \\ \searrow \rho_f \circ_0^2 \iota^1(g) & & \swarrow \iota^1(f) \circ_0^2 \lambda_g \\ & f \circ_0^1 g & \end{array}$$

We refer to the respective entry [coherence in n-Lab] for references and details about the proof of this result. \square

2.2.2 Monads in a Bicategory

The notion of *monad* (in a strict 2-category) originated in a concrete adjunction case in [Godement 1958], with the name “standard construction”; it is a powerful instrument that allows to generalize algebraic structures.

The abstract notion of formal monad over an object of a strict 2-category and in a bicategory are introduced in [Street 1972] and more recently discussed, for example, in [Chikhladze 2015].

Definition 2.6. Let $(\mathcal{B}, \circ, \iota, \alpha, \lambda, \rho)$ be a bicategory and $\mathbb{B} \in \mathcal{B}^0$ an object of \mathcal{B} .

A **monad** (T, μ, η) over \mathbb{B} consists of a 1-cell $\mathbb{B} \xrightarrow{T} \mathbb{B}$ together with a pair of 2-cells $\mu, \eta \in \mathcal{B}^2$:

► the **monadic multiplication** $T \circ_0^1 T \xrightarrow{\mu} T$,

► the **monadic unit** $\iota^0(\mathbb{B}) \xrightarrow{\eta} T$,

such that the following unitality and associativity diagrams of \circ_1^2 -compositions of

2-cells are commuting:

$$\begin{array}{ccc}
 \iota^0(\mathcal{B}) \circ_0^1 T & \xrightarrow{\eta \circ_0^2 \iota^1(T)} & T \circ_0^1 T \xleftarrow{\iota^1(T) \circ_0^2 \eta} T \circ_0^1 \iota^0(\mathcal{B}) \\
 \lambda \downarrow & & \mu \downarrow \quad \rho \downarrow \\
 T & \xrightarrow{\iota^1(T)} & T \xleftarrow{\iota^1(T)} T
 \end{array}
 \quad (2.3)$$

$$\begin{array}{ccc}
 (T \circ_0^1 T) \circ_0^1 T & \xrightarrow{\alpha} & T \circ_0^1 (T \circ_0^1 T) \xrightarrow{\iota^1(T) \circ_0^2 \mu} T \circ_0^1 T \\
 \mu \circ_0^2 \iota^1(T) \downarrow & & \mu \downarrow \\
 T \circ_0^1 T & \xrightarrow{\mu} & T
 \end{array}$$

Given two objects $\mathcal{A}, \mathcal{B} \in \mathcal{B}^0$ and two monads, (T, μ, η) over \mathcal{A} and (S, μ', η') over \mathcal{B} , a T - S **bimodule** is a 1-cell $\mathcal{A} \xleftarrow{\mathcal{M}} \mathcal{B}$ in \mathcal{B}^1 together with a pair of **left/right evaluations** 2-cells $T \circ_0^1 \mathcal{M} \xrightarrow{\theta} \mathcal{M}$ and $\mathcal{M} \circ_0^1 S \xrightarrow{\vartheta} \mathcal{M}$ such that the following diagrams involving vertical composition of 2-cells all commute:

$$\begin{array}{ccc}
 T \circ_0^1 (T \circ_0^1 \mathcal{M}) & \xleftarrow{\alpha_{T,T,\mathcal{M}}} & (T \circ_0^1 T) \circ_0^1 \mathcal{M} \xrightarrow{\mu \circ_0^2 \iota^1(\mathcal{M})} T \circ_0^1 \mathcal{M} \\
 \iota^1(T) \circ_0^2 \theta \downarrow & & \theta \downarrow \\
 T \circ_0^1 \mathcal{M} & \xrightarrow{\theta} & \mathcal{M}
 \end{array}$$

$$\begin{array}{ccc}
 \iota^0(\mathcal{A}) \circ_0^1 \mathcal{M} & \xrightarrow{\eta \circ_0^2 \iota^1(\mathcal{M})} & T \circ_0^1 \mathcal{M} \\
 \lambda_{\mathcal{A}} \downarrow & & \theta \downarrow \\
 \mathcal{M} & \xrightarrow{\iota^1(\mathcal{M})} & \mathcal{M}
 \end{array}$$

$$\begin{array}{ccc}
 (\mathcal{M} \circ_0^1 S) \circ_0^1 S & \xrightarrow{\alpha_{\mathcal{M},S,S}} & \mathcal{M} \circ_0^1 (S \circ_0^1 S) \xrightarrow{\iota^1(\mathcal{M}) \circ_0^2 \mu'} \mathcal{M} \circ_0^1 S \\
 \vartheta \circ_0^2 \iota^1(S) \downarrow & & \vartheta \downarrow \\
 \mathcal{M} \circ_0^1 S & \xrightarrow{\vartheta} & \mathcal{M}
 \end{array}$$

$$\begin{array}{ccc}
\mathcal{M} \circ_0^1 \iota^0(\mathcal{B}) & \xrightarrow{\iota^1(\mathcal{M}) \circ_0^2 \eta'} & \mathcal{M} \circ_0^1 \mathcal{S} \\
\rho_{\mathcal{B}} \downarrow & & \downarrow \vartheta \\
\mathcal{M} & \xrightarrow{\iota^1(\mathcal{M})} & \mathcal{M}
\end{array}$$

$$\begin{array}{ccccc}
(T \circ_0^1 \mathcal{M}) \circ_0^1 \mathcal{S} & \xrightarrow{\alpha_{T, \mathcal{M}, \mathcal{S}}} & T \circ_0^1 (\mathcal{M} \circ_0^1 \mathcal{S}) & \xrightarrow{\iota^1(T) \circ_0^2 \vartheta} & T \circ_0^1 \mathcal{M} \\
\theta \circ_0^2 \iota^1(\mathcal{S}) \downarrow & & & & \downarrow \theta \\
\mathcal{M} \circ_0^1 \mathcal{S} & \xrightarrow{\quad \quad \quad} & & \xrightarrow{\quad \quad \quad} & \mathcal{M}
\end{array}$$

We will need to apply monads two consecutive times: in the first case it will be a monad \hat{T} , over \mathcal{Q} (respectively \hat{T}^* over \mathcal{Q}^*), in the strict bicategory \mathcal{C} of natural transformations between covariant functors between (not necessarily small) strict 1-categories; in the second case it will be a monad, over a terminal object, in the bicategory $\mathcal{E}_{\hat{T}}$ (respectively $\mathcal{E}_{\hat{T}^*}$) that will be subsequently defined in proposition 2.12.

Remark 2.7. When applied to the 2-category \mathcal{C} of natural transformations between covariant functors between (small) strict 1-categories, the previous definition of monad over an object $\mathcal{C} \in \mathcal{C}^0$ reproduces the traditional monad endofunctor $T \in [\mathcal{C}; \mathcal{C}]$ with its multiplication and unit natural transformations.

Given a small strict 1-category $\mathcal{C} \in \mathcal{C}^0$, a monad (T, μ, η) over \mathcal{C} is an endofunctor $\mathcal{C} \xrightarrow{T} \mathcal{C}$, $T \in \mathcal{C}^1$ equipped with natural transformations $T \circ T \xrightarrow{\mu^T} T$, the monad multiplication $\mu^T \in \mathcal{C}^2$ and $\iota^1(\mathcal{C}) \xrightarrow{\eta^T} T$, the monad unit $\eta^T \in \mathcal{C}^2$, that satisfy, for every object $X \in \mathcal{C}^0$, the following properties:

$$\mu_X^T \circ T^1(\eta_X^T) = \iota_{T^0(X)}^0 = \mu_X^T \circ \eta_{T^0(X)}^T, \quad \mu_X^T \circ T^1(\mu_X^T) = \mu_X^T \circ \mu_{T^0(X)}^T. \quad (2.4)$$

An **algebra for the monad** (T, μ^T, η^T) over the 1-category $(\mathcal{C}, \circ, 1)$ consists of an object $A \in \mathcal{C}^0$ together with an evaluation morphism $T(A) \xrightarrow{\theta^A} A$, $\theta^A \in \mathcal{C}^1$, such that $\theta^A \circ \eta^A = \iota_A^0$ and $\theta^A \circ T^1(\theta^A) = \theta^A \circ \mu^A$.

Considering a functor $\bullet \xrightarrow{A} \mathcal{C}$ from a terminal 1-category $\bullet \in \mathcal{C}^0$, the definition of bimodule left for the monad T over \mathcal{C} and right for the identity endofunctor of \bullet reproduces the usual definition of T -algebra. \lrcorner

We recall these essential properties of monads and adjunctions, see for example [Riehl 2016, chapter 5].

Remark 2.8. Every adjunction $\mathcal{C} \begin{array}{c} \xrightarrow{\mathfrak{U}} \\ \xleftarrow{\mathfrak{Y}} \end{array} \mathcal{D}$, $\mathfrak{Y} \dashv \mathfrak{U}$, between small 1-categories \mathcal{D}, \mathcal{C} , with unit $\text{Id}_{\mathcal{D}} \xrightarrow{\eta} \mathfrak{U} \circ \mathfrak{Y}$ and co-unit $\mathfrak{Y} \circ \mathfrak{U} \xrightarrow{\epsilon} \text{Id}_{\mathcal{C}}$, induces a monad $T := \mathfrak{U} \circ \mathfrak{Y}$, in

the strict 2-category of natural transformations between functors, over the 1-category \mathcal{Q} , with multiplication $\mu := \mathfrak{U} \circ \epsilon \circ \mathfrak{F}$ and unit η (see for example [Riehl 2016, lemma 5.1.3] for further details).

Every monad (T, μ, η) on a category \mathcal{C} , induces an adjunction $\mathcal{C}^T \begin{array}{c} \xrightarrow{\mathfrak{U}^T} \\ \xleftarrow{\mathfrak{F}^T} \end{array} \mathcal{C}$ where \mathfrak{F}^T

is the free T -algebra functor, left-adjoint to the forgetful functor \mathfrak{U}^T defined on the category \mathcal{C}^T of T -algebras. The monad T coincides with the monad induced by the above adjunction $\mathfrak{F}^T \dashv \mathfrak{U}^T$ (see for example [Riehl 2016, lemma 5.2.8]).

Given a monad (T, μ, η) over \mathcal{C} , the adjunction $\mathcal{C}^T \begin{array}{c} \xrightarrow{\mathfrak{U}^T} \\ \xleftarrow{\mathfrak{F}^T} \end{array} \mathcal{C}$ is a terminal object

(see [Riehl 2016, proposition 5.2.12]) in the category whose objects are adjunctions

$\mathcal{D} \begin{array}{c} \xrightarrow{\mathfrak{U}} \\ \xleftarrow{\mathfrak{F}} \end{array} \mathcal{C}$, $\mathfrak{F} \dashv \mathfrak{U}$, over \mathcal{C} and whose morphisms $\mathcal{C} \begin{array}{c} \xrightarrow{\mathfrak{F}} \\ \xleftarrow{\mathfrak{U}} \end{array} \mathcal{D} \xrightarrow{\mathfrak{S}} \hat{\mathcal{D}} \begin{array}{c} \xrightarrow{\hat{\mathfrak{U}}} \\ \xleftarrow{\hat{\mathfrak{F}}} \end{array} \mathcal{C}$ are

functors $\mathcal{D} \xrightarrow{\mathfrak{S}} \hat{\mathcal{D}}$ such that $\hat{\mathfrak{F}} = \mathfrak{S} \circ \mathfrak{F}$ and $\hat{\mathfrak{U}} \circ \mathfrak{S} = \mathfrak{U}$.

A **monadic adjunction** $\mathfrak{F} \dashv \mathfrak{U}$ is an adjunction $\mathcal{D} \begin{array}{c} \xrightarrow{\mathfrak{U}} \\ \xleftarrow{\mathfrak{F}} \end{array} \mathcal{C}$ such that the terminal

morphism $\mathcal{D} \xrightarrow{\mathfrak{U}} \mathcal{C}^T$, in the previous category of adjunctions over \mathcal{C} , is an equivalence of categories (see [Riehl 2016, definition 5.3.1]). A **monadic functor** is a functor $\mathcal{D} \xrightarrow{\mathfrak{U}} \mathcal{C}$ with a left adjoint $\mathcal{D} \xleftarrow{\mathfrak{F}} \mathcal{C}$ such that the adjunction $\mathfrak{F} \dashv \mathfrak{U}$ is monadic. \square

From the adjunctions described in the previous proposition 2.3 we have the following.

Corollary 2.9. *On the 1-category \mathcal{Q} of morphisms of globular ω -sets, we have the following*

- *free strict globular ω -category monad $\hat{T} := \mathfrak{U} \circ \mathfrak{F}$.*

Certain conditions of “closure under pull-backs” are required to define Leinster’s generalized operads.

Definition 2.10. [Leinster 2004, definition II.4.4.1] A category \mathcal{E} is **Cartesian** if it admits all pull-backs: for any co-span $A \xrightarrow{\alpha} X \xleftarrow{\beta} B$ in \mathcal{E} , there exists a span $A \xleftarrow{\hat{\alpha}} \hat{X} \xrightarrow{\hat{\beta}} B$ in \mathcal{E} , with $\alpha \circ \hat{\alpha} = \beta \circ \hat{\beta}$, satisfying the following universal factorization property: for any other span $A \xleftarrow{\alpha'} P \xrightarrow{\beta'} B$ in \mathcal{E} , with $\alpha \circ \alpha' = \beta \circ \beta'$, there exists a unique $P \xrightarrow{\gamma} \hat{X}$ such that $\alpha' = \hat{\alpha} \circ \gamma$ and $\beta' = \hat{\beta} \circ \gamma$.

A functor $\mathcal{E} \xrightarrow{\mathfrak{G}} \hat{\mathcal{E}}$ between 1-categories is a **Cartesian functor** if it preserves pull-backs.

A natural transformation $\mathcal{E} \begin{array}{c} \xrightarrow{\mathfrak{G}} \\ \Downarrow \zeta \\ \xrightarrow{\mathfrak{G}} \end{array} \hat{\mathcal{E}}$ is a **Cartesian natural transformation** if, for

any 1-arrow $A \xrightarrow{x} B$ in \mathcal{E} , the span $\mathfrak{G}(B) \xleftarrow{\mathfrak{G}(x)} \mathfrak{G}(A) \xrightarrow{\zeta_A} \hat{\mathfrak{G}}(A)$ in $\hat{\mathcal{E}}$ is a pull-back of the co-span $\mathfrak{G}(B) \xrightarrow{\zeta_B} \hat{\mathfrak{G}}(B) \xleftarrow{\hat{\mathfrak{G}}(x)} \hat{\mathfrak{G}}(A)$ in $\hat{\mathcal{E}}$.

A monad (T, μ, η) on a category \mathcal{C} is a **Cartesian monad** if the category \mathcal{C} the functor T and the two natural transformations μ and η are all Cartesian in the previously defined senses.

For the free strict globular ω -category adjunction in proposition 2.3 and its associated monad in corollary 2.9 we have these Cartesianity conditions as a consequence of [Leinster 2004, theorem F.2.2].

Proposition 2.11. *The 1-category \mathcal{Q} of small globular ω -sets with morphisms of globular ω -sets is Cartesian. The 1-category \mathcal{C} of small strict globular ω -categories with ω -functors is Cartesian. The forgetful 1-functor $\mathcal{C} \xrightarrow{\mathfrak{U}} \mathcal{Q}$ and the free strict globular ω -category 1-functor $\mathcal{C} \xleftarrow{\mathfrak{F}} \mathcal{Q}$ are Cartesian. The unit $\mathfrak{S}_{\mathcal{Q}} \xrightarrow{\eta} \mathfrak{U} \circ \mathfrak{F}$ of the free strict globular ω -category functor is a Cartesian natural transformation. The free strict globular ω -category monad $\hat{T} := \mathfrak{U} \circ \mathfrak{F}$ is Cartesian (this means that also the multiplication natural transformation $\hat{T} \circ \hat{T} \xrightarrow{\mu} \hat{T}$ is Cartesian).*

2.3 Leinster Weak Globular ω -categories

Weak ω -categories in the Leinster's approach [Leinster 2004, section III.9.2] are *algebras for a certain monad L* ; the monad L here utilized is a certain *universal (initial) generalized contracted T -operad*, where T is the Cartesian free strict globular ω -category monad \hat{T} of corollary 2.9.

We proceed here to recall all the essential above-mentioned ingredients that are still missing.

2.3.1 Generalized Multicategories and Generalized Operads

A monad T on a Cartesian category \mathcal{E} allows to formulate a generalized notion of multicategory, where the ‘‘arity’’ of the multicategory arrows is specified by $T(\bullet)$, with \bullet an object in \mathcal{E} . The strategy behind such definition originates in [Burrioni 1971], [Hermida 1997] and has been further described in [Leinster 2004, section II.4] whose exposition we are closely following.

Proposition 2.12. [Leinster 2004, section II.4.2]. *Let T be a Cartesian monad on the Cartesian category \mathcal{E} .*

Given a span $E_1 \xrightarrow{i_1} X \xleftarrow{i_2} E_2$ in \mathcal{E} , let us denote by $E_1 \xleftarrow{\pi_1} E_1 \diamond_X E_2 \xrightarrow{\pi_2} E_2$ a choice of pull-back.⁶

There is a bicategory $(\mathcal{E}_T, \circ_T, \iota_T, \alpha_T, \lambda_T, \rho_T)$ defined as follows:

- ▶ 0-cells in \mathcal{E}_T^0 are just objects E in \mathcal{E}^0 ,
- ▶ 1-cells $E_2 \xleftarrow{(t_P, P, s_P)} E_1$ in \mathcal{E}_T^1 are spans in \mathcal{E} of the form: $E_2 \xleftarrow{t_P} P \xrightarrow{s_P} T^0(E_1)$
- ▶ 2-cells $E_2 \begin{array}{c} \xleftarrow{(t_P, P, s_P)} \\ \Downarrow \phi \\ \xleftarrow{(t_Q, Q, s_Q)} \end{array} E_1$ consist of commuting diagrams in \mathcal{E} of the form:

$$\begin{array}{ccc}
 & P & \\
 t_P \swarrow & & \searrow s_P \\
 E_2 & & T^0(E_1) \\
 & \phi \downarrow & \\
 & Q & \\
 t_Q \swarrow & & \searrow s_Q
 \end{array}$$

- ▶ vertical composition \circ_1^2 is just the usual composition of 1-arrows in \mathcal{E} ,
- ▶ vertical identities $\iota_{\mathcal{E}_T}^1 \left(E_2 \xleftarrow{t_P} P \xrightarrow{s_P} T(E_1) \right)$ are just identities $\iota_{\mathcal{E}}^0(P)$ in \mathcal{E} ,
- ▶ horizontal composition $\left(E_1 \xleftarrow{t_{P_1}} P_1 \xrightarrow{s_{P_1}} T(E_2) \right) \circ_0^1 \left(E_2 \xleftarrow{t_{P_2}} P_2 \xrightarrow{s_{P_2}} T(E_3) \right)$ of one arrows in \mathcal{E}_T is given by:

$$\left(E_1 \xleftarrow{t_{P_1} \circ \pi_1} P_1 \diamond_{T(E_2)} T(P_2) \xrightarrow{\mu_{E_3}^T \circ T(s_{P_2}) \circ \pi_2} T(E_3) \right),$$

as specified by the following diagram

$$\begin{array}{ccccccc}
 & & P_1 \diamond_{T(E_2)} T(P_2) & & & & T(E_3) \\
 & & \pi_1 \swarrow & & \searrow \pi_2 & & \uparrow \mu_{E_3}^T \\
 E_1 & \xleftarrow{t_{P_1}} & P_1 & \xrightarrow{s_{P_1}} & T(E_2) & \xleftarrow{T(t_{P_2})} & T(P_2) & \xrightarrow{T(s_{P_2})} & T(T(E_3))
 \end{array}$$

⁶Assume that a specific choice of pull-backs has been done via axiom of choice, for example we will use $E_1 \diamond_X E_2 := E_1 \times_X E_2$.

horizontal composition of 2-cells $E_1 \xrightarrow{(t_{P_1}, P_1, s_{P_1})} E_2 \xrightarrow{(t_{P_2}, P_2, s_{P_2})} E_3$ is given by the 2-cell

$E_1 \xrightarrow{(t_{Q_1}, Q_1, s_{Q_1}) \circ_0^1 (t_{P_2}, P_2, s_{P_2})} E_3$, where $\phi_1 \circ_0^2 \phi_2$ is defined as the unique morphism $P_1 \diamond_{T(E_2)}$

$T(P_2) \xrightarrow{\phi_1 \circ_0^2 \phi_2} Q_1 \diamond_{T(E_2)} T(Q_2)$ in \mathcal{E} induced by the universal factorization property of the pull-back $Q_1 \xleftarrow{\pi'_1} Q_1 \diamond_{T(E_2)} T(Q_2) \xrightarrow{\pi'_2} T(Q_2)$ via the span

$$Q_1 \xleftarrow{\phi_1 \circ \pi_1} P_1 \diamond_{T(E_2)} T(P_2) \xrightarrow{T(\phi_2) \circ \pi_2} T(Q_2)$$

as specified in this commuting diagram:

$$\begin{array}{ccccc}
 & & P_1 & \xrightarrow{\phi_1} & Q_1 \\
 & \nearrow \pi_1 & & & \nwarrow \pi'_1 \\
 P_1 \diamond_{T(E_2)} T(P_2) & & T(E_2) & & Q_1 \diamond_{T(E_2)} T(Q_2) \\
 & \searrow \pi_2 & \nearrow T(t_{P_2}) & \nwarrow T(t_{Q_2}) & \\
 & & T(P_2) & \xrightarrow{T(\phi_2)} & T(Q_2)
 \end{array}$$

► horizontal identities $\iota_T^0(E) := \left(E \xleftarrow{\iota^0(E)} E \xrightarrow{\eta_E^T} T(E) \right)$, for all objects $E \in \mathcal{E}_T^0$.

► left-unitors $E_2 \xrightarrow{\iota^1(E_2) \circ_0^1 P} E_1$ are uniquely determined by universal factorization property of the pull-backs:

$$\begin{array}{ccc}
 E_2 & \xrightarrow{\eta_{E_2}^T} & T(E_2) \xleftarrow{T(t_P)} T(P) \\
 & \searrow t_P & \nearrow \eta_P^T \\
 & & P \\
 & \nearrow \pi_1 & \searrow \pi_2 \\
 & & E_2 \diamond_{T(E_2)} T(P)
 \end{array}
 \quad
 \begin{array}{ccc}
 P & \xrightarrow{t_P} & E_2 \\
 \eta_P^T \downarrow & & \downarrow \eta_{E_2}^T \\
 T(P) & \xrightarrow{T(t_P)} & T(E_2)
 \end{array}$$

(2.5)

(where the second pull-back-diagram above is assured by Cartesianity of η^T and by the diagram

$$\begin{array}{ccccc}
 T(P) & \xrightarrow{T(s_P)} & T^2(E_1) & \xrightarrow{\mu_{E_1}^T} & T(E_1) \\
 \eta_P^T \uparrow & & \eta_{T(E_1)}^T \uparrow & \nearrow \iota^1(T(E_1)) & \\
 P & \xrightarrow{s_P} & T(E_1) & &
 \end{array}$$

that commutes by Cartesianity of η^T and the unital property of the monad T ;

- right-unitors $E_2 \xleftarrow{P \circ_0^1 \iota^1(E_1)} E_1$ are also determined via the universal factorization property of the pull-backs:

$$\begin{array}{ccc}
 P & \xrightarrow{s_P} & T(E_1) \xleftarrow{\iota^1(T(E_1))} T(E_1) \\
 \downarrow \iota^1(P) & & \downarrow \iota^1(T(E_1)) \\
 P & \xrightarrow{T(s_P)} & T(E_1) \\
 \pi_1 \swarrow & \nearrow \pi_2 & \\
 P \circ_{T(E_1)} T(E_1) & &
 \end{array}
 \tag{2.6}$$

(where the second pull-back diagram is due to Cartesianity of T) and by the commuting diagram

$$\begin{array}{ccccc}
 P & \xrightarrow{s_P} & T(E_1) & \xrightarrow{T(\eta_{E_1}^T)} & T^2(E_1) \\
 \searrow s_P & & \downarrow \iota^1(T(E_1)) & \swarrow \mu_{E_1}^T & \\
 & & T(E_1) & &
 \end{array}$$

that is again commuting because of the unital property of the monad T ;

- associators $E_4 \xleftarrow{(P_1 \circ_0^1 P_2) \circ_0^1 P_3} E_1$ for the compositions of

$$E_4 \xleftarrow{P_1} E_3 \xleftarrow{P_2} E_2 \xleftarrow{P_3} E_1$$

are uniquely determined by universal factorization property of the following pair of pull-backs over $T(E_2)$:⁷

$$\begin{array}{ccccc}
 P_1 \diamond T(P_2) & \xrightarrow{\mu_{E_2}^T \circ T(s_{P_2}) \circ \pi_2} & T(E_2) & \xleftarrow{T(t_{P_3})} & T(P_3) \\
 & \swarrow \pi_2 \circ \theta & & \searrow \mu_{P_3}^T \circ T(\pi_1) \circ \pi_1 & \\
 & & P_1 \diamond T(T(P_2) \diamond T(P_3)) & & \\
 & \swarrow \pi_1 & \uparrow \alpha_{P_1, P_2, P_3} & \searrow \pi_2 & \\
 & & (P_1 \diamond T(P_2)) \diamond T(P_3) & &
 \end{array}
 \tag{2.7}$$

where the morphism θ is uniquely determined by universal factorization property in the following diagram, where all the squares (due to composition or to the Cartesianity of T and μ^T) are pull-backs:

$$\begin{array}{ccccccc}
 P_1 \diamond T(P_2 \diamond T(P_3)) & \xrightarrow{\pi_2} & T(P_2 \diamond T(P_3)) & \xrightarrow{T(\pi_2)} & T^2(P_3) & \xrightarrow{\mu_{P_3}^T} & T(P_3) \\
 \downarrow \theta & & \downarrow T(\pi_1) & & \downarrow T^2(t_{P_3}) & & \downarrow T(t_{P_3}) \\
 P_1 \diamond T(P_2) & \xrightarrow{\pi_2} & T(P_2) & \xrightarrow{T(s_{P_2})} & T^2(E_2) & \xrightarrow{\mu_{E_2}^T} & T(E_2) \\
 \downarrow \pi_1 & & \downarrow T(t_{P_2}) & & & & \\
 P_1 & \xrightarrow{s_{P_1}} & T(E_3) & & & &
 \end{array}$$

and where the following triangles diagrams, with target E_4 and source $T(E_1)$

$$\begin{array}{ccccccc}
 P_1 \diamond T(T(P_2) \diamond T(P_3)) & \xrightarrow{T(\pi_2) \circ \pi_2} & T^2(P_3) & \xrightarrow{T^2(s_{P_3})} & T^3(E_1) & \xrightarrow{T(\mu_{E_1}^T)} & T^2(E_1) \\
 \downarrow t_{P_1} \circ \pi_1 & & \downarrow \mu_{P_3}^T & & \downarrow \mu_{T(E_1)}^T & & \downarrow \mu_{E_1}^T \\
 E_4 & \xrightarrow{\alpha_{P_1, P_2, P_3}} & T(P_3) & \xrightarrow{T(s_{P_3})} & T^2(E_1) & \xrightarrow{\mu_{E_3}^T} & T(E_1) \\
 \downarrow t_{P_1} \circ \pi_1 \circ \pi_1 & & \swarrow \pi_2 & & & & \\
 (P_1 \diamond T(P_2)) \diamond T(P_3) & & & & & &
 \end{array}
 \tag{2.8}$$

are commuting because of Cartesianity of T and μ^T and by the associativity property of the monad T .

⁷Here, with a little abuse of notation, we utilize the same symbols π_1, π_2 to denote all the pull-back projections of compositions.

Definition 2.13. Given a Cartesian monad (T, μ^T, η^T) on a Cartesian category \mathcal{E} , a **T -multicategory on E** is a monad (P, μ_P, η_P) over an object E in the bicategory \mathcal{E}_T defined in proposition 2.12.

A **T -operad (P, μ_P, η_P)** is a T -multicategory on a terminal object \bullet in \mathcal{E} .

Remark 2.14. In practice a T -multicategory consists of: an object $E \in \mathcal{E}^0$ (indexing the inputs and outputs of multicategory arrows); a 1-cell $E \xleftarrow{t^P} P \xrightarrow{s^P} T^0(E)$ in \mathcal{E}_T (specifying all the multicategory arrows with multi-input as an element of $T^0(E)$ and only one target in E); a composition of multicategory arrows specified by the monadic multiplication μ_P ; and multicategory identity, specified by the monadic unit η_P . \dashv

Proposition 2.15. For any object $E \in \mathcal{E}$, we have a 1-category, denoted by \mathcal{E}_E^T , whose objects are 1-arrows in \mathcal{E}_T with source and target E and whose morphisms are 2-arrows in \mathcal{E}_T .

For every object $E \in \mathcal{E}$, there is a **category \mathcal{C}_E^T of T -multicategories on E** that is the subcategory of \mathcal{E}_E^T with:

- ▶ objects of \mathcal{C}_E^T are T -multicategories (P, μ_P, η_P) in \mathcal{E}_T ,
- ▶ morphisms $(P, \mu_P, \eta_P) \xrightarrow{\phi} (Q, \mu_Q, \eta_Q)$ in \mathcal{C}_E^T are morphisms in \mathcal{E}_T such that:

$$\begin{array}{ccc}
 P \circ_0^1 P & \xrightarrow{\mu_P} & P \\
 \phi \circ_0^2 \phi \downarrow & & \downarrow \phi \\
 Q \circ_0^1 Q & \xrightarrow{\mu_Q} & Q
 \end{array}
 \qquad
 \begin{array}{ccc}
 \iota_T^0(E) & \xrightarrow{\eta_P} & P \\
 \iota^1(E) \downarrow & & \downarrow \phi \\
 \iota_T^0(E) & \xrightarrow{\eta_Q} & Q
 \end{array}$$

- ▶ composition and identity coincide with those in \mathcal{E}_T .

Definition 2.16. For a terminal object \bullet in a Cartesian category \mathcal{E} with a Cartesian monad T we denote by:

- ▶ \mathcal{E}_\bullet^T the **category of T -collections over \bullet**
- ▶ $\mathcal{O}_\bullet^T := \mathcal{C}_\bullet^T$ the **category of T -operads over \bullet** .

2.3.2 Contractions and Globular Collections

We now specialize the discussion to the bicategory $\mathcal{Q}_{\hat{T}}$ constructed from the Cartesian category $\mathcal{E} := \mathcal{Q}$ of globular ω -sets equipped with the free strict globular ω -category Cartesian monad \hat{T} .

The codification of the algebraic axioms for weak globular ω -categories via ‘‘contractions’’ originates in [Penon 1999]; the following notion of contraction, for globular ω -sets, appears in [Leinster 2001, section II.5] and [Leinster 2004, section III.9.1] and is actually used to formalize, at the same time, the (evaluation of) operations and the algebraic and coherence axioms for weak globular ω -categories.

Definition 2.17. Let $Q \in \mathcal{Q}^0$ be a globular ω -set; we say that $x_1, x_2 \in Q^k$ are **parallel k -cells** if either $k = 0$ or

$$s^{k-1}(x_1) = s^{k-1}(x_2), \quad t^{k-1}(x_1) = t^{k-1}(x_2).$$

We denote by Par_Q the family of pairs of parallel cells of the globular ω -set Q .

Let $Q_1 \xrightarrow{\pi} Q_2$ be a covariant morphism in the category \mathcal{Q} of globular ω -sets, define

$$\text{Par}(\pi) := \left\{ (x^+, y, x^-) \mid (x^+, x^-) \in \text{Par}_{Q_1}, y \in Q_2^n, n \in \mathbb{N}_0, \right. \\ \left. \pi(x^+) = t^{n-1}(y), s^{n-1}(y) = \pi(x^-) \right\}.$$

A **Leinster contraction on π** is a map $\text{Par}(\pi) \xrightarrow{\kappa} Q_1$ such that, $\forall (x^+, y, x^-) \in \text{Par}(\pi)$:

$$s(\kappa(x^+, y, x^-)) = x^-, \quad t(\kappa(x^+, y, x^-)) = x^+, \quad \pi(\kappa(x^+, y, x^-)) = y. \quad (2.9)$$

Definition 2.18. Let $\bullet \in \mathcal{Q}$ denote a terminal object in the category of globular ω -sets and T a monad on \mathcal{Q} .

A **globular T -collection** is a morphism $Q \xrightarrow{\pi} T(\bullet)$ in \mathcal{Q} ; a **globular contracted T -collection** consists of a Leinster contraction $\text{Par}(\pi) \xrightarrow{\kappa} Q$ on a globular T -collection:

$$\text{Par}(\pi) \xrightarrow{\kappa} Q \xrightarrow{\pi} T(\bullet).$$

Lemma 2.19. Given a commuting diagram of covariant functors in the category \mathcal{Q} of globular ω -sets:

$$\begin{array}{ccc} Q_1 & \xrightarrow{\phi} & \hat{Q}_1 \\ & \searrow \pi_1 & \swarrow \pi_2 \\ & Q_2 & \end{array}$$

there is a well-defined induced map $\text{Par}(\pi_1) \xrightarrow{\text{Par}_\phi} \text{Par}(\pi_2)$ given by

$$\text{Par}_\phi : (x^+, y, x^-) \mapsto (\phi(x^+), y, \phi(x^-)).$$

We recall here the notion of category of globular contracted operads from [Leinster 2004, definition III.9.2.3].

Proposition 2.20. For any terminal object $\bullet \in \mathcal{Q}^0$, we have the following categories:

- the category $\mathcal{Q}_\bullet^{\hat{T}}$ of globular \hat{T} -collections over \bullet , specified in definition 2.16;

- the category $\mathcal{Q}_{\bullet}^{\hat{T}, \kappa}$ of globular contracted \hat{T} -collections over \bullet whose objects are globular contracted \hat{T} -collections and whose morphisms

$$(P, \pi_P, \kappa_P) \xrightarrow{\phi} (Q, \pi_Q, \kappa_Q)$$

are globular ω -functors $P \xrightarrow{\phi} Q$ such that

$$\begin{array}{ccc} \text{Par}(\pi_P) & \xrightarrow{\text{Par}_\phi} & \text{Par}(\pi_Q) \\ \kappa_P \downarrow & & \downarrow \kappa_Q \\ P & \xrightarrow{\phi} & Q \end{array} \quad \phi \circ \kappa_P = \kappa_Q \circ \text{Par}_\phi;$$

- the category $\mathcal{O}_{\bullet}^{\hat{T}}$ of globular \hat{T} -operads over \bullet , specified as in definition 2.16;
- the category $\mathcal{O}_{\bullet}^{\hat{T}, \kappa}$ of globular contracted \hat{T} -operads over \bullet with objects consisting of $(P, \pi_P, \kappa_P, \mu_P, \eta_P)$ such that $(P, \pi_P, \mu_P, \eta_P) \in \mathcal{O}_{\bullet}^{\hat{T}}$ is a globular \hat{T} -operad and $(P, \pi_P, \kappa_P) \in \mathcal{Q}_{\bullet}^{\hat{T}}$ is a globular contracted \hat{T} -collection,⁸ and with morphisms $(P, \pi_P, \kappa_P, \mu_P, \eta_P) \xrightarrow{\phi} (Q, \pi_Q, \kappa_Q, \mu_Q, \eta_Q)$ such that

- $(P, \pi_P, \mu_P, \eta_P) \xrightarrow{\phi} (Q, \pi_Q, \mu_Q, \eta_Q)$ is a morphism in $\mathcal{O}_{\bullet}^{\hat{T}}$
- $(P, \pi_P, \kappa_P) \xrightarrow{\phi} (Q, \pi_Q, \kappa_Q)$ is a morphism in $\mathcal{Q}_{\bullet}^{\hat{T}, \kappa}$.

There is a forgetful functor $\mathcal{Q}_{\bullet}^{\hat{T}, \kappa} \xrightarrow{\mathfrak{B}} \mathcal{Q}_{\bullet}^{\hat{T}}$ defined on object by $(Q, \pi, \kappa) \mapsto (Q, \pi)$, for every globular contracted \hat{T} -collection (Q, π, κ) and as identity map on morphisms.

There is a forgetful functor $\mathcal{O}_{\bullet}^{\hat{T}, \kappa} \xrightarrow{\mathfrak{U}} \mathcal{O}_{\bullet}^{\hat{T}}$ defined by $(P, \pi, \kappa, \mu, \eta) \mapsto (P, \pi, \mu, \eta)$ on objects, for every globular contracted \hat{T} -operad $(P, \pi, \kappa, \mu, \eta)$ and as identity map on morphisms.

Remark 2.21. Although T -operads are defined for any operad T on a Cartesian category \mathcal{C} , Leinster's contractions are defined only on globular T -collections, where T is an arbitrary monad on the Cartesian category \mathcal{Q} of globular ω -sets. As a consequence, proposition 2.20 holds actually for any Cartesian operad T on the Cartesian category of globular ω -sets \mathcal{Q} . We will use this fact, substituting the free strict globular ω -category monad \hat{T} with the free involutive strict globular ω -category monad \hat{T}^* in the next section 3.2. \lrcorner

⁸ Notice that at this point we are not imposing any compatibility axiomatic requirement between the contraction κ and the operadic unit η and multiplication μ simultaneously defined on a \hat{T} -collection; we will address this issue in remark 2.22 here below.

Remark 2.22. As anticipated in footnote 8, following the treatment in [Leinster 2004, definition 9.2.1], we did not include in our definition of globular contracted \hat{T} -operad, in the last point of proposition 2.20, any further compatibility axioms between contractions and operadic units/multiplications.

We introduce here a more restrictive notion of **globular \hat{T} -operadic contraction** over \bullet , consisting of data $(P, \pi_P, \kappa_P, \eta_P, \mu_P)$ such that (P, π_P, κ_P) is a globular contracted \hat{T} -collection in $\mathcal{Q}_{\bullet}^{\hat{T}, \kappa}$ and $(P, \pi_P, \eta_P, \mu_P)$ is a globular \hat{T} -operad in $\mathcal{O}_{\bullet}^{\hat{T}}$ that furthermore satisfy the commutativity of the following diagrams in $\mathcal{Q}_{\bullet}^{\hat{T}}$:

$$\begin{array}{ccc} \text{Par}(\pi_{\bullet}) & \xrightarrow{\text{Par}_{\eta_P}} & \text{Par}(\pi_P) \\ \kappa_{\bullet} \downarrow & & \downarrow \kappa_P \\ \bullet & \xrightarrow{\eta_P} & P \end{array} \quad \begin{array}{ccc} \text{Par}(\pi_P \circ_0^2 \pi_P) & \xrightarrow{\text{Par}_{\mu_P}} & \text{Par}(\pi_P) \\ \kappa_P \circ_0^2 \kappa_P \downarrow & & \downarrow \kappa_P \\ P \circ_0^1 P & \xrightarrow{\mu_P} & P \end{array}$$

where $\bullet \xrightarrow{\pi_{\bullet}} \hat{T}(\bullet)$ is $\pi_{\bullet} := \eta_{\bullet}^T$, we have $P \circ_0^1 P \xrightarrow{\pi_P \circ_0^2 \pi_P} \hat{T}(\bullet) \circ_0^1 \hat{T}(\bullet)$ and lemma 2.19 provides Par_{η_P} and Par_{μ_P} .⁹

The category $\mathcal{O}_{\bullet}^{\hat{T}}$ of globular \hat{T} -operadic contractions over \bullet is just the full subcategory of $\mathcal{O}_{\bullet}^{\hat{T}, \kappa}$ determined by the objects defined above.

It is perfectly viable to use the category $\mathcal{O}_{\bullet}^{\hat{T}}$ instead of $\mathcal{O}_{\bullet}^{\hat{T}, \kappa}$ in order to define a slightly more restrictive notion of weak globular ω -category as an algebra for the initial object in $\mathcal{O}_{\bullet}^{\hat{T}}$, whose existence can be obtained following similar steps and in the case examined in the subsequent subsection 2.3.3. \square

2.3.3 Weak ω -categories

The subsequent fundamental result is proved in [Leinster 2004, appendix G].

Theorem 2.23. *The category $\mathcal{O}_{\bullet}^{\hat{T}, \kappa}$ of globular contracted \hat{T} -operads has initial objects.*

The proof provided in the reference above is not direct, and is based on the following definitions and results.

Definition 2.24. *A strict 1-category \mathcal{C} is **filtered** if any finite diagram $\mathcal{D} \xrightarrow{\exists} \mathcal{C}$ admits a co-cone.¹⁰*

⁹Notice that the contraction $\kappa_{\bullet} : (x^+, y, x^-) \mapsto y$ is uniquely determined by the defining axioms of contraction and the definition of π_{\bullet} .

¹⁰This is equivalent to requiring: a) that there exists at least an object in \mathcal{C}^0 (the vertex of a co-cone on the empty diagram); b) for any two objects $A_1, A_2 \in \mathcal{C}^0$, there exists a co-cone $A_1 \xrightarrow{\alpha_1} C \xleftarrow{\alpha_2} A_2$; c) for any finite diagram consisting of pair of parallel arrows $A_1 \xrightarrow{f, g} A_2$, there exists a co-cone $A_1 \xrightarrow{\alpha_1} C \xleftarrow{\alpha_2} A_1$ and hence $\alpha_2 \circ f = \alpha_1 = \alpha_2 \circ g$.

A **filtered co-limit** is a co-limit for a diagram $\mathcal{D} \xrightarrow{\mathfrak{D}} \mathcal{B}$ with filtered 1-category \mathcal{D} .

A **finitary functor** is a functor $\mathcal{A} \xrightarrow{\mathfrak{F}} \mathcal{B}$ that preserves filtered co-limits.

A **finitely presentable object** is an object $A \in \mathcal{C}^0$ whose associated covariant Hom-functor $\text{Hom}_{\mathcal{C}}(A, -)$ is finitary.

A category \mathcal{C} is **locally finite presentable** if the following properties are satisfied:

- ▶ \mathcal{C} is co-complete (it admits all co-limits of diagrams $\mathcal{D} \xrightarrow{\mathfrak{D}} \mathcal{C}$ with small \mathcal{D}),
- ▶ the full subcategory of finitely presentable objects of \mathcal{C} is essentially small,¹¹
- ▶ every object of \mathcal{C} is a filtered co-limit of at least one diagram of finitely presentable objects.

This first lemma is a consequence of [Leinster 2004, theorems 6.5.1, 6.5.2 and 6.5.4 in appendix D].

Lemma 2.25. *If T is a finitary Cartesian monad on a Cartesian category \mathcal{E} , the forgetful functor $\mathcal{O}_{\bullet}^T \xrightarrow{\mathfrak{U}} \mathcal{E}_{\bullet}^T$ has a left adjoint and the adjunction is monadic.*

In particular, from proposition 2.11, we have that $\mathcal{O}_{\bullet}^{\hat{T}} \xrightarrow{\mathfrak{U}} \mathcal{Q}_{\bullet}^{\hat{T}}$ is monadic.

Furthermore \mathfrak{U} is finitary [Leinster 2004, appendix G page 353].

This second lemma follows from a direct construction contained in [Leinster 2004, appendix G page 352].

Lemma 2.26. *The forgetful functor $\mathcal{Q}_{\bullet}^{\hat{T}, \kappa} \xrightarrow{\mathfrak{B}} \mathcal{Q}_{\bullet}^{\hat{T}}$ is finitary, has a left adjoint and the adjunction is monadic.*

This third lemma is dealt with in [Leinster 2004, appendix G page 352] using that $\mathcal{Q}_{\bullet}^{\hat{T}}$ is a pre-sheaf category.

Lemma 2.27. *The category $\mathcal{Q}_{\bullet}^{\hat{T}}$ is locally finitely presentable.*

Lemma 2.28. *[Kelly 1980, proposition 27.1] In the (big) category \mathcal{C} of functors between 1-categories, consider the co-span $\mathcal{A} \xrightarrow{\mathfrak{A}} \mathcal{X} \xleftarrow{\mathfrak{B}} \mathcal{B}$ and assume that:*

- ▶ \mathcal{X} is a locally finitely presentable 1-category,
- ▶ the functors \mathfrak{A} and \mathfrak{B} are both finitary and monadic,

¹¹Recall that in a **locally small category** \mathcal{C} , for all objects $A, B \in \mathcal{C}^0$, $\text{Hom}_{\mathcal{C}}(A, B)$ is a set; a locally small category is **small** if and only if \mathcal{C}^0 is a set; a category is **essentially small** if and only if it equivalent to a small category.

► the span $\mathcal{A} \xleftarrow{\mathfrak{A}} \mathcal{W} \xrightarrow{\mathfrak{B}} \mathcal{B}$ is a strict pull-back of the diagram $\mathcal{A} \xrightarrow{\mathfrak{A}} \mathcal{X} \xleftarrow{\mathfrak{B}} \mathcal{B}$

$$\begin{array}{ccc} \mathcal{W} & \xrightarrow{\mathfrak{A}} & \mathcal{B} \\ \mathfrak{B} \downarrow & & \downarrow \mathfrak{B} \\ \mathcal{A} & \xrightarrow{\mathfrak{A}} & \mathcal{X} \end{array}$$

The functor $\mathcal{W} \xrightarrow{\mathfrak{A} \circ \mathfrak{A} = \mathfrak{B} \circ \mathfrak{B}} \mathcal{X}$ is also monadic.

As a consequence [Leinster 2004, corollary G.1.2] the category \mathcal{W} has an initial object.

Theorem 2.23 follows applying the previous lemmata in the case of the strict pull-back of forgetful functors:

$$\begin{array}{ccc} \mathcal{O}_{\bullet}^{\hat{T}, \kappa} & \xrightarrow{\hat{u}_\kappa} & \mathcal{O}_{\bullet}^{\hat{T}} \\ \hat{u}_\sigma \downarrow & & \downarrow u_\sigma \\ \mathcal{Q}_{\bullet}^{\hat{T}, \kappa} & \xrightarrow{u_\kappa} & \mathcal{Q}_{\bullet}^{\hat{T}} \end{array}$$

Here is the basic notion of weak ω -category from [Leinster 2004, definition III.9.2.3].

Definition 2.29. A *weak ω -category* is an algebra for any monad L , that is an initial object in $\mathcal{O}_{\bullet}^{\hat{T}, \kappa}$.

2.3.4 Examples

The basic way to provide examples of weak globular ω -categories in Leinster's approach is described in [Leinster 2004, Example 9.2.4]:

any contracted \hat{T} -operad $(P, \kappa, \mu, \eta) \in \mathcal{O}_{\bullet}^{\hat{T}, \kappa}$ (this means for every choice of a contraction on a monad over \bullet in the bicategory $\mathcal{Q}_{\hat{T}}$) induces (since L is initial) a unique morphism $L \xrightarrow{\dagger} P$ in $\mathcal{O}_{\bullet}^{\hat{T}, \kappa}$ that produces a strict functor from algebras over P to algebras over L (that are, by definition, weak globular ω -categories). Hence an example of weak globular ω -category is obtained as soon as we are given:

- $(P, \pi, \kappa, \mu, \eta)$ a contracted \hat{T} -operad (that is just a contraction on a monad in $\mathcal{Q}_{\hat{T}}$),
- an algebra $X \in \mathcal{Q}_{\hat{T}}$ for the contracted \hat{T} -operad P (that is just an algebra for the previous monad).

Notable examples of weak globular ω -categories in Leinster's approach include:

- ▶ strict globular ω -categories (see example [Leinster 2004, 9.2.5]): these are just algebras for the terminal contracted \hat{T} -operad $\hat{T}(\bullet) \in \mathcal{O}^{\hat{T}, \kappa}$,¹²
- ▶ the globular ω -homotopy groupoid $\Pi_\omega(S)$ of a topological space S : in [Leinster 2004, example 9.2.7] it is shown that, for every topological space S , the ω -homotopy groupoid $\Pi_\omega(S)$ is an algebra for a certain contracted \hat{T} -operad.

3 Involutive Leinster Weak Globular ω -categories

In the present section we proceed to study how to introduce weak involutions on a weak globular ω -category.

We start in subsection 3.1 from involutions on strict globular ω -categories, as defined in our previous paper [Bejrakarbum Bertozzini 2017] that extended to the ω -category case the notion of fully involutive strict globular n -category introduced in [Bertozzini Conti Lewkeeratiyutkul Suthichitranont 2020, section 4].

Subsection 3.2 contains our tentative definition of involutive weak globular ω -category based on an initial generalized operad L^\star in the bicategory $\mathcal{Q}_{\hat{T}^\star}$ induced by the free involutive globular omega category monad \hat{T}^\star over the same Cartesian bicategory \mathcal{Q} of globular ω -sets.

3.1 Involutive Strict Globular ω -categories

We quickly recall the basic definitions and constructions of free strict involutive globular ω -categories, closely following our previous paper [Bejrakarbum Bertozzini 2017].

Definition 3.1. A (reflexive) ω -quiver $Q := Q^0 \begin{smallmatrix} \xleftarrow{s^0} \\ \xrightarrow{t^0} \end{smallmatrix} Q^1 \begin{smallmatrix} \xleftarrow{s^1} \\ \xrightarrow{t^1} \end{smallmatrix} \cdots \begin{smallmatrix} \xleftarrow{s^{n-2}} \\ \xrightarrow{t^{n-2}} \end{smallmatrix} Q^{n-1} \begin{smallmatrix} \xleftarrow{s^{n-1}} \\ \xrightarrow{t^{n-1}} \end{smallmatrix} Q^n \begin{smallmatrix} \xleftarrow{s^n} \\ \xrightarrow{t^n} \end{smallmatrix} \cdots$

is **fully self-dual** if it is equipped with a family $(\gamma_q)_{q \in \mathbb{N}}$ endomorphisms $Q \xrightarrow{\gamma_q} Q$, for all $q \in \mathbb{N}$, such that:

- ▶ for all $q \in \mathbb{N}$, γ_q is q -contravariant: $\gamma_q^q \circ s^q = t^q \circ \gamma_q^{q+1}$, $\gamma_q^q \circ t^q = s^q \circ \gamma_q^{q+1}$,
- ▶ for all $p, q \in \mathbb{N}$ such that $p \neq q$, γ_q is p -covariant: $\gamma_q^p \circ s^p = s^p \circ \gamma_q^{p+1}$, $\gamma_q^p \circ t^p = t^p \circ \gamma_q^{p+1}$.

A (reflexive) globular ω -set (respectively a (reflexive) globular ω -magma) is said to be **self-dual** whenever its underlying ω -quiver is self-dual in the previous sense.¹³

¹²The terminal object in $\mathcal{O}^{\hat{T}, \kappa}$ is $\bullet \xleftarrow{\hat{T}(\bullet)} \hat{T}(\bullet) \xrightarrow{\hat{T}(\bullet)} \hat{T}(\bullet)$, we will describe in more detail this point for our new monad \hat{T}^\star in remark 3.11.

¹³Here we interpret \circ, ι, γ as partially defined binary, nullary, unary operations, and we do not impose any algebraic axiom for magmas.

A strict globular ω -category $(\mathcal{C}, \circ, \iota)$ that is also a fully self-dual ω -quiver $(\mathcal{C}, *)$ is a **fully involutive strict globular ω -category** if its family $(*_q)_{q \in \mathbb{N}}$ of self-duality endomorphisms (here denoted by $x \mapsto x^{*_q}$, for $x \in \mathcal{C}^k$ and $k \in \mathbb{N}$) satisfies the following algebraic axioms:

- (**q -contravariance**):
 $(x \circ_q^k y)^{*_q} = y^{*_q} \circ_q^k x^{*_q}$, $\forall k, q \in \mathbb{N}$ such that $0 \leq q < k$ and $(x, y) \in \mathcal{C}^k \times_{\mathcal{C}^q} \mathcal{C}^k$,
 $(x \circ_q^k y)^{*_p} = x^{*_p} \circ_q^k y^{*_p}$, $\forall k, p, q \in \mathbb{N}$ such that $0 \leq q \neq p < k$ and $(x, y) \in \mathcal{C}^k \times_{\mathcal{C}^q} \mathcal{C}^k$,
- (**unitality**): $(\iota^k(x))^{*_q} = \iota^k(x^{*_q})$, for all $k, q \in \mathbb{N}$ and for all $x \in \mathcal{C}^k$,
- (**involutivity**): $(x^{*_q})^{*_q} = x$, for all $k, q \in \mathbb{N}$, and all $x \in \mathcal{C}^k$,
- (**commutativity**): $(x^{*_q})^{*_p} = (x^{*_p})^{*_q}$, for all $k, p, q \in \mathbb{N}$, and all $x \in \mathcal{C}^k$,
- (**q -grounding**): $x^{*_q} = x$, for all $k, q \in \mathbb{N}$ such that $k \leq q$ and all $x \in \mathcal{C}^k$.

A **self-dual morphism** between self-dual ω -quivers (and similarly between self-dual (reflexive) globular ω -sets or self-dual (reflexive) globular ω -magmas) is a morphism $(Q, \gamma) \xrightarrow{\phi} (\hat{Q}, \hat{\gamma})$ such that $\hat{\gamma} \circ \phi = \phi \circ \gamma$.

An **involutive ω -functor** $\mathcal{C} \xrightarrow{\phi} \hat{\mathcal{C}}$ between fully involutive strict globular ω -categories is just a self-dual morphism for the underlying self-dual globular ω -magmas.

This involutive version of proposition 2.3 was proved in [Bejrakarbum Bertozzini 2017, propositions 3.1, 3.2].

Proposition 3.2. Let \mathcal{C}^* denote the strict 1-category of covariant involutive ω -functors between fully involutive strict globular ω -categories.

For any globular ω -set Q in \mathcal{Q} , a **free strict involutive globular ω -category over Q** is a morphism of globular ω -sets $Q \xrightarrow{\eta_Q^*} \mathfrak{U}^*(\mathcal{C})$, into the underlying globular ω -set $\mathfrak{U}^*(\mathcal{C})$ of a strict involutive globular ω -category \mathcal{C} , satisfying the following universal factorization property: for any morphism of globular ω -sets $Q \xrightarrow{\phi} \mathfrak{U}^*(\hat{\mathcal{C}})$, into the underlying globular ω -set $\mathfrak{U}^*(\hat{\mathcal{C}})$ of a strict involutive globular ω -category $\hat{\mathcal{C}}$, there exists a unique involutive ω -functor $\mathcal{C} \xrightarrow{\hat{\phi}} \hat{\mathcal{C}}$ of fully involutive strict globular ω -categories such that $\phi = \hat{\phi} \circ \eta_Q^*$.

The **forgetful functor** $\mathcal{C}^* \xrightarrow{\mathfrak{U}^*} \mathcal{Q}$ (forgetting compositions, identities and involutions of objects in \mathcal{C}^*) admits a left-adjoint $\mathfrak{F}^* \dashv \mathfrak{U}^*$ **free strict involutive globular ω -category functor** $\mathcal{C}^* \xleftarrow{\mathfrak{F}^*} \mathcal{Q}$ that is uniquely determined via a specific construction of free involutive strict globular ω -category $Q \xrightarrow{\eta_Q^*} \mathfrak{U}^*(\mathcal{C})$ over a globular ω -set Q .

Proof. For convenience of the reader, we recall an explicit construction of the free strict globular involutive ω -category $Q \xrightarrow{\eta_Q^*} \mathfrak{F}^*(Q)$ over a given globular ω -set Q .

- a. We first produce the free globular self-dual reflexive ω -magma $Q \xrightarrow{\zeta_Q} \mathfrak{M}(Q)$, over the globular ω -set Q , with respect to all the partial binary compositions, the unary involutions and the nullary identities involved in the definition of strict globular involutive ω -category.
- b. Then we consider in $\mathfrak{M}(Q)$ the smallest congruence relation of globular self-dual reflexive ω -magmas $\Xi \subset \mathfrak{M}(Q) \times \mathfrak{M}(Q)$ containing all the pairs of terms appearing into the algebraic axioms that are involved in the definition of strict globular involutive ω -category.
- c. Finally, considering the quotient morphism $\mathfrak{M}(Q) \xrightarrow{\varpi} \mathfrak{M}(Q)/\Xi$ of globular self-dual reflexive ω -magmas and the map of globular ω -sets $Q \xrightarrow{\varpi \circ \zeta_Q} \mathfrak{M}(Q)/\Xi$, we notice that $\mathfrak{F}^*(Q) := \mathfrak{M}(Q)/\Xi$ is a strict involutive globular ω -category and that $\eta_Q^* := \varpi \circ \zeta_Q$ satisfies the universal factorization property for free involutive globular ω -categories over Q .

a.

The free globular self-dual reflexive ω -magma $Q \xrightarrow{\zeta_Q} \mathfrak{M}(Q)$ over the globular ω -set Q is obtained by a recursive definition. We first provide a 1-quiver $\mathfrak{M}(Q)^0 \Leftarrow \mathfrak{M}(Q)^1$.

Starting from Q^0 , we construct $\mathfrak{M}(Q)^0 := Q^0$.

We then introduce $Q^0 := \{(x, \iota^0) \mid x \in Q^0\}$ (that is a disjoint copy of Q^0 representing freely added identities of elements in Q^0) and $\mathfrak{M}(Q)^1[1]^0 := Q^1 \uplus Q^0$ with source/target given by $s_{\mathfrak{M}(Q)}^0(x) := s_Q^0(x)$, $t_{\mathfrak{M}(Q)}^0(x) := t_Q^0(x)$, for all $x \in Q^1$ and $s_{\mathfrak{M}(Q)}^0(x, \iota) := x =: t_{\mathfrak{M}(Q)}^0(x, \iota)$, for all $x \in Q^0$. If $\mathfrak{M}(Q)^1[1]^j$ has been already constructed, we define $\mathfrak{M}(Q)^1[1]^{j+1} := \{(x, \gamma_0) \mid x \in \mathfrak{M}(Q)^1[1]^j\}$ (that introduces freely added self-dualities of elements in $\mathfrak{M}(Q)^1[1]^j$) with source $s_{\mathfrak{M}(Q)}^0(x, \gamma_0) := t_{\mathfrak{M}(Q)}^0(x)$ and target $t_{\mathfrak{M}(Q)}^0(x, \gamma_0) := s_{\mathfrak{M}(Q)}^0(x)$; defining $\mathfrak{M}(Q)^1[1] := \bigcup_{j=1}^{+\infty} \mathfrak{M}(Q)^1[1]^j$, we get a 1-quiver $\mathfrak{M}(Q)^0 \Leftarrow \mathfrak{M}(Q)^1[1]$.

We proceed to define $\mathfrak{M}(Q)^1[2]^0 := \{(x, 0, y) \mid (x, y) \in \mathfrak{M}(Q)^1[1] \times_{\mathfrak{M}(Q)^0} \mathfrak{M}(Q)^1[1]\}$ (whose elements represent freely added compositions) with sources and targets given by $s_{\mathfrak{M}(Q)}^0(x, 0, y) := s_{\mathfrak{M}(Q)}^0(y)$ and $t_{\mathfrak{M}(Q)}^0(x, 0, y) := t_{\mathfrak{M}(Q)}^0(x)$. Exactly as done before, we define $\mathfrak{M}(Q)^1[2] := \bigcup_{j=1}^{+\infty} \mathfrak{M}(Q)^1[2]^j$, that freely introduces arbitrary iterations of γ_0 -self-dualities of the 1-cells in $\mathfrak{M}(Q)^1[2]^0$, obtaining a new 1-quiver as follows $\mathfrak{M}(Q)^0 \Leftarrow \mathfrak{M}(Q)^1[1] \cup \mathfrak{M}(Q)^1[2]$.

Supposing, by recursion, that the 1-quiver $\mathfrak{M}(Q)^0 \Leftarrow \bigcup_{n=1}^k \mathfrak{M}(Q)^1[n]$ is already defined, we extend it freely adding compositions

$$\mathfrak{M}(Q)^1[k+1]^0 := \{(x, 0, y) \mid (x, y) \in \mathfrak{M}(Q)^1[i] \times_{\mathfrak{M}(Q)^0} \mathfrak{M}(Q)^1[j], i+j=k+1\},$$

with sources/targets $s_{\mathfrak{M}(Q)}^0(x, 0, y) := s_{\mathfrak{M}(Q)}^0(y)$, $t_{\mathfrak{M}(Q)}^0(x, 0, y) := t_{\mathfrak{M}(Q)}^0(x)$, and freely adding arbitrary iterations of γ_0 -self-dualities to get $\mathfrak{M}(Q)^1[k+1] := \bigcup_{j=1}^{+\infty} \mathfrak{M}(Q)^1[k]^j$ and the 1-quiver $\mathfrak{M}(Q)^0 \Leftarrow \mathfrak{M}(Q)^1$, where we have $\mathfrak{M}(Q)^1 := \bigcup_{k=1}^{+\infty} \mathfrak{M}(Q)^1[k]$.

The previous 1-quiver, becomes reflexive defining $\iota^0 : \mathfrak{M}(Q)^0 \rightarrow \mathfrak{M}(Q)^1$ as follows $x \mapsto (x, \iota^0)$; it further becomes self-dual 1-quiver defining $\gamma_0 : \mathfrak{M}(Q)^1 \rightarrow \mathfrak{M}(Q)^1$ via $x \mapsto (x, \gamma_0)$; a binary composition $\circ_0^1 : \mathfrak{M}(Q)^1 \times_{\mathfrak{M}(Q)^0} \mathfrak{M}(Q)^1 \rightarrow \mathfrak{M}(Q)^1$ is also present defining $\circ_0^1 : (x, y) \mapsto (x, 0, y)$ and hence we have a reflexive self-dual (globular) 1-magma.

Supposing by recursion that a reflexive self-dual globular magma for the n -quiver $\mathfrak{M}(Q)^0 \Leftarrow \dots \Leftarrow \mathfrak{M}(Q)^n$ has been already defined, we will obtain a reflexive self-dual globular magma $\mathfrak{M}(Q)^0 \Leftarrow \dots \Leftarrow \mathfrak{M}(Q)^n \Leftarrow \mathfrak{M}(Q)^{n+1}$.

We start with $\mathfrak{M}(Q)^{n+1}[0]^0 := Q^{n+1} \cup \mathfrak{M}(Q)_i^n$, where $\mathfrak{M}(Q)_i^n := \{(x, \iota^n) \mid x \in \mathfrak{M}(Q)^n\}$ with sources and targets $s_{\mathfrak{M}(Q)}(x, \iota^n) := x =: t_{\mathfrak{M}(Q)}(x, \iota^n)$; we then introduce the sets $\mathfrak{M}(Q)^{n+1}[0]^{j+1} := \{(x, \gamma_q) \mid x \in \mathfrak{M}(Q)^{n+1}[0]^j, q = 0, \dots, n\}$, with sources/targets $s_{\mathfrak{M}(Q)}/t_{\mathfrak{M}(Q)}(x, \gamma_q) = s_{\mathfrak{M}(Q)}/t_{\mathfrak{M}(Q)}(x)$, if $q < n$, and, for $q = n$, $s_{\mathfrak{M}(Q)}(x, \gamma_n) = t_{\mathfrak{M}(Q)}(x)$, $t_{\mathfrak{M}(Q)}(x, \gamma_n) = s_{\mathfrak{M}(Q)}(x)$; and we get $\mathfrak{M}(Q)^{n+1}[0] := \bigcup_{j=1}^{+\infty} \mathfrak{M}(Q)^{n+1}[0]^j$. Next, assuming for recursion that $\mathfrak{M}(Q)^{n+1}[k]$ (with its source and target maps) have been already defined, we introduce free depth- p compositions

$$\mathfrak{M}(Q)^{n+1}[k+1]^0 := \left\{ (x, p, y) \mid p = 0, \dots, n, (x, y) \in \mathfrak{M}(Q)^{n+1}[i] \times_{\mathfrak{M}(Q)^p} \mathfrak{M}(Q)^{n+1}[j], \right. \\ \left. i + j = k + 1 \right\},$$

with source targets defined by $s_{\mathfrak{M}(Q)}(x, p, y) := (s_{\mathfrak{M}(Q)}(x), p, s_{\mathfrak{M}(Q)}(y))$ and respectively by $t_{\mathfrak{M}(Q)}(x, p, y) := (t_{\mathfrak{M}(Q)}(x), p, t_{\mathfrak{M}(Q)}(y))$, whenever $0 \leq p < n$, and otherwise $s_{\mathfrak{M}(Q)}(x, n, y) := s_{\mathfrak{M}(Q)}(y)$ and $t_{\mathfrak{M}(Q)}(x, n, y) := t_{\mathfrak{M}(Q)}(x)$.

Subsequently we introduce free iterated self-duals of the previous $(n+1)$ -cells by $\mathfrak{M}(Q)^{n+1}[k+1] := \bigcup_{j=0}^{+\infty} \mathfrak{M}(Q)^{n+1}[k+1]^j$ where, as above, we consider

$$\mathfrak{M}(Q)^{n+1}[k+1]^{j+1} := \left\{ (x, \gamma_q) \mid q = 0, \dots, n, x \in \mathfrak{M}(Q)^{n+1}[k+1]^j \right\}$$

with similarly defined source/target maps.

Finally we set $\mathfrak{M}(Q)^{n+1} := \bigcup_{k=0}^{+\infty} \mathfrak{M}(Q)^{n+1}[k]$ obtaining in this way the required globular $(n+1)$ -quiver that is reflexive, with the map $\iota^n(x) := (x, \iota^n)$, for $x \in \mathfrak{M}(Q)^n$; self-dual with the maps $x \mapsto (x, \gamma_q)$, for all $x \in \mathfrak{M}(Q)^{n+1}$ and all $q \in \{0, \dots, n\}$; and a magma for the partial compositions $x \circ_p^{n+1} y := (x, p, y)$, for all $(x, y) \in \mathfrak{M}(Q)^{n+1} \times_{\mathfrak{M}(Q)^p} \mathfrak{M}(Q)^{n+1}$ and all $p \in \{0, \dots, n\}$.

The recursive construction of the globular reflexive self-dual ω -magma

$$\mathfrak{M}(Q)^0 \Leftarrow \dots \Leftarrow \mathfrak{M}(Q)^n \Leftarrow \dots$$

is now completed and we further produce the morphism $Q \xrightarrow{\eta_Q} \mathfrak{M}(Q)$ of globular ω -sets by the inclusion of Q^n into $\mathfrak{M}(Q)^n[0]^0 \subset \mathfrak{M}(Q)^n$, for all $n \in \mathbb{N}$.

We only need to check the universal factorization property for $Q \xrightarrow{\eta_Q} \mathfrak{M}(Q)$. Suppose that $Q \xrightarrow{\phi} \hat{M}$ is a morphism of reflexive self-dual globular ω -sets into a reflexive self-dual globular ω -magma. Any grade-preserving map $\mathfrak{M}(Q) \xrightarrow{\hat{\phi}} \hat{M}$ such that $\phi = \hat{\phi} \circ \eta_Q$ must necessarily satisfy $x \mapsto \phi(x)$, for all $x \in Q^n \subset \mathfrak{M}(Q)^n$. Using the fact the $\hat{\phi}$ must be a morphism of reflexive self-dual globular ω -magmas, by induction, we obtain that, for all $n \in \mathbb{N}$, $\hat{\phi}(x, t^n) = \hat{\phi}_M^n(\phi(x))$, for all $x \in \mathfrak{M}(Q)^n$, $\hat{\phi}(x, p, y) = \phi(x) \hat{\phi}_p^n \phi(y)$, for all $(x, y) \in \mathfrak{M}(Q)^n \times_{\mathfrak{M}(Q)^p} \mathfrak{M}(Q)^n$ and $0 \leq p < n$, and $\hat{\phi}(x, \gamma_q) = \phi(x)^{*q}$, for all $x \in \mathfrak{M}(Q)^n$ and $0 \leq q < n$. This uniquely defined map $\mathfrak{M}(Q) \xrightarrow{\hat{\phi}} \hat{M}$ is a morphism of reflexive self-dual globular ω -magmas as required.

b.

From [Bejrakarbum Bertozzini 2017, section 3.2] we recall that, given globular ω -sets (Q, s_Q, t_Q) , $(\hat{Q}, s_{\hat{Q}}, t_{\hat{Q}})$ the *Cartesian product of globular ω -sets* is the globular ω -set $(Q \times \hat{Q}, s_{Q \times \hat{Q}}, t_{Q \times \hat{Q}})$ defined, for all $n \in \mathbb{N}$, as $(Q \times \hat{Q})^n := Q^n \times \hat{Q}^n$, with source/targets $s_{Q \times \hat{Q}}^n := (s_Q^n, s_{\hat{Q}}^n)$ and $t_{Q \times \hat{Q}}^n := (t_Q^n, t_{\hat{Q}}^n)$ acting componentwise; and that whenever $(Q, s_Q, t_Q, \iota_Q, \gamma_Q, \circ_Q)$, $(\hat{Q}, s_{\hat{Q}}, t_{\hat{Q}}, \iota_{\hat{Q}}, \gamma_{\hat{Q}}, \circ_{\hat{Q}})$ are globular self-dual reflexive ω -magmas, also their Cartesian product $Q \times \hat{Q}$ is a globular self-dual reflexive ω -magma with the componentwise defined nullary $t_{Q \times \hat{Q}}^n := (t_Q^n, t_{\hat{Q}}^n)$, unary $\gamma_q^{(Q \times \hat{Q})^n} := (\gamma_q^Q, \gamma_q^{\hat{Q}})$ and binary $\circ_p^{(Q \times \hat{Q})^n} := (\circ_p^Q, \circ_p^{\hat{Q}})$ operations.

We also recall that a *congruence* X in a globular self-dual reflexive ω -magma M is a globular self-dual reflexive ω -magma X such that $X^n \subset M^n \times M^n$, for all $n \in \mathbb{N}$, in such a way that the inclusion $X \xrightarrow{\phi} M \times M$ is a morphism of globular self-dual reflexive ω -magmas.

Inside the free globular self-dual reflexive ω -magma $\mathfrak{M}(Q)$ over Q constructed in a. above we consider the congruence $\Xi \subset \mathfrak{M}(Q) \times \mathfrak{M}(Q)$ generated¹⁴ by the union of all of the following families of pairs:

$$\begin{aligned} & \left\{ (x \circ_p^n (y \circ_p^n z), (x \circ_p^n y) \circ_p^n z) \mid n > p \in \mathbb{N}, \right. \\ & \qquad \qquad \qquad \left. (x, y, z) \in \mathfrak{M}(Q)^n \times_{\mathfrak{M}(Q)^p} \mathfrak{M}(Q)^n \times_{\mathfrak{M}(Q)^p} \mathfrak{M}(Q)^n \right\} \\ & \left\{ ((t^{n-1} \circ \dots \circ t^p \circ t^p \circ \dots \circ t^{n-1}(x)) \circ_p^n x, x) \mid n > p \in \mathbb{N}, x \in \mathfrak{M}(Q)^n \right\} \\ & \left\{ (x, x \circ_p^n (t^{n-1} \circ \dots \circ t^p \circ s^p \circ \dots \circ s^{n-1}(x))) \mid n > p \in \mathbb{N}, x \in \mathfrak{M}(Q)^n \right\} \end{aligned}$$

¹⁴Since intersection of congruences is a congruence, Ξ is just the intersection of all the congruences containing the given pairs.

$$\begin{aligned}
& \{(\iota^n(x) \circ_p^{n+1} \iota^n(y), \iota^n(x \circ_p^n y)) \mid n > p \in \mathbb{N}, (x, y) \in \mathfrak{M}(Q)^n \times_{\mathfrak{M}(Q)^p} \mathfrak{M}(Q)^n\} \\
& \{((x \circ_p^n y) \circ_q^n (z \circ_p^n w), (x \circ_q^n z) \circ_p^n (y \circ_q^n w)) \mid n > p, q \in \mathbb{N}, \\
& \quad (x, y), (z, w) \in \mathfrak{M}(Q)^n \times_{\mathfrak{M}(Q)^p} \mathfrak{M}(Q)^n, \\
& \quad (x, z), (y, w) \in \mathfrak{M}(Q)^n \times_{\mathfrak{M}(Q)^q} \mathfrak{M}(Q)^n\} \\
& \{(\gamma_q^n(\gamma_q^n(x)), x) \mid n > q \in \mathbb{N}, x \in \mathfrak{M}(Q)^n\} \\
& \{(\gamma_q^n(\gamma_p^n(x)), \gamma_p^n(\gamma_q^n(x))) \mid n > q, p \in \mathbb{N}, x \in \mathfrak{M}(Q)^n\} \\
& \{(\gamma_q^n(x \circ_p^n y), \gamma_q^n(x) \circ_p^n \gamma_q^n(y)) \mid n > p \neq q \in \mathbb{N}, \\
& \quad (x, y) \in \mathfrak{M}(Q)^n \times_{\mathfrak{M}(Q)^p} \mathfrak{M}(Q)^n\} \\
& \{(\gamma_q^n(x \circ_p^n y), \gamma_q^n(y) \circ_p^n \gamma_q^n(x)) \mid n > p = q \in \mathbb{N}, \\
& \quad (x, y) \in \mathfrak{M}(Q)^n \times_{\mathfrak{M}(Q)^p} \mathfrak{M}(Q)^n\} \\
& \{(\iota^n(\gamma_q^n(x)), \gamma_q^{n+1}(\iota_q^n(x))) \mid n > q \in \mathbb{N}, x \in \mathfrak{M}^n(Q)\} \\
& \{(\gamma_q^n(x), x) \mid n \leq q \in \mathbb{N}, x \in \mathfrak{M}(Q)^n\}.
\end{aligned}$$

c.

As quotient of a globular self-dual reflexive ω -magma by a globular ω -congruence, the quotient $\mathfrak{M}(Q)/\Xi$ is a globular ω -set with

$$(\mathfrak{M}(Q)/\Xi)^n := \mathfrak{M}(Q)^n / \Xi^n = \{[x]_{\Xi^n} \mid x \in \mathfrak{M}(Q)^n\}$$

and $s_{\mathfrak{M}(Q)/\Xi}^n([x]_{\Xi^{n+1}}) := [s_{\mathfrak{M}(Q)}^n(x)]_{\Xi^n}$, $t_{\mathfrak{M}(Q)/\Xi}^n([x]_{\Xi^{n+1}}) := [t_{\mathfrak{M}(Q)}^n(x)]_{\Xi^n}$, for $n \in \mathbb{N}$ and $x \in \mathfrak{M}(Q)^{n+1}$; and $\mathfrak{M}(Q)/\Xi$ is actually a globular self-dual reflexive ω -magma with well-defined compositions given by $[x]_{\Xi^n} \circ_p^{(\mathfrak{M}(Q)/\Xi)^n} [y]_{\Xi^n} := [x \circ_p^{\mathfrak{M}(Q)^n} y]_{\Xi^n}$, self-dualities $\gamma_q([x]_{\Xi^n}) := [\gamma_q(x)]_{\Xi^n}$, and reflexive maps $u_{\mathfrak{M}(Q)/\Xi}^n([x]_{\Xi^n}) := [u_{\mathfrak{M}(Q)}^n(x)]_{\Xi^{n+1}}$, whenever $n > p \in \mathbb{N}$, $q \in \mathbb{N}$, $x, y \in \mathfrak{M}(Q)^n$. Furthermore, since all the algebraic axioms for strict globular involutive ω -category are already included in Ξ , we have that the quotient $\mathfrak{M}(Q)/\Xi$ is actually a strict globular involutive ω -category and the quotient morphism $\varpi : \mathfrak{M}(Q) \rightarrow \mathfrak{M}(Q)/\Xi$, defined as $\varpi^n(x) := [x]_{\Xi^n}$, for all $n \in \mathbb{N}$ and $x \in \mathfrak{M}(Q)^n$, is a morphism of globular self-dual reflexive ω -sets.

We only need to check that the morphism of globular ω -sets $Q \xrightarrow{\eta_Q^*} \mathfrak{M}(Q)/\Xi$, with $\eta_Q^* := \varpi \circ \zeta_Q$, into the globular involutive ω -category $\mathfrak{M}(Q)/\Xi$, satisfies the universal factorization property. For any other morphism $Q \xrightarrow{\phi} \mathcal{C}$ of globular self-dual reflexive ω -sets, into a strict globular involutive ω -category \mathcal{C} , since $Q \xrightarrow{\zeta_Q} \mathfrak{M}(Q)$ is a free globular self-dual reflexive ω -magma, there exists a unique morphism $\mathfrak{M}(Q) \xrightarrow{\bar{\phi}} \mathcal{C}$ of globular self-dual reflexive ω -magmas such that $\phi = \bar{\phi} \circ \zeta_Q$. Consider now the *kernel*

globular ω -congruence $\Xi_{\bar{\phi}}$ induced by the morphism $\bar{\phi}$ of ω -magmas: for all $n \in \mathbb{N}$, we have $\Xi_{\bar{\phi}}^n := \{(x, y) \in \mathfrak{M}(Q)^n \times \mathfrak{M}(Q)^n \mid \bar{\phi}(x) = \bar{\phi}(y)\}$ and, since all the axioms of strict globular involutive ω -category are already satisfied in \mathcal{C} , we have $\Xi \subset \Xi_{\bar{\phi}}$ and hence $\mathfrak{M}(Q)/\Xi_{\bar{\phi}}$ is already a strict globular involutive ω -category, furthermore the assignment $\tilde{\phi} : [x]_{\Xi_{\bar{\phi}}} \mapsto \bar{\phi}(x)$ is a well-defined covariant involutive ω -functor $\mathfrak{M}(Q)/\Xi_{\bar{\phi}} \xrightarrow{\tilde{\phi}} \mathcal{C}$, that is actually the unique morphism such that $\bar{\phi} = \tilde{\phi} \circ \varpi_{\bar{\phi}}$, where $\mathfrak{M}(Q) \xrightarrow{\varpi_{\bar{\phi}}} \mathfrak{M}(Q)/\Xi_{\bar{\phi}}$ is the quotient morphism of globular self-dual reflexive ω -magmas defined as usual by $\varpi_{\bar{\phi}} : x \mapsto [x]_{\Xi_{\bar{\phi}}}$, for $x \in \mathfrak{M}(Q)$. From $\Xi \subset \Xi_{\bar{\phi}}$, we obtain a unique involutive ω -functor $\mathfrak{M}(Q)/\Xi_{\bar{\phi}} \xrightarrow{\theta} \mathfrak{M}(Q)/\Xi$, $\theta : [x]_{\Xi_{\bar{\phi}}} \mapsto [x]_{\Xi}$, for $x \in \mathfrak{M}(Q)$, such that $\varpi_{\bar{\phi}} \circ \varpi = \theta$. Defining $\hat{\phi} := \tilde{\phi} \circ \theta$, we have that $\mathfrak{M}(Q)/\Xi \xrightarrow{\hat{\phi}} \mathcal{C}$ is the unique involutive ω -functor such that $\hat{\phi} \circ \eta_Q^* = \tilde{\phi} \circ \theta \circ \varpi \circ \zeta_Q = \tilde{\phi} \circ \varpi_{\bar{\phi}} \circ \eta_Q = \bar{\phi} \circ \eta_Q = \phi$. \square

Remark 3.3. The existence of algebras (with a given signature) that are free over a set is known; ¹⁵ propositions 2.3 and 3.2 are essentially special cases of a general existence theorem for “ ω -algebras” that are free over a globular ω -set that (for the ω -globular setting) is vertically categorifying the case of algebras over sets. \lrcorner

In parallel with corollary 2.9, from proposition 3.2 we obtain a new monad in the involutive ω -category case.

Corollary 3.4. *On the 1-category \mathcal{Q} of morphisms of globular ω -sets, we have the following*

- *free involutive strict globular ω -category monad $\hat{T}^* := \mathfrak{U}^* \circ \mathfrak{F}^*$.*

3.2 Involutive Weak Globular ω -categories

The following is the “involutive case” version of proposition 2.11.

Proposition 3.5. *The 1-category \mathcal{Q} of small globular ω -sets with morphisms of globular ω -sets is Cartesian. The 1-category \mathcal{C}^* of small strict globular involutive ω -categories with involutive ω -functors is Cartesian. The forgetful 1-functor $\mathcal{C}^* \xrightarrow{\mathfrak{U}^*} \mathcal{Q}$ and the free strict globular involutive ω -category 1-functor $\mathcal{C}^* \xleftarrow{\mathfrak{F}^*} \mathcal{Q}$ are Cartesian. The free strict globular involutive ω -category monad $T^* := \mathfrak{U}^* \circ \mathfrak{F}^*$ is Cartesian.*

Proof. The Cartesianity of the 1-category \mathcal{Q} of globular ω -sets is already known by proposition 2.11. Following the exposition in [Bejrakarbun 2023, section 3.2], we recall that, given a span $Q \xrightarrow{\phi} X \xleftarrow{\psi} R$ of globular ω -sets in \mathcal{Q} , a pull-back can be constructed via the co-span of globular ω -sets $Q \xleftarrow{\hat{\psi}} Q \times_X R \xrightarrow{\hat{\phi}} R$ defined by

$$Q \times_X R := \left((Q \times_X R)^n \begin{array}{c} \xleftarrow{s^n} \\ \xrightarrow{r^n} \end{array} (Q \times_X R)^{n+1} \right)_{n \in \mathbb{N}}$$

¹⁵See section 6 in the n-Lab entry: <https://ncatlab.org/nlab/show/variety+of+algebras#literature>.

where we have $(Q \times_X R)^n := Q^n \times_{X^n} R^n := \{(q, r) \in Q^n \times R^n \mid \phi^n(q) = \psi^n(r)\}$, with $s^n(q, r) := (s_Q^n(q), s_R^n(r))$ and $t^n(q, r) := (t_Q^n(q), t_R^n(r))$, with $\hat{\phi}^n(q, r) := r$ and $\hat{\psi}^n(q, r) := q$, for $n \in \mathbb{N}$.

To prove the Cartesianity of the 1-category \mathcal{C}^* , from [Bejrakarbum 2023, section 3.3], we recall that given any co-span $\mathcal{A} \xrightarrow{\phi} \mathcal{X} \xleftarrow{\psi} \mathcal{B}$ in \mathcal{C}^* , a pull-back can be constructed via the previous span $\mathcal{A} \xleftarrow{\hat{\psi}} \mathcal{A} \times_{\mathcal{X}} \mathcal{B} \xrightarrow{\hat{\phi}} \mathcal{B}$ in \mathcal{Q} , noting that the globular ω -set $\mathcal{A} \times_{\mathcal{X}} \mathcal{B}$ becomes a strict involutive globular ω -category, with componentwise compositions $(a_1, b_1) \circ_q^n (a_2, b_2) := (a_1 \circ_{\mathcal{A}_q^n} a_2, b_1 \circ_{\mathcal{B}_q^n} b_2)$, identities $l^n(a, b) := (l_{\mathcal{A}}^n(a), l_{\mathcal{B}}^n(b))$, involutions $(a, b)^{*q} := (a^{*q}, b^{*q})$; and that the above-defined $\hat{\phi}$ and $\hat{\psi}$ turn out to be involutive covariant ω -functors.

From the previous explicit definitions of pull-backs in \mathcal{Q} and \mathcal{C}^* , it follows that the forgetful functor \mathcal{U}^* is Cartesian, since it associates to the standard pull-back of strict involutive globular ω -categories the standard pull-back of their underlying globular ω -sets.

In order to prove the Cartesianity of the free strict involutive globular ω -category functor \mathfrak{F}^* , we simply notice that $\mathfrak{F}^*(Q) \xleftarrow{\widehat{\mathfrak{F}^*(\psi)}} \mathfrak{F}^*(Q \times_X R) \xrightarrow{\widehat{\mathfrak{F}^*(\phi)}} \mathfrak{F}^*(R)$ is canonically \mathcal{C}^* -isomorphic to the standard \mathcal{C}^* -pull-back

$$\mathfrak{F}^*(Q) \xleftarrow{\widehat{\mathfrak{F}^*(\psi)}} \mathfrak{F}^*(Q) \times_{\mathfrak{F}^*(X)} \mathfrak{F}^*(R) \xrightarrow{\widehat{\mathfrak{F}^*(\phi)}} \mathfrak{F}^*(R)$$

of the co-span $\mathfrak{F}^*(Q) \xrightarrow{\widehat{\mathfrak{F}^*(\phi)}} \mathfrak{F}^*(X) \xleftarrow{\widehat{\mathfrak{F}^*(\psi)}} \mathfrak{F}^*(R)$, via an involutive ω -functor.

The composition of Cartesian functors is Cartesian, hence the Cartesianity of the monad $T^* := \mathcal{U}^* \circ \mathfrak{F}^*$.

For the Cartesianity of the natural transformation η^* , we must show that, for any morphism $Q_1 \xrightarrow{\phi} Q_2$ of globular ω -sets, the solid square commuting diagram below is a pull-back in \mathcal{Q} ; for this purpose, for any span $Q_2 \xleftarrow{\beta} R \xrightarrow{\alpha} \hat{T}^*(Q_1)$ such that $\hat{T}^*(\phi) \circ \alpha = \eta_{Q_2}^* \circ \beta$, we must see that there exists a unique morphism $R \xrightarrow{\theta} Q_1$ making commutative the two triangle diagrams below: $\alpha = \eta_{Q_1}^* \circ \theta$, $\beta = \phi \circ \theta$.

$$\begin{array}{ccccc}
 R & & & & \\
 \text{\scriptsize θ} \swarrow & & \text{\scriptsize α} \searrow & & \\
 & Q_1 & \xrightarrow{\eta_{Q_1}^*} & \hat{T}^*(Q_1) & \\
 \text{\scriptsize β} \searrow & \downarrow \phi & & \downarrow \hat{T}^*(\phi) & \\
 & Q_2 & \xrightarrow{\eta_{Q_2}^*} & \hat{T}^*(Q_2) &
 \end{array}$$

From the explicit construction of the free involutive globular ω -category of a globular ω -set recalled in proposition 3.2 we have that $\eta_Q^*(x) = \varpi \circ \zeta_Q(x) = [x]_{\Xi}$, for all $x \in Q$, is a singleton containing only x . Hence, for all $r \in R$, $\eta_{Q_2}^*(\beta(r)) = [x_2]_{\Xi_2}$ is a singleton in $\hat{T}^*(Q_2)$, with $x_2 \in Q_2$; since $\hat{T}^*(\phi)$, using the fact that every morphism of globular ω -sets, is “degree-preserving”, there exists a unique element $\theta(r) \in Q_1^0$ such that $(\hat{T}^*(\phi))([\theta(r)]_{\Xi_1}) = [x_2]_{\Xi_2}$. Such θ satisfies our requirements.

For the Cartesianity of the natural transformation μ_Q , we must show that, for any morphism $Q_1 \xrightarrow{\phi} Q_2$ of globular ω -sets, the solid square commuting diagram below is a pull-back in \mathcal{Q} ; for this purpose, for any span $(\hat{T}^* \circ \hat{T}^*)(Q_2) \xleftarrow{\beta} R \xrightarrow{\alpha} (\hat{T}^* \circ \hat{T}^*)(Q_1)$ such that $\phi \circ \alpha = \mu_{Q_2}^* \circ \beta$, we must see that there exists a unique morphism $R \xrightarrow{\theta} (\hat{T}^* \circ \hat{T}^*)(Q_1)$ making commutative the two triangle diagrams below: $\beta = ((\hat{T}^* \circ \hat{T}^*)(\phi)) \circ \theta$ and $\alpha = \mu_{Q_1}^* \circ \theta$.

$$\begin{array}{ccccc}
 R & & & & \\
 \text{\scriptsize θ} \searrow & & \text{\scriptsize α} \searrow & & \\
 & (\hat{T}^* \circ \hat{T}^*)(Q_1) & \xrightarrow{\mu_{Q_1}^*} & \hat{T}^*(Q_1) & \\
 & \downarrow (\hat{T}^* \circ \hat{T}^*)(\phi) & & \downarrow \hat{T}^*(\phi) & \\
 \text{\scriptsize β} \searrow & (\hat{T}^* \circ \hat{T}^*)(Q_2) & \xrightarrow{\mu_{Q_2}^*} & \hat{T}^*(Q_2) & \\
 & & & &
 \end{array}$$

Since for all $r \in R$, we have $\alpha(r) \in \hat{T}^*(Q_1)$, the only possible element in $(\hat{T}^* \circ \hat{T}^*)(Q_1)$ that maps, via $\mu_{Q_1}^*$ to $\alpha(r)$, must necessarily be $(\alpha(r))$ and the assignment $r \mapsto (\alpha(r))$ is a morphism of globular ω -sets satisfying the required conditions. \square

As direct application of proposition 2.12 to the Cartesian monad \hat{T}^* on the Cartesian category \mathcal{Q} we obtain:

Corollary 3.6. *There is a bicategory $\mathcal{Q}_{\hat{T}^*}$.*

The notion of Leinster contraction in definition 2.17 remains unchanged and, as anticipated in remark 2.21, we have a parallel version of definition 2.18 and proposition 2.20 that reformulate as follows, for the case of the free involutive globular ω -category monad \hat{T}^* .

Definition 3.7. *Let $\bullet \in \mathcal{Q}$ denote a terminal object in the category of globular ω -sets.*

*A **globular \hat{T}^* -collection** is a morphism $Q \xrightarrow{\pi} \hat{T}^*(\bullet)$ in \mathcal{Q} ; a **globular contracted \hat{T}^* -collection** consists of a Leinster contraction κ on a globular \hat{T}^* -collection π : $\text{Par}(\pi) \xrightarrow{\kappa} Q \xrightarrow{\pi} \hat{T}^*(\bullet)$.*

Similarly, using the bicategory $\mathcal{Q}_{\hat{T}^*}$, for the Cartesian monad \hat{T}^* , definition 2.16 already provides the notion of globular (contracted) \hat{T}^* -operad over \bullet . For technical

reasons, in the proof of the subsequent theorem 3.16, we actually need to introduce the following more general “magma structure” internal to $\mathcal{Q}_{\hat{T}^*}$.

Definition 3.8. A **globular \hat{T}^* -operadic magma**¹⁶ $(M, \pi_M, \eta_M, \mu_M)$ over \bullet is a 1-cell $\bullet \xrightarrow{M} \bullet$ in the bicategory $\mathcal{Q}_{\hat{T}^*}$ (hence a globular \hat{T}^* -collection $M \xrightarrow{\pi_M} \hat{T}^*(\bullet)$) that is equipped with a unit 2-cell $\bullet \xrightarrow{\eta_M} M$ and a multiplication 2-cell $M \circ_0^1 M \xrightarrow{\mu_M} M$ as specified in the following commutative diagrams¹⁷ in the category \mathcal{Q}

$$\begin{array}{ccc}
 \begin{array}{c} \bullet \\ \downarrow \eta_M \\ \bullet \end{array} & = & \hat{T}^*(\bullet) \\
 \begin{array}{c} \bullet \\ \downarrow \eta_M \\ \bullet \end{array} & \xrightarrow{\eta_M} & M \\
 \downarrow \eta_M & & \downarrow \pi_M \\
 \bullet & \xrightarrow{\eta_{\hat{T}^*(\bullet)}} & \hat{T}^*(\bullet)
 \end{array} \quad (3.1)$$

$$\begin{array}{ccc}
 \begin{array}{c} M \circ_0^1 M \\ \downarrow \mu_M \\ M \end{array} & = & \hat{T}^*(\bullet) \\
 \begin{array}{c} M \circ_0^1 M \\ \downarrow \mu_M \\ M \end{array} & \xrightarrow{\mu_M} & M \\
 \downarrow \mu_M & & \downarrow \pi_M \\
 \hat{T}^*(\bullet) \circ_0^1 \hat{T}^*(\bullet) & \xrightarrow{\mu_{\hat{T}^*(\bullet)}} & \hat{T}^*(\bullet)
 \end{array} \quad (3.2)$$

that are not necessarily required to satisfy the operadic axioms 2.3.

A **globular \hat{T}^* -operad** is a globular \hat{T}^* -operadic magma with unit and multiplication that satisfy the monadic associativity and unitality axioms 2.3.

Proposition 3.9. For any terminal object $\bullet \in \mathcal{Q}^0$, we have the following categories:

- ▶ the category $\mathcal{Q}_{\hat{T}^*}$ of globular \hat{T}^* -collections over \bullet ,
- ▶ the category $\mathcal{Q}_{\hat{T}^*,k}$ of globular contracted \hat{T}^* -collections over \bullet ,

¹⁶Of course, although the definition is here given in the special case of the bicategory $\mathcal{Q}_{\hat{T}^*}$, it remains perfectly valid when applied to an arbitrary bicategory \mathcal{E}_T , for a given Cartesian monad T ; furthermore one can define *multicategorical magmas over E* using a given object E in place of a terminal \bullet in \mathcal{E}_T .

¹⁷For the description of the notation required in the square diagrams on the right, refer to remark 3.11 below.

- the category $\mathcal{M}_{\bullet}^{\hat{T}^*}$ of globular \hat{T}^* -operadic magmas over \bullet ,
- the category $\mathcal{O}_{\bullet}^{\hat{T}^*}$ of globular \hat{T}^* -operads over \bullet ,
- the category $\mathcal{M}_{\bullet}^{\hat{T}^*, \kappa}$ of globular contracted \hat{T}^* -operadic magmas over \bullet ,
- the category $\mathcal{O}_{\bullet}^{\hat{T}^*, \kappa}$ of globular contracted \hat{T}^* -operads over \bullet .

There are commuting diagram (in the category of functors between 1-categories) of forgetful functors:

$$\begin{array}{ccc}
 \mathcal{O}_{\bullet}^{\hat{T}^*, \kappa} & \xrightarrow{\hat{u}_{\kappa}^*} & \mathcal{O}_{\bullet}^{\hat{T}^*} \\
 \hat{u}_{\mathcal{O}}^* \downarrow & & \downarrow u_{\mathcal{O}}^* \\
 \mathcal{Q}_{\bullet}^{\hat{T}^*, \kappa} & \xrightarrow{u_{\kappa}^*} & \mathcal{Q}_{\bullet}^{\hat{T}^*}
 \end{array}
 \quad
 \begin{array}{ccc}
 \mathcal{M}_{\bullet}^{\hat{T}^*, \kappa} & \xrightarrow{\hat{u}_{\kappa}^*} & \mathcal{M}_{\bullet}^{\hat{T}^*} \\
 \hat{u}_{\mathcal{M}}^* \downarrow & & \downarrow u_{\mathcal{M}}^* \\
 \mathcal{Q}_{\bullet}^{\hat{T}^*, \kappa} & \xrightarrow{u_{\kappa}^*} & \mathcal{Q}_{\bullet}^{\hat{T}^*}
 \end{array}$$

The categories $\mathcal{O}_{\bullet}^{\hat{T}^*}$, respectively $\mathcal{O}_{\bullet}^{\hat{T}^*, \kappa}$ are full subcategories of $\mathcal{M}_{\bullet}^{\hat{T}^*}$, respectively of $\mathcal{M}_{\bullet}^{\hat{T}^*, \kappa}$.

Remark 3.10. Exactly as already noticed in remark 2.22, it is possible to introduce more restrictive notions of globular \hat{T}^* -operadic magma contraction and globular \hat{T}^* -operadic contraction.

A **globular \hat{T}^* -operadic magma contraction** over \bullet $(M, \pi_M, \kappa_M, \eta_M, \mu_M)$ consists of a globular contracted \hat{T}^* -operadic magma over \bullet that further satisfies the commutativity of the following two diagrams in $\mathcal{Q}_{\bullet}^{\hat{T}^*}$:

$$\begin{array}{ccc}
 \text{Par}(\pi_{\bullet}) & \xrightarrow{\text{Par}_{\eta_M}} & \text{Par}(\pi_M) \\
 \kappa_{\bullet} \downarrow & & \downarrow \kappa_M \\
 \bullet & \xrightarrow{\eta_M} & M
 \end{array}
 \quad
 \begin{array}{ccc}
 \text{Par}(\pi_M \circ_0^2 \pi_M) & \xrightarrow{\text{Par}_{\mu_M}} & \text{Par}(\pi_M) \\
 \kappa_M \circ_0^2 \kappa_M \downarrow & & \downarrow \kappa_M \\
 M \circ_0^1 M & \xrightarrow{\mu_M} & M.
 \end{array}
 \quad (3.3)$$

A **globular \hat{T}^* -operadic contraction** over \bullet is just a globular contracted \hat{T}^* -operad $(M, \pi_M, \kappa_M, \eta_M, \mu_M)$ that also satisfies the previous commutative diagrams.

We can introduce the category $\mathcal{MH}_{\bullet}^{\hat{T}^*}$, of globular \hat{T}^* -operadic magma contractions over \bullet , as the full subcategory of $\mathcal{M}_{\bullet}^{\hat{T}^*, \kappa}$; similarly the category $\mathcal{OH}_{\bullet}^{\hat{T}^*}$, of globular \hat{T}^* -operadic contractions over \bullet , as the full subcategory of $\mathcal{O}_{\bullet}^{\hat{T}^*, \kappa}$. \square

The following remark is absolutely crucial for us: it identifies the terminal contracted- \hat{T}^* -operad in $\mathcal{O}_{\bullet}^{\hat{T}^*, \kappa}$.

Remark 3.11. Notice that the globular ω -set $\hat{T}^*(\bullet)$ is naturally a globular \hat{T}^* -collection $\hat{T}^*(\bullet) \xrightarrow{\pi_{\hat{T}^*(\bullet)}} \hat{T}^*(\bullet)$ with projection $\pi_{\hat{T}^*(\bullet)}$ the identity morphism of globular ω -sets,

- for any \hat{T}^* -collection $Q \xrightarrow{\pi} \hat{T}^*(\bullet)$, the projection π is a morphism in $\mathcal{Q}_{\bullet}^{\hat{T}^*}$,

$$\begin{array}{ccc} Q & \xrightarrow{\pi} & \hat{T}^*(\bullet) \\ & \searrow \pi & \swarrow \pi_{\hat{T}^*(\bullet)} \\ & \hat{T}^*(\bullet) & \end{array} \quad \pi_{\hat{T}^*(\bullet)} : x \mapsto x.$$

It also naturally becomes a contracted globular \hat{T}^* -collection

$$\text{Par}(\pi_{\hat{T}^*(\bullet)}) \xrightarrow{\kappa_{\hat{T}^*(\bullet)}} \hat{T}^*(\bullet) \xrightarrow{\pi_{\hat{T}^*(\bullet)}} \hat{T}^*(\bullet)$$

with contraction $\kappa_{\hat{T}^*(\bullet)} : (y^+, y, y^-) \mapsto y$ on

$$\text{Par}(\pi_{\hat{T}^*(\bullet)}) = \left\{ (y^+, y, y^-) \in \hat{T}^*(\bullet) \times \hat{T}^*(\bullet) \times \hat{T}^*(\bullet) \mid \begin{array}{c} y^+ \\ \Downarrow y \\ y^- \end{array} \right\},$$

- for any contracted \hat{T}^* -collection $\text{Par}(\pi) \xrightarrow{\kappa} Q \xrightarrow{\pi} \hat{T}^*(\bullet)$, the projection π is a morphism in $\mathcal{Q}_{\bullet}^{\hat{T}^*, \kappa}$:

$$\begin{array}{ccc} \text{Par}(\pi) & \xrightarrow{\kappa} & Q \\ \text{Par}_{\pi} \downarrow & & \downarrow \pi \\ \text{Par}(\pi_{\hat{T}^*(\bullet)}) & \xrightarrow{\kappa_{\hat{T}^*(\bullet)}} & \hat{T}^*(\bullet) \end{array}$$

$\kappa_{\hat{T}^*(\bullet)} : (y^+, y, y^-) \mapsto y, \quad \text{Par}_{\pi} : (x^+, y, x^-) \mapsto (\pi(x^+), y, \pi(x^-)).$

Furthermore $\hat{T}^*(\bullet)$ is a globular \hat{T}^* -operad with operadic unit $\bullet \xrightarrow{\eta_{\hat{T}^*(\bullet)}} \hat{T}^*(\bullet)$ coinciding with the \hat{T}^* -monadic unit $\eta_{\hat{T}^*(\bullet)} := \eta_{\bullet}^{\hat{T}^*}$, and operadic multiplication


$$\begin{aligned} \hat{T}^*(\bullet) \circ_0^1 \hat{T}^*(\bullet) &\xrightarrow{\mu_{\hat{T}^*(\bullet)}} \hat{T}^*(\bullet) \text{ given by } \mu_{\hat{T}^*(\bullet)} := \nu \circ (\pi_{\hat{T}^*(\bullet)}, \mu_{\bullet}^{\hat{T}^*}), \\ \hat{T}^*(\bullet) \circ_0^1 \hat{T}^*(\bullet) &= \hat{T}^*(\bullet) \times_{\hat{T}^*(\bullet)} \hat{T}^*(\hat{T}^*(\bullet)) \\ &= \hat{T}^*(\bullet) \times_{\hat{T}^*(\bullet)} (\hat{T}^*)^2(\bullet) \xrightarrow{(\pi_{\hat{T}^*(\bullet)}, \mu_{\bullet}^{\hat{T}^*})} \hat{T}^*(\bullet) \times_{\hat{T}^*(\bullet)} \hat{T}^*(\bullet) \xrightarrow{\nu} \hat{T}^*(\bullet), \end{aligned}$$

where $\hat{T}^*(\bullet) \times_{\hat{T}^*(\bullet)} \hat{T}^*(\bullet) \xrightarrow{\nu} \hat{T}^*(\bullet)$ is an isomorphism and $(\hat{T}^*)^2(\bullet) \xrightarrow{\mu_{\bullet}^{\hat{T}^*}} \hat{T}^*(\bullet)$ is the \hat{T}^* -monadic multiplication.

To show that $\hat{T}^*(\bullet)$ is a \hat{T}^* -operad, one verifies (using the definitions (2.7) (2.5) (2.6) of associators and unitors via universal factorization property of pull-backs in the Cartesian category $\mathcal{Q}_{\hat{T}^*}$ and the equations (2.4) for the monad \hat{T}^*) the associativity and unitality properties already described in (2.3), here in the case of $\hat{T}^*(\bullet)$:

$$\begin{aligned} \mu_{\hat{T}^*(\bullet)} \circ_1^2 (\eta_{\hat{T}^*(\bullet)} \circ_0^2 \iota_{\hat{T}^*(\bullet)}^1) &= \nu \circ (\iota_{\hat{T}^*(\bullet)}, \mu_{\bullet}^{\hat{T}^*}) \circ (\eta_{\bullet}^{\hat{T}^*}, \iota_{\hat{T}^*(\bullet)}) = \lambda, \\ \mu_{\hat{T}^*(\bullet)} \circ_1^2 (\iota_{\hat{T}^*(\bullet)}^1 \circ_0^2 \eta_{\hat{T}^*(\bullet)}) &= \nu \circ (\mu_{\bullet}^{\hat{T}^*}, \iota_{\hat{T}^*(\bullet)}) \circ (\iota_{\hat{T}^*(\bullet)}, \eta_{\bullet}^{\hat{T}^*}) = \rho, \end{aligned}$$

can be respectively obtained by the commutativity of the following two diagrams and the unicity of λ and ρ ,



the operadic associativity of $\hat{T}^*(\bullet)$, that consists of the following identity

$$\mu_{\hat{T}^*(\bullet)} \circ_1^2 (\iota_{\hat{T}^*(\bullet)}^1 \circ_0^2 \mu_{\hat{T}^*(\bullet)}) \circ_1^2 \alpha = \mu_{\hat{T}^*(\bullet)} \circ_1^2 (\mu_{\hat{T}^*(\bullet)} \circ_0^2 \iota_{\hat{T}^*(\bullet)}^1),$$

can be obtained reconsidering the unicity of $\alpha := \alpha_{\hat{T}^*(\bullet)\hat{T}^*(\bullet)\hat{T}^*(\bullet)}$ in diagram (2.8),

reproduced here in our case,

$$\begin{array}{c}
\hat{T}^*(\bullet) \diamond \hat{T}^*(\hat{T}^*(\bullet) \diamond (\hat{T}^*)^2(\bullet)) \xrightarrow{\hat{T}^*(\pi_2) \circ_0^1 \pi_2} (\hat{T}^*)^3(\bullet) \xrightarrow{l_{(\hat{T}^*)^3(\bullet)}^1} (\hat{T}^*)^3(\bullet) \xrightarrow{\hat{T}^*(\mu_{\bullet}^{\hat{T}^*})} (\hat{T}^*)^2(\bullet) \\
\downarrow \scriptstyle{IP_1 \circ \pi_1} \quad \downarrow \scriptstyle{\mu_{\hat{T}^*(\bullet)}^{\hat{T}^*}} \quad \downarrow \scriptstyle{\mu_{\hat{T}^*(\bullet)}^{\hat{T}^*}} \quad \downarrow \scriptstyle{\mu_{\bullet}^{\hat{T}^*}} \\
\bullet \xrightarrow{\hat{\alpha}} (\hat{T}^*)^2(\bullet) \xrightarrow{l_{(\hat{T}^*)^2(\bullet)}^1} (\hat{T}^*)^2(\bullet) \xrightarrow{\mu_{\bullet}^{\hat{T}^*}} \hat{T}^*(\bullet) \\
\uparrow \scriptstyle{IP_1 \circ \pi_1 \circ \pi_1} \quad \nearrow \scriptstyle{\pi_2} \\
(\hat{T}^*(\bullet) \diamond (\hat{T}^*)^2(\bullet)) \diamond \hat{T}^*(\bullet)
\end{array}$$

and simply noting the structural properties of the multiplication maps involved:

$$\begin{array}{c}
\begin{array}{ccc}
(\hat{T}^*(\bullet) \diamond (\hat{T}^*)^2(\bullet)) \diamond \hat{T}^*(\bullet) & & (\hat{T}^*(\bullet) \diamond (\hat{T}^*)^2(\bullet)) \diamond \hat{T}^*(\bullet) \\
\downarrow \scriptstyle{\hat{T}^*(\bullet)} & \Downarrow \scriptstyle{\mu_{\hat{T}^*(\bullet)}^{\hat{T}^*} \circ_1^2 (\mu_{\hat{T}^*(\bullet)}^{\hat{T}^*} \circ_0^2 l_{\hat{T}^*(\bullet)}^1)} & \downarrow \scriptstyle{\mu_{\bullet}^{\hat{T}^*} \circ \pi_2} \\
\bullet & \hat{T}^*(\bullet) & \hat{T}^*(\bullet)
\end{array} \\
\hat{T}^*(\bullet) \diamond \hat{T}^*(\hat{T}^*(\bullet) \diamond (\hat{T}^*)^2(\bullet)) & & \hat{T}^*(\bullet) \diamond \hat{T}^*(\hat{T}^*(\bullet) \diamond (\hat{T}^*)^2(\bullet)) \\
\downarrow \scriptstyle{\hat{T}^*(\bullet)} & \Downarrow \scriptstyle{\mu_{\hat{T}^*(\bullet)}^{\hat{T}^*} \circ_1^2 (\mu_{\hat{T}^*(\bullet)}^{\hat{T}^*} \circ_0^2 l_{\hat{T}^*(\bullet)}^1)} & \downarrow \scriptstyle{\mu_{\bullet}^{\hat{T}^*} \circ \hat{T}^*(\mu_{\bullet}^{\hat{T}^*}) \circ (\pi_2 \circ_0^1 \hat{T}^*(\pi_2))} \\
\bullet & \hat{T}^*(\bullet) & \hat{T}^*(\bullet)
\end{array}$$

- for any (contracted) globular \hat{T}^* -operadic magma $(M, \pi_M, \eta_M, \mu_M)$, and hence for any (contracted) globular \hat{T}^* -operad, the projection $M \xrightarrow{\pi_M} \hat{T}^*(\bullet)$ is a morphism in $\mathcal{M}_{\bullet}^{\hat{T}^*}$ (and respectively in $\mathcal{M}_{\bullet}^{\hat{T}^*, K}$):

$$\begin{array}{ccc}
\bullet & \xrightarrow{\eta_M} & M \\
\downarrow \scriptstyle{!} & & \downarrow \scriptstyle{\pi_M} \\
\bullet & \xrightarrow{\eta_{\hat{T}^*(\bullet)}} & \hat{T}^*(\bullet)
\end{array}$$

where $\eta_{\hat{T}^*(\bullet)} := \eta_{\bullet}^{\hat{T}^*}$ and $\bullet \xrightarrow{!} \bullet$ is the terminal morphism;

$$\begin{array}{ccc}
M \circ_0^1 M & \xrightarrow{\mu_M} & M \\
\downarrow \scriptstyle{\pi_M \circ_0^2 \pi_M} & & \downarrow \scriptstyle{\pi_M} \\
\hat{T}^*(\bullet) \circ_0^1 \hat{T}^*(\bullet) & \xrightarrow{\mu_{\hat{T}^*(\bullet)}} & \hat{T}^*(\bullet)
\end{array}$$

with $\mu_{\hat{T}^*(\bullet)} := \nu \circ (\pi_{\hat{T}^*(\bullet)}, \mu_{\bullet}^{\hat{T}^*})$

and $\pi_M \circ_0^2 \pi_M : (x, y) \mapsto (\pi_M(x), \hat{T}^*(\pi_M)(y))$, where we have

$$M \circ_0^1 M = M \times_{\hat{T}^*(\bullet)} \hat{T}^*(M) \xrightarrow{\pi_M \circ_0^2 \pi_M} \hat{T}^*(\bullet) \times_{\hat{T}^*(\bullet)} \hat{T}^*(\hat{T}^*(\bullet)) = \hat{T}^*(\bullet) \circ_0^1 \hat{T}^*(\bullet).$$

The (contracted) operad $\hat{T}^*(\bullet)$ is final in $\mathcal{O}_{\bullet}^{\hat{T}^*, \kappa}$: for any other (contracted) \hat{T}^* -operad $(P, \pi^P, \mu^P, \eta^P, \kappa^P)$ the unique morphism of (contracted) \hat{T}^* -operads into $\hat{T}^*(\bullet)$ is given by the projection $P \xrightarrow{\pi^P} \hat{T}^*(\bullet)$.

Actually $\hat{T}^*(\bullet)$ is also a \hat{T}^* -operadic contraction since it furthermore satisfies, by direct computation, the following compatibility properties between contraction and operad structures (see diagrams (3.4) and (3.5)):

$$\mu_{\hat{T}^*(\bullet)} \circ (\kappa_{\hat{T}^*(\bullet)} \circ_0^2 \kappa_{\hat{T}^*(\bullet)}) = \kappa_{\hat{T}^*(\bullet)} \circ \text{Par}_{\mu_{\hat{T}^*(\bullet)}}, \quad \eta_{\hat{T}^*(\bullet)} \circ \kappa_{\pi_{\bullet}} = \kappa_{\hat{T}^*(\bullet)} \circ \text{Par}_{\eta_{\hat{T}^*(\bullet)}}.$$

- for any globular \hat{T}^* -operadic contraction magma $(M, \pi_M, \eta_M, \mu_M)$ (and for any globular \hat{T}^* -operadic contraction) the projection $M \xrightarrow{\pi_M} \hat{T}^*(\bullet)$ is a morphism in $\mathcal{M}\mathcal{K}_{\bullet}^{\hat{T}^*}$ (and respectively in $\mathcal{O}\mathcal{K}_{\bullet}^{\hat{T}^*}$):

$$\begin{array}{ccc}
 \text{Par}(\pi_{\bullet}) & \xrightarrow{\text{Par}_{\eta_M}} & \text{Par}(\pi_M) \\
 \downarrow \text{Par}_{\kappa_{\bullet}} & \swarrow \kappa_{\bullet} & \searrow \kappa_M \\
 \bullet & \xrightarrow{\eta_M} & M \\
 \downarrow \text{Par}_{\kappa_{\hat{T}^*(\bullet)}} & \downarrow ! & \downarrow \pi_M \\
 \bullet & \xrightarrow{\eta_{\hat{T}^*(\bullet)}} & \hat{T}^*(\bullet) \\
 \downarrow \text{Par}_{\kappa_{\pi_{\bullet}}} & \swarrow \kappa_{\hat{T}^*(\bullet)} & \downarrow \text{Par}_{\kappa_{\hat{T}^*(\bullet)}} \\
 \text{Par}(\pi_{\bullet}) & \xrightarrow{\text{Par}_{\eta_{\hat{T}^*}}} & \text{Par}(\pi_{\hat{T}^*(\bullet)})
 \end{array}$$

where:

(3.4)

$$\text{Par}(\pi_{\bullet}) = \left\{ (x^+, y, x^-) \mid \begin{array}{c} x^+ \\ \Downarrow y \\ x^- \end{array} \in \bullet \right\},$$

$$\text{Par}_{\eta_{\hat{T}^*}} : (x^+, y, x^-) \mapsto (x^+, \eta^{\hat{T}^*}(y), x^-),$$

$$\kappa_{\hat{T}^*(\bullet)} : (x^+, \eta^{\hat{T}^*}(y), x^-) \mapsto \eta^{\hat{T}^*}(y), \quad \kappa_{\pi_{\bullet}} : (x^+, y, x^-) \mapsto y;$$

$$\begin{array}{ccc}
\text{Par}(\pi_M \circ_0^2 \pi_M) & \xrightarrow{\text{Par}_{\mu_M}} & \text{Par}(\pi_M) \\
\downarrow \text{Par}_{\pi_M \circ_0^2 \pi_M} & \searrow \kappa_M \circ_0^2 \kappa_M & \swarrow \kappa_M \\
& M \circ_0^1 M \xrightarrow{\mu_M} M & \\
& \downarrow \pi_M \circ_0^2 \pi_M & \downarrow \pi_M \\
& \hat{T}^*(\bullet) \circ_0^1 \hat{T}^*(\bullet) \xrightarrow{\mu_{\hat{T}^*(\bullet)}} \hat{T}^*(\bullet) & \\
\downarrow \text{Par}_{\pi_{\hat{T}^*(\bullet)} \circ_0^2 \pi_{\hat{T}^*(\bullet)}} & \swarrow \kappa_{\hat{T}^*(\bullet)} \circ_0^2 \kappa_{\hat{T}^*(\bullet)} & \searrow \kappa_{\hat{T}^*(\bullet)} \\
\text{Par}(\pi_{\hat{T}^*(\bullet)} \circ_0^2 \pi_{\hat{T}^*(\bullet)}) & \xrightarrow{\text{Par}_{\mu_{\hat{T}^*(\bullet)}}} & \text{Par}(\pi_{\hat{T}^*(\bullet)}).
\end{array}
\tag{3.5}$$

$$\begin{aligned}
& \text{Par}(\pi_{\hat{T}^*(\bullet)} \circ_0^2 \pi_{\hat{T}^*(\bullet)}) \simeq \text{Par}(\pi_{\hat{T}^*(\bullet)}) \circ_0^1 \text{Par}(\pi_{\hat{T}^*(\bullet)}), \\
& \kappa_{\hat{T}^*(\bullet)} : (\mu(x_1^+, x_2^+), \mu(y_1, y_2), \mu(x_1^-, x_2^-)) \mapsto \mu(y_1, y_2), \\
& \kappa_{\mu_{\hat{T}^*(\bullet)}} : (y_1, y_2) \mapsto \mu(y_1, y_2). \\
& \kappa_{\hat{T}^*(\bullet)} \circ_0^2 \kappa_{\hat{T}^*(\bullet)} : ((x_1^+, y_1, x_1^-), (x_1^+, y_1, x_1^-)) \mapsto (y_1, y_2), \\
& \text{Par}_{\mu_{\hat{T}^*(\bullet)}} : ((x_1^+, y_1, x_1^-), (x_1^+, y_1, x_1^-)) \mapsto (\mu(x_1^+, x_2^+), \mu(y_1, y_2), \mu(x_1^-, x_2^-)).
\end{aligned}$$

The \hat{T}^* -operadic magma contraction (respectively the \hat{T}^* -operadic contraction) $\hat{T}^*(\bullet)$ is final in $\mathcal{M}\mathcal{H}_{\bullet}^{\hat{T}^*}$ (respectively in $\mathcal{O}\mathcal{H}_{\bullet}^{\hat{T}^*}$): the projection $M \xrightarrow{\pi_M} \hat{T}^*(\bullet)$ being the terminal morphism from any other object M . \lrcorner

The following is the fundamental theorem in our paper, allowing the definition of weak involutive ω -categories.

Theorem 3.12. *The category $\mathcal{O}_{\bullet}^{\hat{T}^*, \kappa}$ of globular contracted \hat{T}^* -operads has initial objects.*

Proof. Instead of following Leinster's original line of proof in section 2.3.3, we give a direct argument.

- The category $\mathcal{Q}_{\bullet}^{\hat{T}^*}$ has an initial object I : the empty \hat{T}^* -collection $I \xrightarrow{\pi} \hat{T}^*(\bullet)$ given by $I^n := \emptyset$, for all $n \in \mathbb{N}$, where all the source/target maps and the projection π are empty functions.
- Left-adjoint functors preserve colimits (see [Riehl 2016, theorem 4.5.3]) and hence they preserve initial objects (that are colimits of the empty diagram).

- Hence, if $\mathfrak{U}_\mathcal{O}^* \circ \hat{\mathfrak{U}}_k^* = \mathfrak{U}_k^* \circ \hat{\mathfrak{U}}_\mathcal{O}^*$ has a left-adjoint $\mathcal{Q}_\bullet^{\hat{T}^*} \xrightarrow{\zeta} \mathcal{O}_\bullet^{\hat{T}^*,k}$, the object $L^* := \mathcal{Q}^0(I)$ is initial in $\mathcal{O}_\bullet^{\hat{T}^*,k}$.

The theorem is now reduced to providing the existence of a *free contracted \hat{T}^* -operad over a \hat{T}^* -collection*. This is achieved below, in the theorem 3.16, by an argument substantially similar to that used in our construction of the free self-dual Penon's contractions in [Bejrakarbum Bertozzini 2017, proposition 3.3]. \square

First we need to define suitable free structures over \hat{T}^* -collections. ¹⁸

Definition 3.13. A *free globular contracted \hat{T}^* -operad* (P, ζ) over a \hat{T}^* -collection $Q := Q \xrightarrow{\pi} \hat{T}^*(\bullet)$ consists of a contracted \hat{T}^* -operad $P := (P, \pi_P, \kappa_P, \mu_P, \eta_P)$ and a morphism $Q \xrightarrow{\zeta} \mathfrak{U}_k^* \circ \hat{\mathfrak{U}}_\mathcal{O}^*$ in $\mathcal{Q}_\bullet^{\hat{T}^*}$ that satisfies the following universal factorization property: for every other morphism $Q \xrightarrow{\phi} \mathfrak{U}_k^* \circ \hat{\mathfrak{U}}_\mathcal{O}^*(\hat{P})$ in $\mathcal{Q}_\bullet^{\hat{T}^*}$, where \hat{P} is another contracted \hat{T}^* -operad, there exists a unique morphism $P \xrightarrow{\hat{\phi}} \hat{P}$ in $\mathcal{O}_\bullet^{\hat{T}^*,k}$ such that $\phi = \hat{\phi} \circ \zeta$.

A *free globular contracted \hat{T}^* -operadic magma* $M := (M, \pi_M, \kappa_M, \mu_M, \eta_M)$ over a \hat{T}^* -collection, is defined in a similar way, via a morphism $Q \xrightarrow{\xi_Q} \mathfrak{U}_k^* \circ \hat{\mathfrak{U}}_{\mathcal{M}}^*(M)$, substituting “operads” with “operadic magmas” above.

As in any universal factorization property construct, free contracted \hat{T}^* -operad(ic magma)s are unique, modulo a unique isomorphism compatible with the universal factorization property. Existence is shown in theorem 3.16.

Before embarking on the proof, we need to introduce relevant notions of *congruence* and *quotient structure*.

Definition 3.14. A *globular ω -equivalence relation* is an equivalence relation \mathcal{E} in a globular ω -set Q that is graded ¹⁹ $\mathcal{E} \subset Q \times_{\mathbb{N}} Q := \{(x, y) \in Q \times Q \mid \exists n \in \mathbb{N} : x, y \in Q^n\}$ and source/target preserving: ²⁰

$$(x_1, x_2) \in \mathcal{E} \Rightarrow (s(x_1), s(x_2)) \in \mathcal{E}, \quad (x_1, x_2) \in \mathcal{E} \Rightarrow (t(x_1), t(x_2)) \in \mathcal{E}, \quad \forall x, y \in Q. \quad (3.6)$$

A \hat{T}^* -collection *congruence* is a globular ω -equivalence relation \mathcal{E} in the globular ω -set Q of a globular \hat{T}^* -collection $Q \xrightarrow{\pi} \hat{T}^*(\bullet)$ that is projection-preserving:

$$(x, y) \in \mathcal{E} \Rightarrow \pi(x) = \pi(y)$$

¹⁸Of course the definitions, that for convenience are here stated for the specific case of \hat{T}^* , work for any Cartesian monad T .

¹⁹This is equivalent to say that \mathcal{E} consists of a sequence $\mathcal{E}^n \subset Q^n \times Q^n$ of equivalence relations in Q^n , for all $n \in \mathbb{N}$.

²⁰In this way, the globular source and target product maps $(s, s), (t, t) : Q \times_{\mathbb{N}} Q \rightarrow Q \times_{\mathbb{N}} Q$ restrict to (necessarily globular) source and target maps on \mathcal{E} and hence $(\mathcal{E}, (s, s)|_{\mathcal{E}}^{\mathcal{E}}, (t, t)|_{\mathcal{E}}^{\mathcal{E}})$ becomes a globular ω -set canonically included in $Q \times_{\mathbb{N}} Q$.

and hence $\mathcal{E} \subset Q \times_{\pi} Q$.²¹

A **congruence of contracted \hat{T}^* -collection** is a \hat{T}^* -collection congruence \mathcal{E} in a contracted \hat{T}^* -collection $\text{Par}(\pi) \xrightarrow{\kappa} Q \xrightarrow{\pi} \hat{T}^*(\bullet)$ that is also contraction-preserving:²²

$$\begin{aligned} \forall (x_1^+, x_2^+), (x_1^-, x_2^-) \in \mathcal{E} : \\ (x_1^+, y_1, x_1^-), (x_2^+, y_2, x_2^-) \in \text{Par}(\pi) \Rightarrow (\kappa(x_1^+, y_1, x_1^-), \kappa(x_2^+, y_2, x_2^-)) \in \mathcal{E}. \end{aligned} \quad (3.7)$$

A **congruence of (contracted) \hat{T}^* -operadic magma** (M, μ_M, η_M) is a congruence \mathcal{E} of the underlying (contracted) \hat{T}^* -collection $\text{Par}(\pi) \xrightarrow{\kappa_M} M \xrightarrow{\pi_M} \hat{T}^*(\bullet)$ that is unit-preserving and multiplication-preserving:²³

$$(x, y) \in \mathcal{E} \circ_0^1 \mathcal{E} = \mathcal{E} \times_{\hat{T}^*(\bullet)} \hat{T}^*(\mathcal{E}) \quad \Rightarrow \quad ((\mu_M, \mu_M) \circ \tau_M \circ (\mathcal{E} \circ_0^1 \mathcal{E}))(x, y) \in \mathcal{E}, \quad (3.8)$$

$$\begin{array}{ccc} \mathcal{E} \circ_0^1 \mathcal{E} & \xrightarrow{\mathcal{E} \circ_0^2 \mathcal{E}} & (M \times_{\pi_M} M) \circ_0^1 (M \times_{\pi_M} M) \xrightarrow{\tau_M} (M \circ_0^1 M) \times_{\pi_M \circ_0^2 \pi_M} (M \circ_0^1 M) \\ & & \downarrow (\mu_M, \mu_M) \\ & & M \times_{\pi_M} M \end{array}$$

with τ_M denoting the canonical isomorphism of \hat{T}^* -collections between pull-back of products and product of pull-backs in $\mathcal{Q}_{\bullet}^{\hat{T}^*}$ and $!$ as the unique isomorphism of terminal objects.²⁴

The previous congruences have been defined, for our convenience, for 1-cells in $\mathcal{Q}_{\bullet}^{\hat{T}^*}$, but they actually work for 1-cells in the bicategory \mathcal{Q}_T , where T is any Cartesian monad on the category of globular ω -sets \mathcal{Q} .

As usual, congruences produce quotients of the corresponding algebraic structures.

²¹This means that the globular ω -set $(\mathcal{E}, (s, s), (t, t))$ is a \hat{T}^* -collection $\mathcal{E} \xrightarrow{\pi_{\mathcal{E}}} \hat{T}^*(\bullet)$ equipped with the projection $\pi_{\mathcal{E}} : (x, y) \mapsto \pi(x) = \pi(y)$; and we have a canonical inclusion morphism of \hat{T}^* -collections $\mathcal{E} \xrightarrow{\mathcal{E}_{\mathcal{E}}} Q \times_{\pi} Q$ into the **product \hat{T}^* -collection** $Q \times_{\pi} Q \xrightarrow{(\pi, \pi)} \hat{T}^*(\bullet)$.

²²This entails that the \hat{T}^* -collection $\mathcal{E} \xrightarrow{\pi_{\mathcal{E}}} \hat{T}^*(\bullet)$ has a contraction $\kappa_{\mathcal{E}} : \text{Par}(\pi_{\mathcal{E}}) \rightarrow \mathcal{E}$ with $\kappa_{\mathcal{E}} := (\kappa, \kappa)$.

²³The congruence \mathcal{E} is always unit-preserving since the product morphism $\bullet \times_{\eta_{\hat{T}^*}} \bullet \xrightarrow{(\eta_M, \eta_M)} M \times_{\pi_M} M$ has always image inside \mathcal{E} ; hence \mathcal{E} is equipped with a canonical unit $\bullet \xrightarrow{\eta_{\mathcal{E}}} \mathcal{E}$ given by composing the maps $\bullet \xrightarrow{!} \bullet \times_{\eta_{\hat{T}^*}} \bullet \xrightarrow{(\eta_M, \eta_M)^{\mathcal{E}}} \mathcal{E}$ with isomorphism $!$ of terminal objects.

²⁴In this way, denoting by $\mathcal{E} \circ_0^1 \mathcal{E} \xrightarrow{\mu_{\mathcal{E}}} \mathcal{E}$ the restriction $\mu_{\mathcal{E}} := (\mu_M, \mu_M) \circ \tau_M \circ (\mathcal{E} \circ_0^1 \mathcal{E})|_{\mathcal{E} \circ_0^1 \mathcal{E}}$ and by $\bullet \xrightarrow{\eta_{\mathcal{E}}} \mathcal{E}$ the restriction $\eta_{\mathcal{E}} := (\eta_M, \eta_M) \circ !|_{\mathcal{E}}$, we have that $(\mathcal{E}, \eta_{\mathcal{E}}, \mu_{\mathcal{E}})$ is itself a (contracted) \hat{T}^* -operadic magma and that the inclusion $\mathcal{E} \xrightarrow{\mathcal{E}} M \times_{\pi_M} M$ is a morphism of (contracted) \hat{T}^* operadic magmas. Whenever $(M, \pi_M, \kappa_M, \eta_M, \mu_M)$ is a (contracted) \hat{T}^* -operad, also $(\mathcal{E}, \pi_{\mathcal{E}}, \kappa_{\mathcal{E}}, \eta_{\mathcal{E}}, \mu_{\mathcal{E}})$ is.

Proposition 3.15. *Given a globular ω -relation $\mathcal{E} \subset Q \times_{\mathbb{N}} Q$ on a globular ω -set Q , the family of quotients $Q/\mathcal{E} := (Q^n/\mathcal{E}^n)_{n \in \mathbb{N}}$ becomes a globular ω -set with well-defined sources and targets $s_{Q/\mathcal{E}}([x]_{\mathcal{E}}) := [s_Q(x)]$, $t_{Q/\mathcal{E}}([x]_{\mathcal{E}}) := [t_Q(x)]_{\mathcal{E}}$; furthermore the quotient map $Q \xrightarrow{\varpi_{\mathcal{E}}} Q/\mathcal{E}$, defined as usual by $\varpi_{\mathcal{E}} : x \mapsto [x]_{\mathcal{E}}$, is a morphism of globular ω -sets.*

Given a congruence $\mathcal{E} \subset Q \times_{\pi} Q$ of (contracted) \hat{T}^ -collection $Q \xrightarrow{\pi} \hat{T}^*(\bullet)$, the quotient globular ω -set Q/\mathcal{E} becomes a (contracted) \hat{T}^* -collection $Q/\mathcal{E} \xrightarrow{\pi_{Q/\mathcal{E}}} \hat{T}^*(\bullet)$ with projection $\pi_{Q/\mathcal{E}} : [x]_{\mathcal{E}} \mapsto \pi(x)$ (contraction $\kappa_{Q/\mathcal{E}} : ([x^+]_{\mathcal{E}}, [x^-]_{\mathcal{E}}) \mapsto [\kappa(x^+, y, x^-)]_{\mathcal{E}}$ on $\text{Par}(\pi_{Q/\mathcal{E}}) = \{([x^+]_{\mathcal{E}}, y, [x^-]_{\mathcal{E}}) \mid (x^+, y, x^-) \in \text{Par}(\pi)\}$); furthermore the quotient map $Q \xrightarrow{\varpi_{\mathcal{E}}} Q/\mathcal{E}$ is a morphism of (contracted) \hat{T}^* -collections.*

Given $\mathcal{E} \subset M \times_{\pi_M} M$, a congruence on $(M, \pi_M, \eta_M, \mu_M)$, a (contracted) \hat{T}^ -operadic magma, the quotient (contracted) \hat{T}^* -collection $M/\mathcal{E} \xrightarrow{\pi_{M/\mathcal{E}}} \hat{T}^*(\bullet)$ becomes a (contracted) \hat{T}^* -operadic magma with operadic unit $\bullet \xrightarrow{\eta_{M/\mathcal{E}}} M/\mathcal{E}$ given by $\eta_{M/\mathcal{E}} := \pi_{\mathcal{E}} \circ \eta_M$ and with operadic multiplication $M/\mathcal{E} \circ_0^1 M/\mathcal{E} \xrightarrow{\mu_{M/\mathcal{E}}} M/\mathcal{E}$ that is well-defined by $\mu_{M/\mathcal{E}} : ([x]_{\mathcal{E}}, [y]_{\mathcal{E}}) \mapsto [\mu_M(x, y)]_{\mathcal{E}}$, for all $(x, y) \in M \circ_0^1 M = M \times_{\hat{T}^*(\bullet)} \hat{T}^*(M)$; furthermore the quotient map $M \xrightarrow{\varpi_{\mathcal{E}}} M/\mathcal{E}$ is a morphism of (contracted) \hat{T}^* -operadic magmas.*

Given a morphism $Q_1 \xrightarrow{\phi} Q_2$ of the categories in proposition 3.9, a congruence \mathcal{E} of the respective type in Q_2 naturally induces a congruence, of the same type, in Q_1 :

$$\mathcal{E}_{\phi} := \{(x, y) \in Q_1 \times Q_1 \mid (\phi(x), \phi(y)) \in \mathcal{E}\}.$$

Furthermore if \mathcal{E}_1 is a congruence of the respective type in Q_1 such that $\mathcal{E}_1 \subset \mathcal{E}_{\phi}$, there exists a unique well-defined quotient morphism $Q_1/\mathcal{E}_1 \xrightarrow{\hat{\phi}} Q_2$, well-defined by $\hat{\phi} : [x]_{\mathcal{E}_1} \mapsto \phi(x)$, such that $\phi = \hat{\phi} \circ \pi_{\mathcal{E}_1}$.

Proof. By property (3.6), the source and target $s_{Q/\mathcal{E}}^n, t_{Q/\mathcal{E}}^n : Q^{n+1}/\mathcal{E}^{n+1} \rightarrow Q^n/\mathcal{E}^n$, for all $n \in \mathbb{N}$, are well-defined and their globularity property follows from the globularity of Q . For all $n \in \mathbb{N}$, the quotient function $\varpi_{\mathcal{E}}^n : Q^n \rightarrow Q^n/\mathcal{E}^n$ is well-defined by $x \mapsto [x]_{\mathcal{E}^n}$; and $\varpi_{\mathcal{E}} := (\pi_{\mathcal{E}}^n)_{\mathbb{N}} : Q \rightarrow Q/\mathcal{E}$ becomes a morphism in \mathcal{Q} since we have $s_{Q/\mathcal{E}}^n(\varpi_{\mathcal{E}}^{n+1}(x)) = s_{Q/\mathcal{E}}^n([x]_{\mathcal{E}^{n+1}}) = [s_Q^n(x)]_{\mathcal{E}^n} = \varpi_{\mathcal{E}}^n(s_Q^n(x))$, for all $x \in Q^{n+1}$, and similarly for targets.

A congruence $\mathcal{E} \subset Q \times_{\pi} Q$ of a collection $Q \xrightarrow{\pi} \hat{T}^*(\bullet)$ is necessarily a congruence of globular ω -sets, hence we already have a quotient morphism $Q \xrightarrow{\varpi_{\mathcal{E}}} Q/\mathcal{E}$ in \mathcal{Q} . Since $\mathcal{E} \subset Q \times_{\pi} Q$, the assignment $[x]_{\mathcal{E}} \mapsto \pi(x)$ is a well-defined graded function $\pi_{Q/\mathcal{E}} : Q/\mathcal{E} \rightarrow \hat{T}^*(\bullet)$ that is actually a morphism of globular ω -sets in \mathcal{Q} : for all $[x]_{\mathcal{E}} \in Q/\mathcal{E}$, $\pi_{Q/\mathcal{E}}(s_{Q/\mathcal{E}}[x]_{\mathcal{E}}) = \pi_{Q/\mathcal{E}}([s_Q(x)]_{\mathcal{E}}) = \pi(s_Q(x)) = s(\pi(x)) = s_{\hat{T}^*(\bullet)}(\pi_{Q/\mathcal{E}}[x]_{\mathcal{E}})$ and similarly for targets. Since $\pi_{Q/\mathcal{E}} \circ \varpi_{\mathcal{E}}(x) = \pi_{Q/\mathcal{E}}([x]_{\mathcal{E}}) = \pi(x)$, for all $x \in Q$, the quotient $Q \xrightarrow{\varpi_{\mathcal{E}}} Q/\mathcal{E}$ is a morphism of \hat{T}^* -collections in $\mathcal{Q}^{\hat{T}^*}$.

Suppose now that $\text{Par}(\pi) \xrightarrow{\kappa} Q \xrightarrow{\pi} \hat{T}^*(\bullet)$ is a contracted \hat{T}^* -collection and $\mathcal{E} \subset Q \times_{\pi} Q$ is a congruence of contracted \hat{T}^* -collections. Considering the quotient \hat{T}^* -collection $Q/\mathcal{E} \xrightarrow{\pi_{Q/\mathcal{E}}} \hat{T}^*(\bullet)$, we have that $[x_1]_{\mathcal{E}}, [x_2]_{\mathcal{E}} \in Q/\mathcal{E}$ are $\pi_{Q/\mathcal{E}}$ -parallel if and only if x_1, x_2 are π -parallel and hence $\text{Par}(\pi_{Q/\mathcal{E}}) = \{([x^+]_{\mathcal{E}}, y, [x^-]_{\mathcal{E}}) \mid (x^+, y, x^-) \in \text{Par}(\pi)\}$. Furthermore, since $\text{Par}_{\pi_{Q/\mathcal{E}}} : (x^+, y, x^-) \mapsto ([x^+]_{\mathcal{E}}, y, [x^-]_{\mathcal{E}})$ there exists a unique relation $\text{Par}(\pi_{Q/\mathcal{E}}) \xrightarrow{\kappa_{Q/\mathcal{E}}} Q/\mathcal{E}$ that satisfies $\text{Par}_{\pi_{\mathcal{E}}} \circ \kappa_{Q/\mathcal{E}} = \kappa$ and that is given by

$$\kappa_{Q/\mathcal{E}}([x^+]_{\mathcal{E}}, y, [x^-]_{\mathcal{E}}) := [\kappa(x^+, y, x^-)]_{\mathcal{E}}.$$

Since \mathcal{E} is a congruence of contracted \hat{T}^* -collections, from equation (3.7) we see that $\kappa_{Q/\mathcal{E}}$ is actually a well-defined function and hence a contraction on Q/\mathcal{E} and the map $Q \xrightarrow{\varpi_{\mathcal{E}}} Q/\mathcal{E}$ is a morphism in $\mathcal{Q}_{\bullet}^{\hat{T}^*, \kappa}$.

Let \mathcal{E} be a congruence of the (contracted) operadic magma $(M, \pi_M, \kappa_M, \eta_M, \mu_M)$. We already know that the quotient $(M/\mathcal{E}, \pi_{M/\mathcal{E}}, \kappa_{M/\mathcal{E}})$ is a (contracted) \hat{T}^* -collection and that $M \xrightarrow{\varpi_{\mathcal{E}}} M/\mathcal{E}$ is a morphism in $\mathcal{Q}_{\bullet}^{\hat{T}^*, \kappa}$.

The operadic unit map $\eta_{M/\mathcal{E}} : \bullet \mapsto [\eta_M(\bullet)]_{\mathcal{E}}$ is well defined, and we immediately get $\eta_{M/\mathcal{E}} = \pi_{M/\mathcal{E}} \circ \eta_M$, hence the quotient morphism $\pi_{M/\mathcal{E}}$ preserves units.

To describe the operadic multiplication, we first notice that we have a canonical exchange isomorphism $(M/\mathcal{E}) \circ_0^1 (M/\mathcal{E}) \xrightarrow{\chi} (M \circ_0^1 M)/(\mathcal{E} \circ_0^1 \mathcal{E})$; any multiplication morphism $(M/\mathcal{E}) \circ_0^1 (M/\mathcal{E}) \xrightarrow{\mu_{M/\mathcal{E}}} M/\mathcal{E}$ such that $\mu_{M/\mathcal{E}} \circ (\varpi_{\mathcal{E}} \circ_0^2 \varpi_{\mathcal{E}}) = \varpi_{\mathcal{E}} \circ \mu_M$, must necessarily be given by $\mu_{M/\mathcal{E}} := \hat{\mu}_{M/\mathcal{E}} \circ \chi$ where, using the congruence property (3.8) of \mathcal{E} , we have that $\hat{\mu}_{M/\mathcal{E}} : [(x, y)]_{\mathcal{E} \circ_0^1 \mathcal{E}} \mapsto \varpi_{\mathcal{E}} \circ \mu_M(x, y)$, for $(x, y) \in M \circ_0^1 M$, is a well-defined morphism of (contracted) \hat{T}^* -collections.

Since $\mu_{M/\mathcal{E}} \circ (\varpi_{\mathcal{E}} \circ_0^2 \varpi_{\mathcal{E}}) = \hat{\mu}_{M/\mathcal{E}} \circ \chi \circ (\varpi_{\mathcal{E}} \circ_0^2 \varpi_{\mathcal{E}}) = \hat{\mu}_{M/\mathcal{E}} \circ \varpi_{\mathcal{E} \circ_0^1 \mathcal{E}} = \varpi_{\mathcal{E}} \circ \mu_M$, we see that $\pi_{\mathcal{E}}$ is a morphism of (contracted) \hat{T}^* -operadic magmas.

The family $\mathcal{E}_{\phi} \subset Q_1 \times Q_1$ is an equivalence relation in Q_1 and, since ϕ is grade-preserving, we also have $\mathcal{E}_{\phi} \subset Q_1 \times_{\mathbb{N}} Q_1$ and hence \mathcal{E}_{ϕ} consists of a family of equivalence relations $\mathcal{E}_{\phi}^n \subset Q_1^n \times Q_1^n$, for all $n \in \mathbb{N}$. Since ϕ is always a morphism of globular ω -sets, $s_{Q_2}^n \circ \phi^{n+1} = \phi^n \circ s_{Q_1}^n$ (and similarly for targets), for all $n \in \mathbb{N}$, and hence property (3.6) holds and \mathcal{E}_{ϕ} is a globular ω -equivalence relation in Q_1 .

If $Q_1 \xrightarrow{\phi} Q_2$ is a morphism of \hat{T}^* -collections in $\mathcal{Q}_{\bullet}^{\hat{T}^*}$,

$$(x, y) \in \mathcal{E}_{\phi} \Rightarrow \pi_{Q_2}(\phi(x)) = \pi_{Q_2}(\phi(y)) \Rightarrow \pi_{Q_1}(x) = \pi_{Q_1}(y)$$

and hence $\mathcal{E}_{\phi} \subset Q_1 \times_{\pi_{Q_1}} Q_1$ is a congruence of \hat{T}^* -collections in Q_1 .

Suppose that ϕ is a morphism of contracted \hat{T}^* -collections in $\mathcal{Q}_{\bullet}^{\hat{T}^*, \kappa}$, consider a pair of elements $(x_1^+, y_1, x_1^-), (x_2^+, y_2, x_2^-) \in \text{Par}(\pi_1)$ with $(x_1^+, x_2^+), (x_1^-, x_2^-) \in \mathcal{E}_{\phi}$, we must show

$$(\kappa_1(x_1^+, y_1, x_1^-), \kappa_1(x_2^+, y_2, x_2^-)) \in \mathcal{E}_{\phi}$$

and this is equivalent to prove that

$$(\kappa_2 \circ \text{Par}_\phi(x_1^+, y_1, x_1^-), \kappa_2 \circ \text{Par}(x_1^+, y_1, x_1^-)) = (\phi \circ \kappa_1(x_1^+, y_1, x_1^-), \phi \circ \kappa_1(x_1^+, y_1, x_1^-)) \in \mathcal{E}.$$

This final statement is true since we have $(\phi(x_1^+), y_1, \phi(x_1^-)), (\phi(x_2^+), y_2, \phi(x_2^-)) \in \text{Par}(\pi_2)$ and $(\phi(x_1^+), \phi(x_2^+)), (\phi(x_1^-), \phi(x_2^-)) \in \mathcal{E}$ and hence also

$$\begin{aligned} & (k_2(\phi(x_1)^+, y_1, \phi(x_1)^-), \kappa_2(\phi(x_2)^+, y_2, \phi(x_2)^-)) \\ &= (\kappa_2 \circ \text{Par}_\phi(x_1^+, y_1, x_1^-), \kappa_2 \circ \text{Par}(x_1^+, y_1, x_1^-)) \in \mathcal{E}. \end{aligned}$$

This shows that \mathcal{E}_ϕ is also a congruence of contracted \hat{T}^* -collection in Q_1 .

Finally if \mathcal{E} is a congruence of (contracted) \hat{T}^* -operadic magma in Q_2 and $Q_1 \xrightarrow{\phi} Q_2$ is a morphism in $\mathcal{M}_\bullet^{\hat{T}^*}$:

$$\begin{aligned} (x, y) \in \mathcal{E}_\phi \circ_0^1 \mathcal{E}_\phi &\Rightarrow (\phi \circ_0^2 \phi)(x, y) \in \mathcal{E} \circ_0^1 \mathcal{E} \Rightarrow \mu_{\mathcal{E}} \circ (\phi \circ_0^2 \phi)(x, y) \in \mathcal{E} \\ &\Rightarrow (\phi \circ_0^2 \phi)(\mu_{\mathcal{E}}(x, y)) \in \mathcal{E} \Rightarrow \mu_{\mathcal{E}}(x, y) \in \mathcal{E}_\phi, \end{aligned}$$

hence \mathcal{E}_ϕ is a congruence \hat{T}^* -operadic magmas. The same argument assures that, if ϕ is a morphism in $\mathcal{M}_\bullet^{\hat{T}^*, \kappa}$, \mathcal{E}_ϕ is a congruence of contracted \hat{T}^* -operadic magmas.

Finally, whenever $\mathcal{E}_1 \subset \mathcal{E}_\phi$ is a congruence (of the ‘‘same type’’ of the morphism ϕ) in Q_1 , any relation $Q_1/\mathcal{E}_1 \xrightarrow{\hat{\phi}} Q_2$ such that $\phi = \hat{\phi} \circ \pi_{\mathcal{E}_1}$ must necessarily associate $[x]_{\mathcal{E}_1} \mapsto \phi(x)$ and this is a well defined function since $[x]_{\mathcal{E}_1} \subset [x]_{\mathcal{E}_\phi}$ and hence $\phi(x)$ does not depend on the representative element.

We must show that $Q_1/\mathcal{E}_1 \xrightarrow{\hat{\phi}} Q_2$ is a morphism in the same category of ϕ . From its definition $\hat{\phi}$ is already a graded map: $\hat{\phi}(Q_1^n/\mathcal{E}_1^n) \subset Q_2^n$, for all $n \in \mathbb{N}$. For all $x \in Q_1$, we have $s_{Q_2}(\hat{\phi}([x]_{\mathcal{E}_1})) = s_{Q_2}(\phi(x)) = \phi(s_{Q_1}(x)) = \hat{\phi}([s_{Q_1}(x)]_{\mathcal{E}_1}) = \hat{\phi}(s_{Q_1/\mathcal{E}_1}([x]_{\mathcal{E}_1}))$ and similarly for the target; hence $\hat{\phi}$ is a morphism in \mathcal{Q} .

Since $\pi_{Q_2}(\hat{\phi}([x]_{\mathcal{E}_1})) = \pi_{Q_2}(\phi(x)) = \pi_{Q_1}(x) = \pi_{Q_1/\mathcal{E}_1}([x]_{\mathcal{E}_1})$, for all $x \in Q_1$, we also have that $\hat{\phi}$ is a morphism in $\mathcal{Q}_\bullet^{\hat{T}^*}$. Since $\hat{\phi}$ is already a morphism of \hat{T}^* -collections it naturally induces a map $\text{Par}_{\hat{\phi}} : \text{Par}(\pi_{Q_1/\mathcal{E}_1}) \rightarrow \text{Par}(\pi_{Q_2})$ and, whenever ϕ is a morphism in $\mathcal{Q}^{\hat{T}^*, \kappa}$, we show that $\kappa_{Q_2} \circ \text{Par}_{\hat{\phi}} = \hat{\phi} \circ \kappa_{Q_1/\mathcal{E}_1}$:

$$\begin{aligned} \kappa_{Q_2}(\text{Par}_{\hat{\phi}}([x^+]_{\mathcal{E}_1}, y, [x^-]_{\mathcal{E}_1})) &= \kappa_{Q_2}(\text{Par}_{\hat{\phi}} \circ \text{Par}_{\pi_{\frac{Q_1}{\mathcal{E}_1}}}(x^+, y, x^-)) \\ &= \kappa_{Q_2}(\text{Par}_\phi(x^+, y, x^-)) = \phi(\kappa_{Q_1}(x^+, y, x^-)) \\ &= \hat{\phi}(\pi_{\frac{Q_1}{\mathcal{E}_1}}(x^+, y, x^-)) = \hat{\phi}(\kappa_{\frac{Q_1}{\mathcal{E}_1}}([x^+]_{\mathcal{E}_1}, y, [x^-]_{\mathcal{E}_1})), \end{aligned}$$

for all $(x^+, y, x^-) \in \text{Par}(\pi_{Q_1})$, and hence $\hat{\phi}$ is a morphism in $\mathcal{Q}_\bullet^{\hat{T}^*}$.

Supposing now that $Q_1 \xrightarrow{\phi} Q_2$ is a morphism in $\mathcal{M}_\bullet^{\hat{T}^*}$ and \mathcal{E}_1 is a congruence of \hat{T}^* -operadic magmas, we have $\hat{\phi} \circ \eta_{Q_1/\mathcal{E}_1} = \hat{\phi} \circ \pi_{Q_1/\mathcal{E}_1} \circ \eta_{Q_1} = \phi \circ \eta_{Q_1} = \eta_{Q_2}$. As regards

multiplication, from $\phi = \hat{\phi} \circ \pi_{Q_1/\mathcal{E}_1}$ we obtain $(\phi \circ_0^2 \phi) = (\hat{\phi} \circ_0^2 \hat{\phi}) \circ (\pi_{Q_1/\mathcal{E}_1} \circ_0^2 \pi_{Q_1/\mathcal{E}_1})$; since $\mu_{Q_2} \circ (\phi \circ_0^2 \phi) = \phi \circ \mu_{Q_1}$ and $\mu_{Q_1/\mathcal{E}_1} \circ (\pi_{Q_1/\mathcal{E}_1} \circ_0^2 \pi_{Q_1/\mathcal{E}_1}) = \pi_{Q_1/\mathcal{E}_1} \circ \mu_{Q_1}$, we get:

$$\begin{aligned} \mu_{Q_2} \circ (\hat{\phi} \circ_0^2 \hat{\phi}) \circ (\pi_{Q_1/\mathcal{E}_1} \circ_0^2 \pi_{Q_1/\mathcal{E}_1}) &= \mu_{Q_2} \circ (\phi \circ_0^2 \phi) = \phi \circ \mu_{Q_1} = \hat{\phi} \circ \pi_{Q_1/\mathcal{E}_1} \circ \mu_{Q_1} \\ &= \hat{\phi} \circ \mu_{Q_1/\mathcal{E}_1} \circ (\pi_{Q_1/\mathcal{E}_1} \circ_0^2 \pi_{Q_1/\mathcal{E}_1}). \end{aligned}$$

Since $(\pi_{Q_1/\mathcal{E}_1} \circ_0^2 \pi_{Q_1/\mathcal{E}_1})$ is an epimorphism, we finally have $\mu_{Q_2} \circ (\hat{\phi} \circ_0^2 \hat{\phi}) = \hat{\phi} \circ \mu_{Q_1/\mathcal{E}_1}$ and hence $\hat{\phi}$ is a morphism of (contracted) \hat{T}^* -operadic magmas. \square

Theorem 3.16. *A free contracted \hat{T}^* -operad over a \hat{T}^* -collection always exists.*

Proof. We proceed with a direct iterative construction followed by a quotient.

- a. starting from a globular \hat{T}^* -collection $Q \xrightarrow{\pi_Q} \hat{T}^*(\bullet)$ a new globular \hat{T}^* -collection $\mathfrak{M}(Q) \xrightarrow{\pi_{\mathfrak{M}(Q)}} \hat{T}^*(\bullet)$ is inductively constructed together with a morphism $\xi_Q : Q \rightarrow \mathfrak{M}(Q)$ of globular \hat{T}^* -collections;
- b. we show that the \hat{T}^* -collection $\mathfrak{M}(Q) \xrightarrow{\pi_{\mathfrak{M}(Q)}} \hat{T}^*(\bullet)$ can be equipped with a contraction $\kappa_{\mathfrak{M}(Q)} : \text{Par}(\pi_{\mathfrak{M}(Q)}) \rightarrow \mathfrak{M}(Q)$, a unit $\eta_{\mathfrak{M}(Q)} : \bullet \rightarrow \mathfrak{M}(Q)$ and a multiplication $\mu_{\mathfrak{M}(Q)} : \mathfrak{M}(Q) \circ_0^1 \mathfrak{M}(Q) = \mathfrak{M}(Q) \times_{\hat{T}^*(\bullet)} \hat{T}^*(\mathfrak{M}(Q)) \rightarrow \mathfrak{M}(Q)$;
- c. it is shown that $Q \xrightarrow{\xi_Q} \mathfrak{M}(Q)$ is a free contracted \hat{T}^* -operadic magma over the \hat{T}^* -collection $Q \xrightarrow{\pi_Q} \hat{T}^*(\bullet)$;
- d. we establish the existence of the smallest congruence \mathcal{E}_χ generated by operadic associativity and unitality axioms on the contracted operadic \hat{T}^* -magma $\mathfrak{M}(Q)$, so that the quotient $\mathfrak{M}(Q) \xrightarrow{\pi_{\mathcal{E}_\chi}} \mathfrak{P}(Q) := \mathfrak{M}(Q)/\mathcal{E}_\chi$ becomes a morphism of contracted operadic \hat{T}^* -magmas onto a contracted \hat{T}^* -operad $\mathfrak{P}(Q) \xrightarrow{\pi_{\mathfrak{P}(Q)}} \hat{T}^*(\bullet)$ with operadic multiplication $\mu_{\mathfrak{P}(Q)} : \mathfrak{P}(Q) \circ_0^1 \mathfrak{P}(Q) \rightarrow \mathfrak{P}(Q)$ operadic unit $\eta_{\mathfrak{P}(Q)} : \bullet \rightarrow \mathfrak{P}(Q)$ and contraction $\kappa_{\mathfrak{P}(Q)} : \text{Par}(\pi_{\mathfrak{P}(Q)}) \rightarrow \mathfrak{P}(Q)$, as explained in proposition 3.15;
- e. the universal factorization property for free contracted \hat{T}^* -operads is proved for $Q \xrightarrow{\zeta_Q := \pi_{\mathcal{E}_\chi} \circ \xi_Q} \mathfrak{P}(Q)$.

a.

We start by explicitly providing $\mathfrak{M}(Q)^j \xrightarrow{\pi_{\mathfrak{M}(Q)}^j} \hat{T}^*(\bullet)^j$, for $j = 0, 1$.

Define $\mathfrak{M}(Q)^0 := Q^0 \uplus Q_\eta^0$, disjoint union of Q^0 and a singleton $Q_\eta^0 := \{\bullet_\eta^0\}$, and $\pi_{\mathfrak{M}(Q)}^0 : \mathfrak{M}(Q)^0 \rightarrow \hat{T}^*(\bullet)^0 = \{\bullet^0\}$ as the terminal map, necessarily coinciding with the terminal map π_Q^0 on Q^0 and given by $\bullet_\eta^0 \mapsto \bullet^0$ on the singleton. Notice that, at this level-0 stage, no contraction-cells or operadic multiplication-cells are added, but only a “free operadic-unit 0-cell” \bullet_η^0 .

Passing to the level-1, we define $\mathfrak{M}(Q)^1[0] := Q^1 \uplus Q_\eta^1 \uplus Q_\kappa^1$, where $Q_\eta^1 := \{\bullet_\eta^1\}$ is again a singleton (corresponding to a free operadic-unit 1-cell); $Q_\kappa^1 := \text{Par}(\pi_{\mathfrak{M}(Q)}^0)$ consists of a copy of the set of free contraction 1-cells for the parallel 0-cells induced by the projection $\pi_{\mathfrak{M}(Q)}^0$ at the level-0; we set $\pi_{\mathfrak{M}(Q)}^1[0] : \mathfrak{M}(Q)^1[0] \rightarrow \hat{T}^*(\bullet)^1$ coinciding with π_Q^1 on Q^1 , as a terminal map $\bullet_\eta^1 \mapsto \bullet^1 \in \hat{T}^*(\bullet)^1$ on Q_η^1 , and as the map $(x^+, y, x^-) \mapsto y$ on all $(x^+, y, x^-) \in Q_\kappa^1$; furthermore we introduce new source/target maps $s_{\mathfrak{M}(Q)}^0[0], t_{\mathfrak{M}(Q)}^0[0] : \mathfrak{M}(Q)^1[0] \rightarrow \mathfrak{M}(Q)^0$ coinciding with the original source/target maps on Q^1 ; as $\bullet_\eta^1 \mapsto \bullet_\eta^0$ on Q_η^1 ; and as $s_{\mathfrak{M}(Q)}^0[0] : (x^+, y, x^-) \mapsto x^-$, respectively $t_{\mathfrak{M}(Q)}^0[0] : (x^+, y, x^-) \mapsto x^+$ on Q_κ^1 .

We introduce free operadic multiplication 1-cells by $\mathfrak{M}(Q)^1 := \uplus_{k=0}^{+\infty} \mathfrak{M}(Q)^1[k]$, where we inductively have

$$\mathfrak{M}(Q)^1[k+1] := \{(x, \mu, y) \mid (x, y) \in \mathfrak{M}(Q)^1[k_1] \times_{\hat{T}^*(\bullet)} \hat{T}^*(\mathfrak{M}(Q)^1[k_2]), k_1 + k_2 = k\};$$

for all $(x, y) \in \mathfrak{M}(Q)^1[k_1] \times_{\hat{T}^*(\bullet)} \hat{T}^*(\mathfrak{M}(Q)^1[k_2])$, we recursively define the projection, target and source maps:

$$\begin{aligned} \pi_{\mathfrak{M}(Q)}^1[k+1] : (x, \mu, y) &\mapsto \pi_{\mathfrak{M}(Q)}^1[k_1](x) = \mu_\bullet^* \left(\hat{T}^*(\pi_{\mathfrak{M}(Q)}^1[k_2])(y) \right), \\ t_{\mathfrak{M}(Q)}^0[k+1] : (x, \mu, y) &\mapsto (t_{\mathfrak{M}(Q)}^0[k_1])(x), \\ s_{\mathfrak{M}(Q)}^0[k+1] : (x, \mu, y) &\mapsto \mu_{\mathfrak{M}(Q)}^* \left(\hat{T}^*(s^0[k_2])(y) \right), \end{aligned}$$

where $\mu_{\mathfrak{M}(Q)}^*$ denotes the operadic multiplication that is inductively specified in the following subsection b.²⁵

Assuming now inductively the existence of $\mathfrak{M}(Q)^n \xrightarrow{\pi_{\mathfrak{M}(Q)}^n} \hat{T}^*(\bullet)^n$, we go to construct $\mathfrak{M}(Q)^{n+1} \xrightarrow{\pi_{\mathfrak{M}(Q)}^{n+1}} \hat{T}^*(\bullet)^{n+1}$: starting from $\mathfrak{M}(Q)^{n+1}[0] := Q^{n+1} \uplus Q_\eta^{n+1} \uplus Q_\kappa^{n+1}$, where $Q_\eta^{n+1} := \{\bullet_\eta^{n+1}\}$ is a singleton free operadic-unit $(n+1)$ -cell and $Q_\kappa^{n+1} := \text{Par}(\pi_{\mathfrak{M}(Q)}^n)$ is a family of free contraction $(n+1)$ -cells induced by the already defined projection map $\pi_{\mathfrak{M}(Q)}^n : \mathfrak{M}(Q)^n \rightarrow \hat{T}^*(\bullet)$, then we proceed to introduce free operadic-multiplication $(n+1)$ -cells by the recursive nesting

$$\mathfrak{M}(Q)^{n+1}[k+1] := \{(x, \mu, y) \mid (x, y) \in \mathfrak{M}(Q)^n[k_1] \times_{\hat{T}^*(\bullet)} \hat{T}^*(\mathfrak{M}(Q)^n[k_2]), k_1 + k_2 = k\}$$

and we get $\mathfrak{M}(Q)^{n+1} := \uplus_{k=0}^{+\infty} \mathfrak{M}(Q)^{n+1}[k]$.

The projection map $\pi_{\mathfrak{M}(Q)}^{n+1}$ is separately defined on each set of the disjoint union: it coincides with π_Q^{n+1} on $Q^{n+1} \subset \mathfrak{M}(Q)^{n+1}[0]$; it is $\bullet_\eta^{n+1} \mapsto \bullet^{n+1}$ on Q_η^{n+1} ; it is $(x^+, y, x^-) \mapsto y$ on Q_κ^{n+1} ; and it is recursively given by

$$\pi_{\mathfrak{M}(Q)}^{n+1} : (x, \mu, y) \mapsto \pi_{\mathfrak{M}(Q)}^n[k_1](x) = \mu_{\hat{T}^*} \left(\hat{T}^*(\pi_{\mathfrak{M}(Q)}^n[k_2])(y) \right)$$

²⁵Notice that, since $\mathfrak{M}(Q)$ is inductively defined and the functor \hat{T}^* preserves the inductive grading, the definitions of the source $s_{\mathfrak{M}(Q)}$ and of the operadic multiplication $\mu_{\mathfrak{M}(Q)}$ are both perfectly sound.

on the elements $(x, \mu, y) \in \mathfrak{M}(Q)^{n+1}[k+1]$. Finally we obtain a globular \hat{T}^* -collection with target/source maps given, for all $(x, y) \in \mathfrak{M}(Q)^n[k_1] \times_{\hat{T}^*(\bullet)} \hat{T}^*(\mathfrak{M}(Q)^n[k_2])$, by

$$\begin{aligned} t_{\mathfrak{M}(Q)}^{n+1}[k+1] : (x, \mu, y) &\mapsto (t_{\mathfrak{M}(Q)}^{n+1}[k_1])(x), \\ s_{\mathfrak{M}(Q)}^{n+1}[k+1](x, \mu, y) &\mapsto \mu_{\mathfrak{M}(Q)}^* \left((\hat{T}^*(s_{\mathfrak{M}(Q)}^{n+1}[k_2]))(y) \right), \end{aligned}$$

where again $\mu_{\mathfrak{M}(Q)}$ denotes the operadic multiplication described below, in section b.

The inclusions $\xi_Q^n : Q^n \rightarrow Q^n \uplus Q_\eta^n \uplus Q_\kappa^n = \mathfrak{M}(Q)^n[0] \subset \mathfrak{M}(Q)^n = \uplus_{k=0}^{+\infty} \mathfrak{M}(Q)^n[k]$, for all $n \in \mathbb{N}$ define level-by-level the map $\xi_Q : Q \rightarrow \mathfrak{M}(Q)$ that is already a morphism of globular \hat{T}^* -collections.

b.

We construct level-by-level the several structural maps involved in the definition of contracted \hat{T}^* -operadic magma: the contraction $\kappa_{\mathfrak{M}(Q)}$, the unit $\eta_{\mathfrak{M}(Q)}$, the multiplication $\mu_{\mathfrak{M}(Q)}$.

The operadic unit $\eta_{\mathfrak{M}(Q)} : \bullet \rightarrow \mathfrak{M}(Q)$ is defined as $\eta_{\mathfrak{M}(Q)}^n : \bullet^n \mapsto \bullet_\eta^n$, for all $n \in \mathbb{N}$.

The operadic multiplication $\mu_{\mathfrak{M}(Q)} : \mathfrak{M}(Q) \times_{\hat{T}^*(\bullet)} \hat{T}^*(\mathfrak{M}(Q)) \rightarrow \mathfrak{M}(Q)$ is given, at each level $n \in \mathbb{N}$, for all $k, k_1, k_2 \in \mathbb{N}$ with $k_1 + k_2 = k$, by the maps

$$\mu_{\mathfrak{M}(Q)}^n[k+1] : \mathfrak{M}(Q)^n[k_1] \times_{\hat{T}^*(\bullet)} \hat{T}^*(\mathfrak{M}(Q)^n[k_2]) \rightarrow \mathfrak{M}(Q)^n[k+1]$$

given as $\mu_{\mathfrak{M}(Q)}^n[k+1] : (x, y) \mapsto (x, \mu, y)$.

The contraction $\kappa_{\mathfrak{M}(Q)} : \text{Par}(\pi_{\mathfrak{M}(Q)}) \rightarrow \mathfrak{M}(Q)$ is provided by the maps

$$\kappa_{\mathfrak{M}(Q)}^n : \text{Par}(\pi_{\mathfrak{M}(Q)}^n) \rightarrow \mathfrak{M}(Q)^{n+1}$$

defined, for all $n \in \mathbb{N}$, as inclusions $\kappa_{\mathfrak{M}(Q)}^n : Q_\kappa^n \rightarrow \mathfrak{M}(Q)^{n+1}$.

c.

Here we deal with the universal factorization property of $Q \xrightarrow{\xi_Q} \mathfrak{M}(Q)$: given another contracted \hat{T}^* -operadic magma $\hat{M} \xrightarrow{\hat{\pi}} \hat{T}^*(\bullet)$ with contraction $\hat{\kappa} : \text{Par}(\hat{\pi}) \rightarrow \hat{M}$, operadic unit $\hat{\eta} : \bullet \rightarrow \hat{M}$ and operadic multiplication $\hat{\mu} : \hat{M} \times_{\hat{T}^*(\bullet)} \hat{T}^*(\hat{M}) \rightarrow \hat{M}$, we show the existence of a unique morphism $\hat{\phi} : \mathfrak{M}(Q) \rightarrow \hat{M}$ of contracted \hat{T}^* -operadic magmas such that $\phi = \hat{\phi} \circ \xi_Q$.

Since, by construction, the inclusion $Q \xrightarrow{\xi_Q} \mathfrak{M}(Q)$, maps every element $x \in Q^n$ to the same element $x \in \mathfrak{M}(Q)^n$, for all $n \in \mathbb{N}$, we must necessarily have that $\hat{\phi}(x) := x$, for all $x \in Q \subset \mathfrak{M}(Q)$. Since $\hat{\phi}$ should preserve the unit, $\hat{\phi} \circ \eta_{\mathfrak{M}(Q)} = \hat{\eta}$, the explicit construction of $\eta_{\mathfrak{M}(Q)}$, necessarily entails $\hat{\phi} : \eta_{\mathfrak{M}(Q)}(\bullet) \mapsto \hat{\eta}(\bullet)$, for all the elements $\eta_{\mathfrak{M}(Q)}(\bullet) \subset \mathfrak{M}(Q)$.

Similarly, since $\hat{\phi}$ should be contraction preserving, $\hat{\phi} \circ \kappa_{\mathfrak{M}(Q)} = \hat{\kappa} \circ (\hat{\phi}, \hat{\phi})$, the only possible choice for the restriction of $\hat{\phi}$ on $\kappa_{\mathfrak{M}(Q)}(\text{Par}(\pi_{\mathfrak{M}(Q)})) \subset \mathfrak{M}(Q)$ is given by: $\hat{\phi}(x^+, y, x^-) \mapsto \hat{\kappa}(\hat{\phi}(x^+), y, \hat{\phi}(x^-))$.²⁶

Since $\hat{\phi}$ should preserve multiplications $\hat{\phi}^n \circ (\mu_{\mathfrak{M}(Q)})^n = \hat{\mu}^n \circ (\hat{\phi}^n, (\hat{T}^*(\hat{\phi}))^n)$, for all $n \in \mathbb{N}$, the map $\hat{\phi}^n$ should be uniquely defined as $(x, \mu, y) \mapsto \hat{\mu}(\hat{\phi}^n(x), \hat{T}^*(\hat{\phi})^n(y))$ on the elements in $\mu_{\mathfrak{M}(Q)}(\mathfrak{M}(Q) \times_{\hat{T}^*(\bullet)} \hat{T}^*(\mathfrak{M}(Q))^n) \subset \mathfrak{M}(Q)^n$.²⁷ The already uniquely constructed $\hat{\phi} : \mathfrak{M}(Q) \rightarrow \hat{M}$ is a morphism of globular contracted \hat{T}^* -operadic magmas that also satisfies $\phi = \hat{\phi} \circ \xi_Q$.

d.

The previously constructed free contracted \hat{T}^* -operadic magma $\mathfrak{M}(Q)$ is not yet a free contracted operad because its free unit $\eta_{\mathfrak{M}(Q)} : \bullet \rightarrow \mathfrak{M}(Q)$ and free multiplication $\mu_{\mathfrak{M}(Q)} : \mathfrak{M}(Q) \circ_0^1 \mathfrak{M}(Q) \rightarrow \mathfrak{M}(Q)$ fail to satisfy the unitality and associativity axioms for a monad in the bicategory $\mathcal{Q}_{\hat{T}^*}$ as specified by the commuting diagrams in definition 2.6, in detail, denoting the associators and left/right unitors morphisms by $(\mathfrak{M}(Q) \circ_0^1 \mathfrak{M}(Q)) \circ_0^1 \mathfrak{M}(Q) \xrightarrow{\alpha_{\mathfrak{M}(Q)}} \mathfrak{M}(Q) \circ_0^1 (\mathfrak{M}(Q) \circ_0^1 \mathfrak{M}(Q))$ and $\iota^1(\bullet) \circ_0^1 \mathfrak{M}(Q) \xrightarrow{\lambda_{\mathfrak{M}(Q)}} \mathfrak{M}(Q) \xleftarrow{\rho_{\mathfrak{M}(Q)}} \mathfrak{M}(Q) \circ_0^1 \iota^1(\bullet)$, we need to get identified all the pairs of terms in $\mathfrak{M}(Q)$ contained in the following family $\mathcal{X} \subset \mathfrak{M}(Q) \times \mathfrak{M}(Q)$:

$$\begin{aligned} \mathcal{X} := & \left\{ \left(\lambda_{\mathfrak{M}(Q)}(x_1), \mu_{\mathfrak{M}(Q)} \left(\eta_{\mathfrak{M}(Q)} \circ_0^2 \iota_{\mathfrak{M}(Q)}^1(x_1) \right) \right) \mid x_1 \in \iota^1(\bullet) \circ_0^1 \mathfrak{M}(Q) \right\} \cup \\ & \left\{ \left(\rho_{\mathfrak{M}(Q)}(x_2), \mu_{\mathfrak{M}(Q)} \left(\iota_{\mathfrak{M}(Q)}^1 \circ_0^2 \eta_{\mathfrak{M}(Q)}(x_2) \right) \right) \mid x_2 \in \mathfrak{M}(Q) \circ_0^1 \iota^1(\bullet) \right\} \cup \\ & \left\{ \left(\mu_{\mathfrak{M}(Q)} \left(\iota_{\mathfrak{M}(Q)}^1 \circ_0^2 \mu_{\mathfrak{M}(Q)} \right) \circ \alpha_{\mathfrak{M}(Q)}(x_3), \mu_{\mathfrak{M}(Q)} \left(\mu_{\mathfrak{M}(Q)} \circ_0^2 \iota_{\mathfrak{M}(Q)}^1(x_3) \right) \right) \mid \right. \\ & \left. x_3 \in (\mathfrak{M}(Q) \circ_0^1 \mathfrak{M}(Q)) \circ_0^1 \mathfrak{M}(Q) \right\}. \end{aligned} \quad (3.9)$$

To solve this problem, we define $\mathcal{E}_{\mathcal{X}}$ as the *smallest congruence of contracted \hat{T}^* -operadic magma* in $\mathfrak{M}(Q)$ containing all the pairs of terms in \mathcal{X} . From remark 3.11 $\hat{T}^*(\bullet) \xrightarrow{\pi_{\hat{T}^*(\bullet)}} \hat{T}^*(\bullet)$ is a contracted \hat{T}^* -operad and, since $\mathfrak{M}(Q) \xrightarrow{\pi_{\mathfrak{M}(Q)}} \hat{T}^*(\bullet)$ is morphism of contracted \hat{T}^* -operadic magmas, by proposition 3.15 we know that the canonical equivalence relation

$$\mathcal{E}_{\pi_{\mathfrak{M}(Q)}} := \mathfrak{M}(Q) \times_{\pi_{\mathfrak{M}(Q)}} \mathfrak{M}(Q) = \{(x, y) \in \mathfrak{M}(Q) \times \mathfrak{M}(Q) \mid \pi_{\mathfrak{M}(Q)}(x) = \pi_{\mathfrak{M}(Q)}(y)\}$$

is itself a congruence of contracted \hat{T}^* -operadic magmas with $\mathcal{X} \subset \mathcal{E}_{\pi_{\mathfrak{M}(Q)}}$ and hence $\mathcal{E}_{\mathcal{X}}$ can be taken as the intersection of the (non-empty) family of all such congruences containing \mathcal{X} in $\mathfrak{M}(Q)$.

²⁶Notice that, if $(x^+, y, x^-) \in \text{Par}(\pi_{\mathfrak{M}(Q)})^n$, we have $x^+, x^- \in \mathfrak{M}(Q)^{n-1}$ and, since the construction of $(\mathfrak{M}(Q), \pi_{\mathfrak{M}(Q)}, \kappa_{\mathfrak{M}(Q)})$, is produced inductively, for all $n \in \mathbb{N}$, the definition of $\hat{\phi}$ on the elements $\kappa_{\mathfrak{M}(Q)}(\text{Par}(\pi_{\mathfrak{M}(Q)}))^n \subset \mathfrak{M}(Q)^n$ requires only the knowledge of the already available map $\hat{\phi}^{n-1}$.

²⁷Notice again that, because of the inductive definition of $\mathfrak{M}(Q)$, the elements in $\mathfrak{M}(Q) \times_{\hat{T}^*(\bullet)} \hat{T}^*(\mathfrak{M}(Q))^n$ only require the knowledge of already defined $\hat{\phi}^k$, for all $k \leq n$ on .

By proposition 3.15, the quotient map $\mathfrak{M}(Q) \xrightarrow{\pi_{\mathcal{E}_X}} \mathfrak{F}(Q) := \mathfrak{M}(Q)/\mathcal{E}_X$ is a morphism of contracted \hat{T}^* -operadic magmas and, since $X \subset \mathcal{E}_X$, furthermore we have that the quotient $\mathfrak{F}(Q) := \mathfrak{M}(Q)/\mathcal{E}_X \xrightarrow{\pi_{\mathfrak{F}(Q)}} \hat{T}^*(\bullet)$, where $\pi_{\mathfrak{F}(Q)}$ is the only morphism of contracted \hat{T}^* -operadic magmas such that $\pi_{\mathfrak{M}(Q)} = \pi_{\mathfrak{F}(Q)} \circ \pi_{\mathcal{E}_X}$, is not only a contracted \hat{T}^* -operadic magma, but it is already a contracted \hat{T}^* -operad.

The inclusion $Q \xrightarrow{\zeta_Q} \mathfrak{F}(Q)$ given by $\zeta_Q := \pi_{\mathcal{E}_X} \circ \xi_Q$ is, by composition, a morphism of globular ω -sets.

e.

Suppose that $Q \xrightarrow{\phi} \hat{P}$ is a morphism of \hat{T}^* -collections into another contracted \hat{T}^* -operad $\hat{P} \xrightarrow{\hat{\pi}_P} \hat{T}^*(\bullet)$ with contraction $\hat{\kappa}_P : \text{Par}(\hat{\pi}_P) \rightarrow \hat{P}$, operadic unit $\hat{\eta}_P : \bullet \rightarrow \hat{P}$ and operadic multiplication $\hat{\mu}_P : \hat{P} \circ_0^1 \hat{P} \rightarrow \hat{P}$.

Since (forgetting the associativity and unitality axioms) every contracted \hat{T}^* -operad is a contracted \hat{T}^* -operadic magma, by the previous point c. above, there exists a unique morphism $\mathfrak{M}(Q) \xrightarrow{\tilde{\phi}} \hat{P}$ of contracted \hat{T}^* -operadic magmas, defined on the free contracted \hat{T}^* -operadic magma $\mathfrak{M}(Q)$, such that $\phi = \tilde{\phi} \circ \xi_Q$.

The equivalence relation $\mathcal{E}_{\tilde{\phi}} := \mathfrak{M}(Q) \times_{\tilde{\phi}} \mathfrak{M}(Q) = \{(x, y) \in \mathfrak{M}(Q) \times \mathfrak{M}(Q) \mid \tilde{\phi}(x) = \tilde{\phi}(y)\}$ induced by the morphism $\mathfrak{M}(Q) \xrightarrow{\tilde{\phi}} \hat{P}$ into the contracted \hat{T}^* -operad \hat{P} , is a congruence of contracted \hat{T}^* -operadic magma that contains all the terms X generating \mathcal{E}_X and hence, by the minimality of \mathcal{E}_X , we have $\mathcal{E}_X \subset \mathcal{E}_{\tilde{\phi}}$. It follows, by proposition 3.15, that there exists a unique quotient morphism $\mathfrak{F}(Q) \xrightarrow{\hat{\phi}} \hat{P}$ of contracted \hat{T}^* -operads, given by $\hat{\phi}(\pi_{\mathcal{E}_X}(x)) := \tilde{\phi}(x)$, for all $x \in \mathfrak{M}(Q)$. The universal factorization property for free \hat{T}^* -operads over Q is satisfied since: $\hat{\phi} \circ \zeta_Q = \hat{\phi} \circ \pi_{\mathcal{E}_X} \circ \xi_Q = \tilde{\phi} \circ \xi_Q = \phi$. \square

Remark 3.17. The proof of theorem 3.16 could be obtained mimicking Leinster's techniques in section 2.3.3. Specifically, applying lemma 2.28 after showing that: a) the category $\mathcal{Q}_{\hat{T}^*}$ is locally finitely presentable; b) \hat{T}^* is a finitary Cartesian monad (and hence $\mathfrak{U}_{\hat{T}^*}$ is monadic) c) $\mathfrak{U}_{\hat{T}^*}$ is finitary d) $\mathfrak{U}_{\hat{T}^*}$ is monadic and finitary. \dashv

Our main definition in the \hat{T}^* case, is in perfect analogy with the Leinster's definition 2.29:

Definition 3.18. A *weak involutive globular ω -category* is an algebra for an initial object L^* in $\mathcal{O}_{\bullet}^{\hat{T}^*, \kappa}$.

Remark 3.19. It is perfectly possible to utilize the category $\mathcal{O}_{\bullet}^{\hat{T}^*, \kappa}$ introduced in remark 3.10 instead of $\mathcal{O}_{\bullet}^{\hat{T}^*, \kappa}$ in order to define a slightly more restrictive notion of involutive weak globular ω -category as an algebra for the initial object in $\mathcal{O}_{\bullet}^{\hat{T}^*, \kappa}$.

The existence of such initial object can be obtained with techniques perfectly similar to those utilized in theorems 3.12 and in the proof of theorem 3.16, just adding to

the family X in equation (3.9) all the pairs of terms of $\mathfrak{M}(Q)$ required for the validity of the additional axioms imposed by the commuting diagrams (3.3) and quotienting by the smallest congruence of \hat{T}^* -operadic contraction containing X (such minimal congruence always exists because $\hat{T}^*(\bullet)$ is a terminal object in $\mathcal{O}\mathcal{H}_{\bullet}^{\hat{T}^*}$). \dashv

3.3 Examples

The strategy used to provide examples of weak involutive globular ω -categories is perfectly parallel to the one described in section 2.3.4 and consists of producing: a contracted \hat{T}^* -operad P^* and an algebra X^* over it.

We just mention here some immediate available examples of involutive weak globular ω -categories.

- **strict involutive globular ω -categories:** again, these coincide with algebras for the monad given by the terminal contracted- \hat{T}^* -operad $\hat{T}^*(\bullet) \in \mathcal{O}\mathcal{H}_{\bullet}^{\hat{T}^*}$ described in remark 3.11.
- **globular ω -spans:** following the same notation introduced in [Bertozzini Conti Lewkeeratiyutkul Suthichitranont 2020, example 4.7] we define an ω -span as a sequence $(A^n \xleftarrow{r^n} Q^{n+1} \xrightarrow{s^n} B^n)_{n \in \mathbb{N}}$ of 1-spans, where $Q^n := A^n \cup B^n$; a globular ω -span is an ω -span $Q := (Q^{n+1} \rightrightarrows Q^n)$ that is a globular ω -set (in this way any $x \in Q^n$ determines, with all its sources/targets a unique globular n -cell). Introducing level-by-level the equivalence relation that identifies $(n+1)$ -cells having the same source/target sets, we obtain a quotient ω -globular set X . A family of globular ω -spans is hence determined by the quotient morphism $Q \xrightarrow{X} X$. Considering such family $Q \xrightarrow{X} X \rightarrow \bullet$ as “generators”, we apply the free involutive magma functor (as constructed in point a. of the proof of proposition 3.2) to get $\mathfrak{M}(Q) \xrightarrow{\mathfrak{M}(X)} \mathfrak{M}(X) \xrightarrow{\mathfrak{M}(!)} \mathfrak{M}(\bullet)$, applying the forgetful functor to the category of globular ω -sets (that we omit to indicate) and using the quotient projection (constructed in point b. of proposition 3.2) onto $\hat{T}^*(Q) \xrightarrow{\hat{T}^*(X)} \hat{T}^*(X) \xrightarrow{\hat{T}^*(!)} \hat{T}^*(\bullet)$, we obtain a morphism of \hat{T}^* -collections as in the first diagram below:

$$\begin{array}{ccc}
 \mathfrak{M}(Q) & \xrightarrow{\mathfrak{M}(X)} & \mathfrak{M}(X) \\
 & \searrow & \swarrow \\
 & \hat{T}^*(\bullet) &
 \end{array}
 \quad \mapsto \quad
 \begin{array}{ccc}
 \mathfrak{F}(Q) & \xrightarrow{\mathfrak{F}(X)} & \mathfrak{F}(X)
 \end{array}$$

The *involutive weak globular ω -category of globular ω -spans generated by $Q \xrightarrow{X} X$* is given by the morphism of free globular contracted \hat{T}^* -operads to the right of the diagram above (considered as algebras over themselves). Keeping track of the projection onto X and $\mathfrak{F}(X)$ is necessary to “coarse grain” recovering the original spans and operations between them.

- **homotopy ω -groupoids** $\Pi_\omega(X)$ of a topological space X : here, since the involutions coincide with the weak inverse homotopies, one just show that the contracted \hat{T} -operad utilized in [Leinster 2004, example 9.2.7] is actually a contracted \hat{T}^* -operad.

4 Outlook

The construction of algebraic involutive versions of weak (globular) ω -categories (either in Penon’s or in Leinster’s approaches), as done in our previous work [Bejrakarbum Bertozzini 2017] and in the present paper, is only the very first step in the direction of a full operator algebraic categorical environment suitable for the needs of *categorical non-commutative geometry* [Bertozzini Conti Lewkeeratiyutkul 2008], [Bertozzini Conti Lewkeeratiyutkul 2012].

Involutions, in the case of (weak) cubical ω -categories, are currently under investigation,²⁸ following recent work by [Kachour 2022], including the study of conditions (see [Al-Agl Brown, Steiner 2002]) assuring the equivalence between cubical and globular *involutive* settings, extending previous still unpublished results already achieved in the case of involutive 2-categories / double categories [Bertozzini Conti Dawe Martins 2014].

Immediate further developments of the present work will concentrate on possible algebraic definitions and examples of involutive weak ω -algebroids; subsequently the treatment of uniform structures related to completeness and norms necessary for the formulation of weak ω - C^* -algebroids will have to be addressed, generalizing (and possibly modifying) the strict n - C^* -categorical notions tentatively put forward in [Bertozzini Conti Lewkeeratiyutkul Suthichitranont 2020, section 5].

In the present paper, for simplicity, we have only considered monads and operads that do not possess involutive symmetries, but it is already evident that certain “covariant” involutions could have been introduced at the level of the operads in $\hat{\mathcal{O}}^*$. Certain involutive monads and operads (see [Yau 2020, chapter 4] and references therein) can be used for this purpose. A full treatment of *involutive bicategories* and the discussion of *covariant vs contravariant involutive monads and operads in a bicategory* will be separately addressed in a companion paper²⁹ making direct use of hybrid-categories as put forward in [Bertozzini Puttirungroj 2014].

In this work we have not considered any relaxing of the usual axioms for globular higher categories, in particular we did not formulate a definition of weak involutive globular ω -categories with *non-commutative exchange property*, as already proposed in [Bertozzini Conti Lewkeeratiyutkul Suthichitranont 2020, section 3.3] for strict

²⁸Bejrakarbum P, Bertozzini P, Theesoongnern S, *Involutive Weak Cubical/Globular ω -categories* (work in progress).

²⁹Bejrakarboom P, Bertozzini P, Puttirungroj C, *Involutive Monads and Hybrid Categories* (work in progress).

globular n -categories. We suspect that treatments of versions of *non-commutative derived geometries* (homotopies, cobordisms, holonomies) will require some axiomatic modification in such direction.

Notes and Acknowledgments: P. Bejrakarbum is supported by the “Research Professional Development Project” under the “Science Achievement Scholarship of Thailand” (SAST).

P. Bertozzini thanks Starbucks Coffee (Langsuan, Jasmine City, Gaysorn Plaza and Emquartier Sky Garden) where he spent most of the time dedicated to this research project; he thanks Fiorentino Conte of “The Melting Clock” for the great hospitality during many crucial on-line dinner-time meetings.

References

- [Al-Agl Brown, Steiner 2002]
Al-Agl F A, Brown R, Steiner R (2002) **Multiple Categories: the Equivalence of a Globular and a Cubical Approach** *Adv Math* 170(1):71-118
- [Baez 1997]
Baez J C (1997) **An Introduction to n -categories** *Category Theory and Computer Science (Santa Margherita Ligure 1997)* Moggi E, Rosolini G (eds), *Lecture Notes in Computer Science* 1290:1-33 Springer arXiv:q-alg/9705009
- [Baez Dolan 1995]
Baez J, Dolan J (1995) **Higher-dimensional Algebra and Topological Quantum Field Theory** *J Math Phys* 36:6073-6105 arXiv:q-alg/9503002
- [Bénabou 1967]
Bénabou J (1967) **Introduction to Bicategories** *Reports of the Midwest Category Seminar* Bénabou J et al (eds) *Lecture Notes in Mathematics* 47 Springer
- [Batatin 1998]
Batatin M (1998) **Monoidal Globular Categories as a Natural Environment for the Theory of Weak n -categories** *Advances in Mathematics* 136(1):39-103
- [Batatin 2022]
Batatin M (2002) **On the Penon Method of Weakening Algebraic Structures** *Journal of Pure and Applied Algebra* 172(1):1-23
- [Bejrakarbum 2016]
Bejrakarbum P (2016) **Involutive Weak Globular Higher Categories** MSc Thesis Thammasat University
- [Bejrakarbum 2023]
Bejrakarbum P (2023) **Involutive Weak Globular Higher Categories: Leinster’s Approach** PhD Thesis Thammasat University
- [Bejrakarbum Bertozzini 2017]
Bejrakarbum P, Bertozzini P (2017) **Involutive Weak Globular Higher Categories** arXiv:1709.09336 [math.CT]

- [Bertozzini Conti Dawe Martins 2014]
 Bertozzini P, Conti R, Dawe Martins R (2014) **Involutive Double Categories** (manuscript)
- [Bertozzini Conti Lewkeeratiyutkul 2008]
 Bertozzini P, Conti R, Lewkeeratiyutkul W (2008) **Non-commutative Geometry Categories and Quantum Physics** *Contributions in Mathematics and Applications II East-West J Math* special volume 2007:213-259
 arXiv:0801.2826v2
- [Bertozzini Conti Lewkeeratiyutkul 2012]
 Bertozzini P, Conti R, Lewkeeratiyutkul W (2012) **Categorical Non-commutative Geometry** *J Phys: Conf Ser* 346:012003 arXiv: 1409. 1337
- [Bertozzini Conti Lewkeeratiyutkul Suthichitranont 2020]
 Bertozzini P, Conti R, Lewkeeratiyutkul W, Suthichitranont N (2020) **On Strict Higher C*-categories** *Cahiers de Topologie et Géométrie Différentielle Catégoriques* LXI(3):239-348
- [Bertozzini Puttirungroj 2014]
 Bertozzini P, Puttirungroj C (2014) **Hybrid Categories** *Proceedings of AMM2014 (19th Annual Meeting in Mathematics, 20-22 March 2014, A-One Hotel, Pattaya, Thailand* 119-128
- [Borceux 1994]
 Borceux F (1994) **Handbook of Categorical Algebra I-II-III** Cambridge University Press
- [Burroni 1971]
 Burroni A (1971) **T-catégories (Catégories Dans un Triple)** *Cahiers de Topologie et Géométrie Différentielle* 12:215-321
- [Cheng Lauda 2004]
 Cheng E, Lauda A (2004) **Higher-Dimensional Categories: an Illustrated Guide Book** IMA Workshop
- [Chikhladze 2015]
 Chikhladze D (2015) **Lax Formal Theory of Monads, Monoidal Approach to Bicategorical Structures and Generalized Operads** *Theory and Applications of Categories* 30(10), 2015, pp. 332–386. arXiv:1412.4628
- [Eilenberg Mac Lane 1945]
 Eilenberg S, Mac Lane S (1945) **General Theory of Natural Equivalences** *Trans Am Math Soc* 58:231-294
- [Godement 1958]
 Godement R (1958) **Topologie Algébrique et Théorie des Faisceaux** *Actualités Sci Ind Publ Math Univ Strasbourg* 1252 Hermann
- [Hermida 1997]
 Hermida C (1997) **Higher-dimensional Multicategories** (slides)
- [Kachour 2022]
 Kachour C (2022) **Algebraic Models of Cubical Weak ∞ -categories with Connections** *Categ Gen Algebr Struct Appl* 16(1):143-187
 Kachour C (2022) **Algebraic Models of Cubical Weak Higher Structures** *Categ Gen Algebr Struct Appl* 16(1):189-220

[Kelly 1980]

Kelly G M (1980) **A Unified Treatment of Transfinite Constructions for Free Algebras, Free Monoids, Colimits, Associated Sheaves, and So On** *Bulletin of the Australian Mathematical Society* 22:1-83

[Leinster 2001]

Leinster T (2001) **Structures in Higher-dimensional Category Theory**
arXiv:math/0109021

[Leinster 1998]

Leinster T (1998) **Basic Bicategories** arXiv:math/9810017

[Leinster 2004]

Leinster T (2004) **Higher Operads, Higher Categories** Cambridge University Press
arXiv:math/0305049 [math.CT]

[coherence in n-Lab]

n-Lab <https://ncatlab.org/nlab/show/coherence+theorem+for+monoidal+categories>

[Penon 1999]

Penon J (1999) **Approche Polygraphique des ω -Categories Non Strictes** *Cahiers de Topologie et Géométrie Différentielle* 40(1):31-80

[Riehl 2016]

Riehl E (2016) **Category Theory in Context** Dover

[Street 1972]

Street R (1972) **The Formal Theory of Monads** *Journal of Pure and Applied Algebra* 2(2):149-168

[Yau 2020]

Yau D (2020) **Involutive Category Theory** *Lecture Notes in Mathematics* 2279 Springer