

EDGE-ODD GRACEFUL LABELINGS OF SOME PRISMS AND PRISM-LIKE GRAPHS.

Ratinan Boonklurb¹, Sirirat Singhun²
and
Akkaporn Wongpradit³

¹*Department of Math. and Computer Science, Faculty of Science,
Chulalongkorn University, Bangkok 10330, Thailand
e-mail: ratinan.b@chula.ac.th*

²*Department of Mathematics, Faculty of Science,
Ramkhamhaeng University, Bangkok 10240, Thailand
e-mail: sin_sirirat@ru.ac.th*

³*The Demonstration School, Silpakorn University,
Sanamchandra Palace Campus, Nakhon Phathom 73000, Thailand
e-mail: aw_np@hotmail.com*

Abstract

A simple graph G with q edges is called an *edge-odd graceful graph*, EOGG, if there is a bijection f from the edge set of the graph to the set $\{1, 3, 5, \dots, 2q - 1\}$ such that, when each vertex is assigned the sum of all values of the edges incident to it modulo $2q$, the resulting vertex labels are distinct.

In this paper, we define new graphs called a prism of star S_n , $\text{Prism}(S_n)$, a prism-like graph, $\text{Prism}_3(S_n)$, and a prism of wheel graph W_n , $\text{Prism}(W_n)$. We give necessary conditions on n that force these graphs to be EOGG, namely, (i) if $n \geq 3$, then $\text{Prism}(S_n)$ is an EOGG; (ii) if $n \geq 3$ and $n \equiv 2 \pmod{6}$, then $\text{Prism}_3(S_n)$ is an EOGG; (iii) if $n \geq 3$ and $2|n$, then $\text{Prism}(W_n)$ is an EOGG.

Key words: edge-odd graceful labeling, prism, star, wheel.

2010 AMS Mathematics Subject Classification: 05C78.

¹ Corresponding author

1 Introduction

Let G be a simple graph with q edges. In this article, we let $V(G)$ and $E(G)$ denote the vertex set and the edge set of G , respectively. In 1967, Rosa [3] gave a definition of a *graceful labeling* of G which is an injection f from $V(G)$ to the set $\{0, 1, 2, \dots, q\}$ such that each edge xy is assigned the label $|f(x) - f(y)|$, the resulting edge labels are distinct. In 1991, Gnanajothi [2] introduced an *odd-graceful labeling* concept for a graph, that is an injection f from $V(G)$ to the set $\{0, 1, 2, \dots, 2q - 1\}$ such that, when each edge xy is assigned the label $|f(x) - f(y)|$, the resulting edges labels are in the set $\{1, 3, 5, \dots, 2q - 1\}$. In 2009, Solairaju and Chithra [6] reversed the concepts of those two previous vertex labelings by defining an *edge-odd graceful labeling* as the following.

Definition 1.1. Let G be a simple graph an *edge-odd graceful labeling* of G is a bijection f from $E(G)$ to the set $\{1, 3, 5, \dots, 2q - 1\}$ so that the induced mapping f^+ from $V(G)$ to the set $\{0, 1, 2, \dots, 2q - 1\}$ given by $f^+(x) = \sum f(xy) \pmod{2q}$ where the vertex x is adjacent to the vertex y . The edge labels and vertex labels are distinct. A graph that admitted an edge-odd graceful labeling is called an *edge-odd graceful graph* denoted by EOGG.

Solairaju and Chithra [6] showed edge-odd graceful labelings of graphs related to paths. Later, Singhun [4] showed edge-odd graceful labelings of graphs related to cycles, $SF(n, m)$, where n is an odd integer and m is an even integer such that $n \geq 3$ and $n|m$ and a wheel graph W_n , where n is even. In 2013, Boonklurb et. al. [1] showed edge-odd graceful labelings of prism of cycle C_n , where $n \geq 3$ and $\text{Shaft}(n, 1)$, where $n \geq 3$ and $n \equiv 1 \pmod{2}$. Note that $\text{Shaft}(n, 1)$ is a graph consists of 2 copies of wheel graphs joining at the middle. Here, we define new graphs from star, S_n , and wheel, W_n . Thus, for ease of reference, we give definitions of them as follow.

Definition 1.2. Let $n \in \mathbb{N}$. A complete bipartite $K_{1,n}$ is called *star*, denoted by S_n .

In this article, we usually let the first partite with one element be $\{u\}$ and the second partite with n elements be $\{u_1, u_2, u_3, \dots, u_n\}$.

Definition 1.3. A *wheel* graph W_n is a graph with $n + 1$ vertices obtained by connecting a single vertex u to all vertices of a cycle $u_1u_2u_3 \dots u_nu_1$. Then, the vertex set of W_n is the set $\{u, u_1, u_2, u_3, \dots, u_n\}$ and the edge set of W_n is the set $\{uu_i | i \in \{1, 2, 3, \dots, n\}\} \cup \{u_iu_{i+1} | i \in \{1, 2, 3, \dots, n - 1\}\} \cup \{u_1u_n\}$.

In Section 2, a prism of star, $\text{Prism}(S_n)$, is defined, algorithms for constructing edge-odd graceful labelings are given and we can prove that $\text{Prism}(S_n)$ is an EOGG for $n \geq 3$. In Section 3, we extend the idea of prism to a prism-like graph by defining $\text{Prism}_3(S_n)$. After that, we can show that if $n \geq 3$ and $n \equiv 2 \pmod{6}$, then we can be able to label the edge of $\text{Prism}_3(S_n)$ in such a way that $\text{Prism}_3(S_n)$ is an EOGG. In Section 4, we give a definition of a prism of

wheel, $\text{Prism}(W_n)$. If we impose the conditions: $n \geq 3$ and $n \equiv 0 \pmod{2}$, then we can find an algorithm for labeling $\text{Prism}(W_n)$ and can prove that this labeling is an edge-odd graceful labeling for $\text{Prism}(W_n)$. In the last section, we give a conclusion of our results and some discussion on the on-going research.

2 Prism of star

Definition 2.1. For $n \geq 3$. Let S_n be a star and S'_n be a copy of S_n . Define $\text{Prism}(S_n)$, called the *prism* of S_n , by joining u of S_n to the corresponding vertex u' of S'_n and each u_i of S_n to the corresponding vertex u'_i of S'_n for all $i \in \{1, 2, 3, \dots, n\}$. Thus,

$$E(\text{Prism}(S_n)) = E(S_n) \cup E(S'_n) \cup \{u_i u'_i \mid i \in \{1, 2, 3, \dots, n\}\} \cup \{uu'\}.$$




Figure 1: $\text{Prism}(S_4)$

Note that, $\text{Prism}(S_n)$ can be called a *book* graph. Solairaju et.al. [5] gave algorithms to construct an edge-odd graceful labeling for book graphs without rigorous proofs. The following algorithms are different from those given in [5] and later we give rigorous proofs to show that the labelings from the following algorithms are edge-odd graceful labelings for $\text{Prism}(S_n)$.

Algorithm 2.1. If $n = 3$, we label each edge as shown in Figure 2.




Figure 2: Edge-labelings for $\text{Prism}(S_3)$

Algorithm 2.2. Let $n \geq 4$ and $n \equiv 0 \pmod{2}$. Then, $q = |E(\text{Prism}(S_n))| = 3n + 1$. Define $f : E(\text{Prism}(S_n)) \rightarrow \{1, 3, 5, \dots, 6n + 1\}$ by

- i $f(u_i u'_i) = 2i - 1$, for $i \in \{1, 2, 3, \dots, n\}$;
- ii $f(u_i u) = 2n + 4i - 3$, for $i \in \{1, 2, 3, \dots, n\}$;
- iii $f(u'_i u') = 2n + 4i - 1$, for $i \in \{1, 2, 3, \dots, n\}$;
- iv $f(uu') = 6n + 1$.




Figure 3: Edge-labelings for $\text{Prism}(S_6)$

For an odd integer n such that $n > 3$, we separate it into 3 cases as follow.

Algorithm 2.3. Let $n > 3$ and $n \equiv 5 \pmod{6}$. Then, $q = |E(\text{Prism}(S_n))| = 3n + 1$. Define $f : E(\text{Prism}(S_n)) \rightarrow \{1, 3, 5, \dots, 6n + 1\}$ by

- i $f(u_i u'_i) = 2n + 2i - 1$, for $i \in \{1, 2, 3, \dots, n\}$;
- ii $f(u_n u) = 1$;
- iii $f(u_i u) = 2i + 1$, for $i \in \{1, 2, 3, \dots, n - 1\}$;
- iv $f(u'_i u') = 4n + 2i - 1$, for $i \in \{1, 2, 3, \dots, n\}$;
- v $f(uu') = 6n + 1$.





Figure 4: Edge-labelings for $\text{Prism}(S_5)$


Algorithm 2.4. Let $n > 3$ and $n \equiv 1 \pmod{6}$. Then, $q = |E(\text{Prism}(S_n))| = 3n + 1$. Define $f : E(\text{Prism}(S_n)) \rightarrow \{1, 3, 5, \dots, 6n + 1\}$ by

- i $f(u_i u'_i) = 4n + 2i + 1$, for $i \in \{1, 2, 3, \dots, n\}$;
- ii $f(u_i u) = 2i + 1$, for $i \in \{1, 2, 3, \dots, n\}$;
- iii $f(u'_1 u') = 1$;
- iv $f(u'_i u') = 2n + 2i + 1$, for $i \in \{2, 3, 4, \dots, n\}$;
- v $f(uu') = 2n + 3$.


Figure 5: Edge-labelings for $\text{Prism}(S_7)$

Algorithm 2.5. Let $n > 3$ and $n \equiv 3 \pmod{6}$. Then, $q = |E(\text{Prism}(S_n))| = 3n + 1$. Define $f : E(\text{Prism}(S_n)) \rightarrow \{1, 3, 5, \dots, 6n + 1\}$ by

- i $f(u_i u'_i) = 4n + 2i + 1$, for $i \in \{1, 2, 3, \dots, n\}$;
- ii $f(u_i u) = 2i + 1$, for $i \in \{1, 2, 3, \dots, n\}$;
- iii $f(u'_i u') = 2n + 2i + 1$, for $i \in \{1, 2, 3, \dots, n\}$;
- iv $f(uu') = 1$.

Figure 6: Edge-labelings for $\text{Prism}(S_9)$

Lemma 2.1. $\text{Prism}(S_3)$ is an EOGG.

Figure 7: The vertex-labeling for $\text{Prism}(S_3)$ induced by Figure 2

Proof. The vertex labeling of $\text{Prism}(S_3)$ induced by the edge labeling given in Algorithm 2.1 is shown in Figure 7. Therefore, the function f defined in Algorithm 2.1 is an edge-odd graceful labeling and this implies that $\text{Prism}(S_3)$ is an EOGG. $\square$

Lemma 2.2. *If $n \geq 3$ and $n \equiv 0 \pmod{2}$, then $\text{Prism}(S_n)$ is an EOGG.*

Proof. From Algorithm 2.2(i-iv), the edge labels are arranged in the set $\{1, 3, 5, \dots, 2n-1\}$, $\{2n+1, 2n+5, 2n+9, \dots, 6n-3\}$, $\{2n+3, 2n+7, 2n+11, \dots, 6n-1\}$, and $\{6n+1\}$, respectively. Then, f is a bijection from $E(\text{Prism}(S_n))$ to $\{1, 3, 5, \dots, 6n-1\}$.

Next, from Algorithm 2.2, we have

$$f^+(u_i) = (f(u_i u'_i) + f(u_i u)) \pmod{6n+2} = 2n + 6i - 4 \pmod{6n+2} \text{ for } i \in \{1, 2, 3, \dots, n\};$$

$$f^+(u'_i) = (f(u_i u'_i) + f(u'_i u')) \pmod{6n+2} = 2n + 6i - 2 \pmod{6n+2} \text{ for } i \in \{1, 2, 3, \dots, n\};$$

$$f^+(u) = ((\sum_{i=1}^n f(u_i u)) + f(u u')) \pmod{6n+2} = (4n^2 + 5n + 1) \pmod{6n+2};$$

$$f^+(u') = ((\sum_{i=1}^n f(u'_i u')) + f(u u')) \pmod{6n+2} = (4n^2 + 7n + 1) \pmod{6n+2}.$$

Since n is even, we can see from the division algorithm easily that there is an integer t for which $f^+(u) = (4n^2 + 5n + (6n+2)t) + 1$. Thus, $f^+(u)$ is an odd integer. Similarly, we can use the division algorithm to prove that $f^+(u')$ is an odd integer and $f^+(u_i)$ and $f^+(u'_i)$ are even integers. Since the sequences $\{6i-4\}_{i=1}^n$ and $\{6i-2\}_{i=1}^n$ are all distinct, it clearly that $\{f^+(u_i)\}_{i=1}^n$ and $\{f^+(u'_i)\}_{i=1}^n$ are also distinct.

Next, we will show that $f^+(u)$ and $f^+(u')$ are distinct. Suppose in the contrary that $f^+(u) \equiv f^+(u') \pmod{6n+2}$. Then, $4n^2 + 5n + 1 \equiv 4n^2 + 7n + 1 \pmod{6n+2}$. This implies that $2n \equiv 0 \pmod{6n+2}$, which is a contradiction. Thus, for all $i \in \{1, 2, 3, \dots, n\}$, $f^+(u_i)$, $f^+(u'_i)$, $f^+(u)$ and $f^+(u')$ are all distinct and they are in $\{0, 1, 2, \dots, 6n+1\}$. Therefore, the function f defined in Algorithm 2.2 is an edge-odd graceful labeling and this implies that $\text{Prism}(S_n)$ is an EOGG. $\square$

Lemma 2.3. *If $n \geq 3$ and $n \equiv 5 \pmod{6}$, then $\text{Prism}(S_n)$ is an EOGG.*

Proof. From Algorithm 2.3(i-v), the edge labels are arranged in the set $\{2n + 1, 2n + 3, 2n + 5, \dots, 4n - 1\}$, $\{1\}$, $\{3, 5, 7, \dots, 2n - 1\}$, $\{4n + 1, 4n + 3, 4n + 5, \dots, 6n - 1\}$ and $\{6n + 1\}$, respectively. Then, f is a bijection from $E(\text{Prism}(S_n))$ to $\{1, 3, 5, \dots, 6n - 1\}$.

Next, from Algorithm 2.3, we have

$$\begin{aligned} f^+(u_i) &= (f(u_i u'_i) + f(u_i u)) \pmod{6n+2} = 2n + 4i, \text{ for } i \in \{1, 2, 3, \dots, n-1\}; \\ f^+(u_n) &= 4n; \\ f^+(u'_i) &= (f(u_i u'_i) + f(u'_i u')) \pmod{6n+2} = 4i - 4, \text{ for } i \in \{1, 2, 3, \dots, n\}; \\ f^+(u) &= ((\sum_{i=1}^n f(u_i u)) + f(uu')) \pmod{6n+2} = (n^2 - 1) \pmod{6n+2}; \\ f^+(u') &= ((\sum_{i=1}^n f(u'_i u')) + f(uu')) \pmod{6n+2} = (5n^2 - 1) \pmod{6n+2}. \end{aligned}$$

Since $n = 6k - 1$ for some $k \in \mathbb{N}$, $f^+(u_i) = 2(6k - 1) + 4i = 4(3k + i - 1) + 2$ for all $i \in \{1, 2, 3, \dots, n - 1\}$. Then, $4 \nmid f^+(u_i)$. However, $4 \mid f^+(u'_i)$ for all $i \in \{1, 2, 3, \dots, n\}$, $4 \mid f^+(u_n)$, and $\max_{1 \leq i \leq n} \{f^+(u'_i)\} = 4n - 4$. We can conclude that $\{f^+(u_i) \mid i \in \{1, 2, 3, \dots, n\}\}$ and $\{f^+(u'_i) \mid i \in \{1, 2, 3, \dots, n\}\}$ are disjoint. Since $n = 6k - 1$ for some $k \in \mathbb{N}$, we can use the division algorithm to prove that $4 \mid f^+(u)$ and $4 \mid f^+(u')$. Then, to complete the prove, we will show that the values of $f^+(u)$ and $f^+(u')$ under the integer modulo $6n + 2$ are greater than $4n$.

Let k be an integer such that $n = 6k - 1$, then $6n + 2 = 36k - 4$, $n^2 - 1 = 36k^2 - 12k$ and $5n^2 - 1 = 180k^2 - 60k + 4$. Thus, $f^+(u) = (36k^2 - 12k) \pmod{36k - 4} = 28k - 4 > 24k - 4 = 4n$, and $f^+(u') = (5n^2 - 1) \pmod{6n + 2} = 32k - 4 > 24k - 4 = 4n$. Hence, for all $i \in \{1, 2, 3, \dots, n\}$, $f^+(u_i)$, $f^+(u'_i)$, $f^+(u)$ and $f^+(u')$ are distinct and they are in $\{0, 1, 2, \dots, 6n + 1\}$. Therefore, the function f defined in Algorithm 2.3 is an edge-odd graceful labeling and this implies that $\text{Prism}(S_n)$ is an EOGG. $\square$

Lemma 2.4. *If $n \geq 3$ and $n \equiv 1 \pmod{6}$, then $\text{Prism}(S_n)$ is an EOGG.*

Proof. From Algorithm 2.4(i-iv), the edge labels are arranged in the set $\{4n + 3, 4n + 5, 4n + 7, \dots, 6n + 1\}$, $\{3, 5, 7, \dots, 2n + 1\}$, $\{1\}$, $\{2n + 5, 2n + 7, 2n + 9, \dots, 4n + 1\}$ and $\{2n + 3\}$, respectively. Then, f is a bijection from $E(\text{Prism}(S_n))$ to $\{1, 3, 5, \dots, 6n - 1\}$.

Next, from Algorithm 2.4, we have

$$\begin{aligned} f^+(u_i) &= (f(u_i u'_i) + f(u_i u)) \pmod{6n+2} = (4n + 4i + 2) \pmod{6n+2}, \\ &\text{for } i \in \{1, 2, 3, \dots, n\}; \\ f^+(u) &= ((\sum_{i=1}^n f(u_i u)) + f(uu')) \pmod{6n+2} = (n^2 + 4n + 3) \pmod{6n+2}; \\ f^+(u'_1) &= 4n + 4; \\ f^+(u'_i) &= (f(u_i u'_i) + f(u'_i u')) \pmod{6n+2} = 4i, \text{ for } i \in \{2, 3, 4, \dots, n\}; \\ f^+(u') &= ((\sum_{i=1}^n f(u'_i u')) + f(uu')) \pmod{6n+2} = (3n^2 + 2n + 1) \pmod{6n+2}. \end{aligned}$$

Since $n = 6k + 1$ for some $k \in \mathbb{N}$,

$$\begin{aligned} \{f^+(u_i) \mid i \in \{1, 2, 3, \dots, \frac{n-1}{2}\}\} &\cup \{f^+(u_i) \mid i \in \{\frac{n+1}{2}, \frac{n+3}{2}, \frac{n+5}{2}, \dots, n\}\} \\ &= \{4n + 6, 4n + 10, 4n + 14, \dots, 6n\} \cup \{2, 6, 10, \dots, 2n\} \\ &= \{24k + 10, 24k + 14, 24k + 18, \dots, 36k + 6\} \cup \{2, 6, 10, \dots, 12k + 2\} \text{ and} \\ &\{f^+(u'_i) \mid i \in \{1, 2, 3, \dots, n\}\} \end{aligned}$$

$$= \{8, 12, 16, \dots, 4n, 4n+4\} = \{8, 12, 16, \dots, 24k+4, 24k+8\}.$$

Since $n = 6k + 1$ for some $k \in \mathbb{N}$, we have $6n + 2 = 36k + 8$, $n^2 + 4n + 3 = 36k^2 + 36k + 8 \equiv 28k + 8 \pmod{36k + 8}$ and $3n^2 + 2n + 1 = 108k^2 + 48k + 6 \equiv 24k + 6 \pmod{36k + 8}$. Then, $f^+(u) = 28k + 8$ and $f^+(u') = 24k + 6$. Hence, for all $i \in \{1, 2, 3, \dots, n\}$, $f^+(u_i)$, $f^+(u'_i)$, $f^+(u)$ and $f^+(u')$ are distinct and they are in $\{0, 1, 2, \dots, 6n + 1\}$. Therefore, the function f defined in Algorithm 2.4 is an edge-odd graceful labeling and this implies that $\text{Prism}(S_n)$ is an EOGG. $\square$

Lemma 2.5. *If $n \geq 3$ and $n \equiv 3 \pmod{6}$, then $\text{Prism}(S_n)$ is an EOGG.*

Proof. From Algorithm 2.5, the edge labels are arranged in the set $\{4n+3, 4n+5, 4n+7, \dots, 6n+1\} \cup \{3, 5, 7, \dots, 2n+1\} \cup \{2n+3, 2n+5, 2n+7, \dots, 4n+1\} \cup \{1\}$, respectively. Then f is a bijection from $E(\text{Prism}(S_n))$ to $\{1, 3, 5, \dots, 6n-1\}$.

Next, from Algorithm 2.5, we have

$$f^+(u_i) = (f(u_i u'_i) + f(u_i u)) \pmod{6n+2} = (4n+4i+2) \pmod{6n+2}, \text{ for } i \in \{1, 2, 3, \dots, n\};$$

$$f^+(u) = ((\sum_{i=1}^n f(u_i u)) + f(u u')) \pmod{6n+2} = (n^2 + 2n + 1) \pmod{6n+2};$$

$$f^+(u'_i) = (f(u_i u'_i) + f(u'_i u')) \pmod{6n+2} = 4i, \text{ for } i \in \{1, 2, 3, \dots, n\};$$

$$f^+(u') = ((\sum_{i=1}^n f(u'_i u')) + f(u u')) \pmod{6n+2} = (3n^2 + 2n + 1) \pmod{6n+2}.$$

Since $n = 6k + 3$ for some $k \in \mathbb{N}$, the similar argument as in Lemma 2.4 can show that $f^+(u_i)$, $f^+(u'_i)$, $f^+(u)$ and $f^+(u')$ are distinct and they are subsets of $\{0, 1, 2, \dots, 6n + 1\}$. Therefore, the function f defined in Algorithm 2.5 is an edge-odd graceful labeling and this implies that $\text{Prism}(S_n)$ is an EOGG. $\square$

The results from those of Lemma 2.1-2.5 can be concluded as the following theorem.

Theorem 2.1. *The $\text{Prism}(S_n)$ is an EOGG for every $n \geq 3$.*







Figure 8: The vertex-labeling for $\text{Prism}(S_6)$ induced by Algorithm 2.2

Figure 9: The vertex-labeling for $\text{Prism}(S_5)$ induced by Algorithm 2.3Figure 10: The vertex-labeling for $\text{Prism}(S_7)$ induced by Algorithm 2.4Figure 11: The vertex-labeling for $\text{Prism}(S_9)$ induced by Algorithm 2.5

3 Prism-like graph

In this section, we give a definition of $\text{Prism}_3(S_n)$ and an algorithm of each edge labeling of $\text{Prism}_3(S_n)$ for $n \geq 3$ and $n \equiv 2 \pmod{6}$. After that we prove

that the labeling given in this algorithm is an edge-odd graceful labeling.

Definition 3.1. For $n \geq 3$, let $S_n^{(1)}$ be a star and $S_n^{(2)}$ and $S_n^{(3)}$ be copies of $S_n^{(1)}$. Define $\text{Prism}_3(S_n)$ by joining $u^{(1)}$ of $S_n^{(1)}$ to the corresponding vertex $u^{(2)}$ of $S_n^{(2)}$, $u^{(2)}$ of $S_n^{(2)}$ to the corresponding vertex $u^{(3)}$ of $S_n^{(3)}$, each $u_i^{(1)}$ of $S_n^{(1)}$ to the corresponding vertex $u_i^{(2)}$ of $S_n^{(2)}$, and each $u_i^{(2)}$ of $S_n^{(2)}$ to the corresponding vertex $u_i^{(3)}$ of $S_n^{(3)}$ for all $i \in \{1, 2, 3, \dots, n\}$. Thus,

$$E(\text{Prism}_3(S_n)) = E(S_n^{(1)}) \cup E(S_n^{(2)}) \cup E(S_n^{(3)}) \cup \{u_i^{(1)}u_i^{(2)} \mid i \in \{1, 2, 3, \dots, n\}\} \\ \cup \{u_i^{(2)}u_i^{(3)} \mid i \in \{1, 2, 3, \dots, n\}\} \cup \{u^{(1)}u^{(2)}\} \cup \{u^{(2)}u^{(3)}\}.$$

Algorithm 3.1. Let $n \geq 3$ be an integer and $n \equiv 2 \pmod{6}$. We can easily obtain that $q = |E(\text{Prism}_3(S_n))| = 5n + 2$. Define $f : E(\text{Prism}_3(S_n)) \rightarrow \{1, 3, 5, \dots, 10n + 3\}$ by


- i $f(u^{(1)}u_i^{(1)}) = 4n + 2i - 1$ for $i \in \{1, 2, 3, \dots, n\}$;
- ii $f(u^{(2)}u_i^{(2)}) = 2i - 1$ for $i \in \{1, 2, 3, \dots, n\}$;
- iii $f(u^{(3)}u_i^{(3)}) = 6n + 2i - 1$ for $i \in \{1, 2, 3, \dots, n\}$;
- iv $f(u_i^{(1)}u_i^{(2)}) = 8n + 2i - 1$ for $i \in \{1, 2, 3, \dots, n\}$;
- v $f(u_i^{(2)}u_i^{(3)}) = 2n + 2i - 1$ for $i \in \{1, 2, 3, \dots, n\}$;
- vi $f(u^{(1)}u^{(2)}) = 10n + 3$;
- vii $f(u^{(2)}u^{(3)}) = 10n + 1$.

Theorem 3.1. If $n \geq 3$ and $n \equiv 2 \pmod{6}$, then $\text{Prism}_3(S_n)$ is an EOGG.

Proof. From Algorithm 3.1(i-vii), the edge labels are arranged in the set $\{4n + 1, 4n + 3, 4n + 5, \dots, 6n - 1\}$, $\{1, 3, 5, \dots, 2n - 1\}$, $\{6n + 1, 6n + 3, 6n + 5, \dots, 8n - 1\}$, $\{8n + 1, 8n + 3, 8n + 5, \dots, 10n - 1\}$, $\{2n + 1, 2n + 3, 2n + 5, \dots, 4n - 1\}$, $\{10n + 3\}$, and $\{10n + 1\}$, respectively. Then, f is a bijection from $E(\text{Prism}_3(S_n))$ to $\{1, 3, 5, \dots, 10n + 4\}$.

Next, from Algorithm 3.1, we have

$$\begin{aligned} f^+(u^{(1)}) &= ((\sum_{i=1}^n f(u_i^{(1)}u^{(1)})) + f(u^{(1)}u^{(2)})) \pmod{10n + 4} \\ &= (5n^2 - 1) \pmod{10n + 4}; \\ f^+(u^{(2)}) &= ((\sum_{i=1}^n f(u_i^{(2)}u^{(2)})) + f(u^{(1)}u^{(2)}) + f(u^{(2)}u^{(3)})) \pmod{10n + 4} \\ &= (n^2 - 4) \pmod{10n + 4}; \\ f^+(u^{(3)}) &= ((\sum_{i=1}^n f(u_i^{(3)}u^{(3)})) + f(u^{(2)}u^{(3)})) \pmod{10n + 4} \\ &= (7n^2 - 3) \pmod{10n + 4}; \\ f^+(u_1^{(2)}) &= (f(u_1^{(1)}u_1^{(2)}) + f(u_1^{(2)}u_1^{(3)}) + f(u_1^{(2)}u^{(2)})) \pmod{10n + 4} = 10n + 3; \end{aligned}$$

Figure 12: Edge-labelings for $\text{Prism}_3(S_8)$

$$f^+(u_i^{(2)}) = (f(u_i^{(1)}u_i^{(2)}) + f(u_i^{(2)}u_i^{(3)}) + f(u_i^{(2)}u^{(2)})) \pmod{10n+4} = 6i - 7 \text{ for } i \in \{2, 3, 4, \dots, n\};$$

$$f^+(u_i^{(1)}) = (f(u_i^{(1)}u_i^{(2)}) + f(u_i^{(1)}u^{(1)})) \pmod{10n+4} = 2n + 4i - 6;$$

$$\begin{aligned} f^+(u_i^{(3)}) &= (f(u_i^{(2)}u_i^{(3)}) + f(u_i^{(3)}u^{(3)})) \pmod{10n+4} \\ &= (8n + 4i - 2) \pmod{10n+4}. \end{aligned}$$

Since $n \equiv 2 \pmod{6}$, we can see from the division algorithm that for all $i \in \{1, 2, 3, \dots, n\}$, $f^+(u^{(2)})$, $f^+(u_i^{(1)})$, and $f^+(u_i^{(3)})$ are even integers and $f^+(u^{(1)})$, $f^+(u^{(3)})$, and $f^+(u_i^{(2)})$ are odd integers. First, we claim that $f^+(u^{(2)})$, $f^+(u_i^{(1)})$, and $f^+(u_i^{(3)})$ are distinct for all $i \in \{1, 2, 3, \dots, n\}$. Since $n = 6k + 2$ for some $k \in \mathbb{N}$, we have

$$f^+(u^{(2)}) = (36k^2 + 24k) \pmod{60k + 24};$$

$$f^+(u_i^{(1)}) = (12k + 4i - 2) \pmod{60k + 24};$$

$$f^+(u_i^{(3)}) = (48k + 4i + 14) \pmod{60k + 24}.$$

By the division algorithm, $4 \mid f^+(u^{(2)})$, $4 \nmid f^+(u_i^{(1)})$ and $4 \nmid f^+(u_i^{(3)})$. Suppose that $f^+(u_i^{(1)}) \equiv f^+(u_j^{(3)}) \pmod{10n+4}$ for some $i, j \in \{1, 2, 3, \dots, n\}$ and $i \neq j$. Then, $2n + 4i - 6 \equiv 8n + 4j - 2 \pmod{10n+4}$. This implies that $4(i - j - 1) = 6n + (10n + 4)t$ for some integer t . Since $i, j \in \{1, 2, 3, \dots, n\}$ and $i \neq j$, we have $-4 \leq 4(i - j - 1) \leq 4n - 8$. However, if $t < 0$, then $4(i - j - 1) = 6n + (10n + 4)t < -4n - 4$ and if $t \geq 0$, then $4(i - j - 1) = 6n + (10n + 4)t \geq 6n$. This is a contradiction.

Next, we claim that $f^+(u^{(1)})$, $f^+(u^{(3)})$, and $f^+(u_i^{(2)})$ are distinct. Since $n = 6k + 2$ for some $k \in \mathbb{N}$, $f^+(u_1^{(2)}) = 60k + 23$, $f^+(u^{(1)}) = (180k^2 + 120k + 19) \pmod{60k + 24} = 48k + 19$ and $f^+(u^{(3)}) = (252k^2 + 168k + 25) \pmod{60k + 24} = (12k^2 + 12k + 1) \pmod{60k + 24}$. Since $f^+(u^{(3)}) =$

$(12k^2 + 12k + 1) \pmod{60k + 24}$, by the division algorithm, there exist unique integers q and r such that $12k^2 + 12k + 1 = (60k + 24)q + r$. Then, $r = 6(2k^2 + 8k - 4)q + 1$. Notice that $f^+(u_1^{(2)}) = 6(10k + 3) + 5$, $f^+(u_i^{(2)}) = 6(i - 2) + 5$ for all $i \in \{2, 3, 4, \dots, n\}$, and $f^+(u^{(1)}) = 6(8k + 3) + 1$. Thus, $\{f^+(u_i^{(2)}) | i \in \{1, 2, 3, \dots, n\}\}$ and $\{f^+(u^{(1)}), f^+(u^{(3)})\}$ are distinct. Moreover, since $\max_{2 \leq i \leq n} \{6i - 7\} = 6n - 7 < 10n + 3 = f^+(u_1^{(2)})$, It's clearly that $f^+(u_1^{(2)})$ and $f^+(u_i^{(2)})$ are distinct.

Finally, to obtain the second claim we have to show that $f^+(u^{(1)})$ and $f^+(u^{(3)})$ are distinct. Suppose that $f^+(u^{(1)}) \equiv f^+(u^{(3)}) \pmod{10n + 4}$. Then, $5n^2 - 1 \equiv 7n^2 - 3 \pmod{10n + 4}$. This implies that $n^2 - 1 \equiv 0 \pmod{5n + 2}$ that is there exist $m \in \mathbb{Z}$ such that $n^2 - 1 = (5n + 2)m$. Since n is even, $n^2 - 1$ is odd and $5n + 2$ is even integer. This is a contradiction. Hence, $f^+(u^{(1)})$ and $f^+(u^{(3)})$ are distinct.

Therefore, the function f defined in Algorithm 3.1 is an edge-odd graceful labeling and this implies that $\text{Prism}_3(S_n)$ is an EOGG. $\square$




Figure 13: The vertex-labeling for $\text{Prism}_3(S_8)$ induced by Algorithm 3.1

4 Prism of wheel

In this section, we give a definition of $\text{Prism}(W_n)$ and an algorithm to label each edge of $\text{Prism}(W_n)$. Then, we prove that the $\text{Prism}(W_n)$ is an edge-odd graceful graph if $2|n$.

Definition 4.1. For $n \geq 3$, let W_n be a wheel graph and W'_n be a copy of W_n . Define $\text{Prism}(W_n)$, called the *prism* of W_n , by joining u of W_n to the corresponding vertex u' of W'_n and each u_i of W_n to the corresponding vertex u'_i of W'_n for all $i \in \{1, 2, 3, \dots, n\}$. Thus,

$$E(\text{Prism}(W_n)) = E(W_n) \cup E(W'_n) \cup \{u_i u'_i \mid i \in \{1, 2, 3, \dots, n\}\} \cup \{uu'\}.$$

Algorithm 4.1. Let $n \geq 3$ and $2|n$. We can easily to obtain that $q = |E(\text{Prism}(W_n))| = 5n + 1$. Define $f : E(G) \rightarrow \{1, 3, 5, \dots, 10n + 1\}$ by

- i $f(u_i u'_i) = 2i - 1$ for $i \in \{1, 2, 3, \dots, n\}$;
- ii $f(u_1 u_n) = 2n + 1$;
- iii $f(u_i u_{i+1}) = 4n - 2i + 1$ for $i \in \{1, 2, 3, \dots, n - 1\}$;
- iv $f(u'_1 u'_n) = 6n + 1$;
- v $f(u'_i u'_{i+1}) = 8n - 2i + 1$ for $i \in \{1, 2, 3, \dots, n - 1\}$;
- vi $f(u_1 u) = 4n + 1$;
- vii $f(u_i u) = 6n - 2i + 3$ for $i \in \{2, 3, 4, \dots, n\}$;
- viii $f(u'_1 u') = 8n + 1$;
- ix $f(u'_i u') = 10n - 2i + 3$ for $i \in \{2, 3, 4, \dots, n\}$;
- x $f(uu') = 10n + 1$.




Figure 14: Edge-labelings for $\text{Prism}(W_4)$

Theorem 4.1. If $n \geq 3$ and $2|n$, then $\text{Prism}(W_n)$ is an EOGG.

Proof. From Algorithm 4.1(i-x), the edge labels are arranged in the set $\{1, 3, 5, \dots, 2n - 1\}$, $\{2n + 1\}$, $\{4n - 1, 4n - 3, 4n - 5, \dots, 2n + 5, 2n + 3\}$, $\{6n + 1\}$, $\{8n - 1, 8n - 3, 8n - 5, \dots, 6n + 3\}$, $\{4n + 1\}$, $\{6n - 1, 6n - 3, 6n - 5, \dots, 4n + 3\}$, $\{8n + 1\}$, $\{10n - 1, 10n - 3, 10n - 5, \dots, 8n + 3\}$, and $\{10n + 1\}$, respectively. Then, f is a bijection from $E(\text{Prism}(W_n))$ to $\{1, 3, 5, \dots, 10n + 1\}$.

Next, from Algorithm 4.1, we have

$$\begin{aligned} f^+(u_1) &= (f(u_1u_n) + f(u_1u_2) + f(u_1u) + f(u_1u'_1)) \pmod{10n+2} = 0; \\ f^+(u_i) &= (f(u_{i-1}u_i) + f(u_iu_{i+1}) + f(u_iu) + f(u_iu'_i)) \pmod{10n+2} = 4n - 4i + 4 \\ &\text{for } i \in \{2, 3, 4, \dots, n\}; \end{aligned}$$

$$\begin{aligned} f^+(u'_1) &= (f(u'_1u'_n) + f(u'_1u'_2) + f(u'_1u') + f(u_1u'_1)) \pmod{10n+2} = 2n - 2; \\ f^+(u'_i) &= (f(u'_{i-1}u'_i) + f(u'_iu'_{i+1}) + f(u'_iu') + f(u_iu'_i)) \pmod{10n+2} = 6n - 4i + 2 \\ &\text{for } i \in \{2, 3, 4, \dots, n\}; \end{aligned}$$

$f^+(u) = ((\sum_{i=1}^n f(u_iu) + f(uu')) \pmod{10n+2} = (5n^2 - 1) \pmod{10n+2}$;
 $f^+(u') = ((\sum_{i=1}^n f(u'_iu') + f(uu')) \pmod{10n+2} = (9n^2 - 1) \pmod{10n+2}$;
 It's clearly that for all $i \in \{1, 2, 3, \dots, n\}$, $f^+(u_i)$ and $f^+(u'_i)$ are even integers and by the division algorithm, we have $f^+(u)$ and $f^+(u')$ are odd integers. First, we show that $f^+(u_i)$ and $f^+(u'_i)$ are distinct. Since the sequences $\{4n - 4i + 4\}_{i=2}^n, \{0\}, \{6n - 4i + 2\}_{i=2}^n$, and $\{2n - 2\}$ are all distinct, it clearly that $\{f^+(u_i)\}_{i=2}^n, \{f^+(u_1)\}, \{f^+(u'_i)\}_{i=2}^n$ and $\{f^+(u'_1)\}$ are also distinct. Next, we claim that $f^+(u)$ and $f^+(u')$ are distinct. Suppose in the contrary that $f^+(u) \equiv f^+(u') \pmod{10n+2}$. Then, $5n^2 - 1 \equiv 9n^2 - 1 \pmod{10n+2}$. This implies that $2n^2 \equiv 0 \pmod{5n+1}$

For $n \in \{4, 6, 8\}$, $2n^2 \pmod{5n+1}$ is 11, 10 and 5, respectively.

If $n = 10k$ for some $k \in \mathbb{N}$, $2n^2 \pmod{5n+1} = 200k^2 \pmod{50k+1} = 46k + 1$.

Similarly,

If $n = 10k + 2$ for some $k \in \mathbb{N}$, $2n^2 \pmod{5n+1} = 36k + 8$.

If $n = 10k + 4$ for some $k \in \mathbb{N}$, $2n^2 \pmod{5n+1} = 26k + 11$.

If $n = 10k + 6$ for some $k \in \mathbb{N}$, $2n^2 \pmod{5n+1} = 16k + 10$.

If $n = 10k + 8$ for some $k \in \mathbb{N}$, $2n^2 \pmod{5n+1} = 6k + 5$.

Thus, $2n^2 \not\equiv 0 \pmod{5n+1}$. Hence, $f^+(u)$ and $f^+(u')$ are distinct.

Therefore, the function f defined in Algorithm 4.1 is an edge-odd graceful labeling and this implies that $\text{Prism}(W_n)$ is an EOGG.

□




Figure 15: The vertex-labeling for $\text{Prism}(W_4)$ induced by Algorithm 4.1

5 Conclusion and Discussion

As we can see in this article that $\text{Prism}(S_n)$ is an EOGG for all $n \in \mathbb{N}$. However, we can only prove that if $n \equiv 2 \pmod{6}$, then $\text{Prism}_3(S_n)$ is an EOGG and if $2|n$, then $\text{Prism}(W_n)$ is an EOGG. One may try to find algorithms for edge labeling of $\text{Prism}_3(S_n)$ and $\text{Prism}(W_n)$ for other cases in such away that force $\text{Prism}_3(S_n)$ and $\text{Prism}(W_n)$ to be EOGG. Another way is extend the definition of $\text{Prism}_3(S_n)$ to be $\text{Prism}_m(S_n)$ for $m \in \mathbb{N}$ and investigate whether we can find the way to have the edge-odd graceful labeling for this type of Prism-like graph or not.

References

- [1] R. Boonklurb, S. Singhun and A. Wongpradit, *Edge-odd graceful labelings of Prism(S_n) and Shaft($n,1$) graphs*, 30th National Graduate Research Conference, 2013, 23–30.
- [2] R. B. Gnanajothi, *Topics in Graph Theory*, Ph.D. Thesis, Madurai Kamaraj University (1991).
- [3] A. Rosa, *Theory of graphs (International Symposium, Rome, July 1966)*, Gordon Breach, N.Y. and Dunod Paris, **170** (1967), 349–355.
- [4] S. Singhun, *Graphs with Edge-Odd Graceful Labelings*, International Mathematical Forum, **8**(2013), 577–582.
- [5] A. Solairaju, C. Vimala and A. Sasikala, *Edge-Odd Gracefulness of the Graph $S_2 \square S_n$* , International Journal of Computer Applications (0975 8887), **9** (2010), 1–3.
- [6] A. Solairaju and K. Chithra, *Edge-Odd Graceful Graphs*, *Electron. Notes Discrete Math*, **33** (2009), 15–20.