***East-West J. of Mathematics:*** Vol. 27, No 1 (2026) pp. 1-22
https://doi.org/10.36853/ewjm0419

# SEVERAL ITERATIVE SCHEMES FOR THE SPLIT FEASIBILITY WITH MULTIPLE OUTPUT SETS AND APPLICATIONS

**Nguyen Thi Quynh Anh**

*Academy of Security Enginneering and Technology,*
*Thuan Thanh, Bac Ninh, Viet Nam.*
*e-mail: quynhanhnguyen0178@gmail.com*

**Abstract**

In this paper, for solving the split feasibility problem with multiple output sets, defined by demiclosed strongly quasi-nonexpansive operators on Hilbert spaces, we propose some block-iterative schemes, using the extrapolated Landweber-type operators. The strong convergence is proved without the boundedly regular condition as well as the closedness property of the range of the transformation operators, assumed recently in the literature for the similar problems. We give a necessary and sufficient condition which ensures that a $k$th iterate is a solution. We also give an application of our results to solve the multiple-sets split feasibility problem (MSSFP) with multiple output level sets with computational experiments for illustration.

## 1 Introduction

In this paper we denote by $I$, $\langle\cdot,\cdot\rangle$, and $\|\cdot\|$, respectively, the identity operator, an inner product, and the corresponding norm in any Hilbert space $\mathcal{H}$. We also denote by $\mathrm{Fix}\,(T)$ the fixed point set of any operator $T$ on $\mathcal{H}$, i.e., $\mathrm{Fix}\,(T) = \{x \in \mathcal{H} : Tx = x\}$.

**Keywords:** Quasi-nonexpasive operator,Landweber-type operator , fixed point , demiclosedness property.
2010 AMS Mathematics Subject Classification: 47J05 , 47H09 , 49J30, 47H07, 47H10

We consider the split feasibility problem with multiple output sets, formulated as finding a point

$$p \in C := \cap_{i\in\mathcal{N}}\mathrm{Fix}(U_i),\ A_jp \in \mathrm{Fix}(V_j)\ \forall j \in \mathcal{M}, \tag{1}$$

where

- $U_i$ is a demiclosed and $\rho_{1,i}$-strongly quasi-nonexpansive operator on a real Hilbert space $\mathcal{H}_0$ for $i \in \mathcal{N} = \{1, ..., r\}$ with any positive integer $r$ such that $\rho_1 = \inf_{i\in\mathcal{N}} \rho_{1,i} > 0$.

- $V_j$ is a demiclosed and $\rho_{2,j}$-strongly quasi-nonexpansive operator on a real Hilbert space $\mathcal{H}_j$ for $j \in \mathcal{M} = \{1, ..., q\}$ where $q$ is any positive integer with $\rho_2 = \inf_{j\in\mathcal{M}} \rho_{2,j} > 0$.

- $A_j$ is a bounded linear operator from $\mathcal{H}_0$ into $\mathcal{H}_j$ with the spectral norm $\|A_j\| > 0$.

Throughout this paper, we assume that the solution set of (1), denoted by $\Gamma$, is non-empty.

This problem with demicontractive operators $U_i$ and $V_j$ has been very recently studied in [1]. To solve it, they suggested a Halpern method in combination with an Armijo-line search. With the additional assumptions: $p \in \Omega$, a closed and convex subset in $\mathcal{H}_0$, and $\mathcal{H}_j = \mathcal{K}$ for $j \in \mathcal{M}$, where $\mathcal{K}$ is another Hilbert space, this problem is called the multiple-operator split common fixed point problem, introduced and studied firstly by Brooke et al. [4]. To solve it, by assuming that $r = q$, designing an operator $R_j = U_jT_j$, where $T_j = I - \gamma_jA_j^*(V_j - I)A_j$ with $\gamma_j \in (0, 1/\|A_j\|^2)$, and using the general iterative scheme, introduced in [26], they suggested weakly convergent dynamic string-averaging CQ-methods.

Problem (1), when $\mathcal{H}_j = \mathcal{K}$ and $A_j = A$ for all $j \in \mathcal{M}$ with an additional requirement $Ap \in \cap_{j\in\mathcal{M}}\mathrm{Fix}\,(V_j)$, is called the multiple split common fixed point problem, that has been first studied by Censor and Segal [16] and developed by Wang with Xu [28] and Tang with Liu [27]. When $r$ and $q$ are sufficiently large, by using the block-iterative scheme [2], Cegielski [11] suggested a general iterative method, a block-iterative variant of Censor and Segal's method. It is well known that any sequence generated by one of these methods, does not converge strongly in general and depends on the value $\|A\|$, that is difficult to be calculated. One important approach for overcoming the difficulty in computation of $\|A\|$ is to use the extrapolated Landweber-type operator, introduced and studied in [12, 14, 8, 9] and references therein. The extrapolated Landweber-type operator related to $V_j$ is determined by

$$\mathcal{L}_{\sigma_j}\{V_j\}x = x + \sigma_j(x)(\mathcal{L}\{V_j\}x - x),\ x \in \mathcal{H}_0, \tag{2}$$

where the extrapolation function $\sigma_j : \mathcal{H}_0 \to [1, \infty)$ is bounded from above by $\tau_j$, defined by

$$\tau_j(x) = \left( \frac{\|A\|.\|(V_j - I)Ax\|}{\|A^*(V_j - I)Ax\|} \right)^2 \tag{3}$$

if $Ax \notin \text{Fix}(V_j)$ and 1, otherwise, where $A^* : \mathcal{K} \to \mathcal{H}_0$ is the adjoint operator to $A$, and

$$\mathcal{L}\{V_j\} = I + \frac{1}{\|A\|^2} A^*(V_j - I)A. \tag{4}$$

Obviously, $\tau_j(x) \geq 1$, since $\|A^*(V_j - I)Ax)\| \leq \|A\|\|(V_j - I)Ax\|$ and thus $\mathcal{L}_{\sigma_j}\{V_j\}$ is well defined. Note that $\mathcal{L}_{\tau_j}\{V_j\}$ does not depend on $\|A\|$. Further, in order to obtain a strong convergence sequence, Cegielski and Al-Musallam [8] combined the above approach with the steepest descent method as follows. Put $U_{j+r} = \mathcal{L}_{\sigma_j}\{T_j\}$ and $\mathcal{I} = \{1, ..., m\}$, where $m = r + q$. Then, $\Gamma = \cap_{i \in \mathcal{I}} \text{Fix}\,(U_i)$. Cegielski and Al-Musallam proposed a hybrid steepest-descent method, generating $x^{k+1} = (I - t_k F)S^k x^k, x^0 \in \mathcal{H}_0$, where $t_k$ satisfies the condition
(t) $t_k \in (0, 1)$ for all $k \geq 0$, $\lim_{k\to\infty} t_k = 0$ and $\sum_{k\geq 0} t_k = \infty$,
$F$ is $\eta$-strongly monotone and $l$-Lipschitz continuous on $\mathcal{H}_0$ with $l \geq \eta > 0$, and $S^k = I + \gamma_k(U_{i_k} - I)$ with $i_k \in \mathcal{I}$. They showed that any sequence, generated by this scheme, converges strongly to a point $p_* \in \mathcal{H}_0$, solving the variational inequality

$$p_* \in \Gamma : \ \langle Fp_*, p_* - p\rangle \leq 0 \ \forall p \in \Gamma, \tag{5}$$

under some conditions, two of which are that $U_i$ for each $i \in \mathcal{I}$ is approximately shrinking and the family $\mathcal{F} := \{\text{Fix}\,(U_i) : i \in \mathcal{I}\}$ is boundedly regular. Recently, in order to obtain a strong convergence result, Cegielski et al. [9] suggested the outer approximation method. They proved strong convergence of any sequence $\{x^k\}$, generated by this method, if $ImA, \text{Fix}(V_1), ..., \text{Fix}\,(V_q)\}$, $\{A^{-1}(Q), \text{Fix}(U_1), ..., \text{Fix}\,(U_r)\}$ are boundedly regular, and $ImA$ is closed, where $Q = \cap_{j\in\mathcal{M}} \text{Fix}(V_j)$.

In the case that $q = r$, problem (1) can be formulated in the following equivalent form: Find a point

$$p \in C_i := \text{Fix}(U_i), \ A_i p \in \text{Fix}(V_i) \ \forall i \in \mathcal{N}.$$

When $U_i$ and $V_j$ are cutters ($\rho_{1,i} = \rho_{2,j} = 1$), the authors of [22] proposed a weak convergent iterative method, $x^{k+1} = U_{i_k,\mu_k} T_{i_k,\lambda_k} x^k$, where the parameters $\mu_k, \lambda_k \in [\varepsilon, 2 - \varepsilon]$, $U_{i,\mu}$ is the $\mu$-relaxation of $U_i$, $T_{i,\lambda}$ is $\lambda$-relaxation of $\mathcal{L}\{V_i\}$, and $\{i_k\}_{k=0}^{\infty}$ is an almost cyclic control (cf. [11], Definition 5.6.10).

Recently, in the case that $r = 1$ and $U_1, T_j$ are demicontractive, in order to avoid the difficulties mentioned above, Wang [29] proposed an iterative method, by using the strategies of self-adaptive stepsizes in combination with the viscosity approximation method [23].

In this paper, still by using the extrapolated Landweber-type operators related to $V_j$ in combination with the steepest-descent method, we introduce some new block-iterative schemes, strong convergence of any sequence generated by one of which is proved without computation of any $\|A_j\|$. Moreover, we give a necessary and sufficient condition which confirms that $k$th iterate is a solution.

We organize the rest of this paper as follows. In Section 2, we list some necessary terminologies and related facts, that will be used in the proof of our results. In Section 3, we first describe our main block-iterative schemes and then show our main results with some modifications and particularities. An application to solving the split feasibility problem with multiple output level sets and numerical experiments are given in Section 4 for illustration.

## 2 Preliminaries and useful lemmas

We remember that an operator $T$ in $\mathcal{H}$ is called (see, [10]):

- contractive if $\|Tx - Ty\| \leq a\|x - y\|$ with $a \in (0, 1)$ for any $x, y \in \mathcal{H}$.
- quasi-nonexpansive, if $\mathrm{Fix}\,(T) \neq \emptyset$ and $\|Tx - p\| \leq \|x - p\|$ for all $x \in \mathcal{H}$ and $p \in \mathrm{Fix}(T)$.
- $\rho$-strongly quasi-nonexpansive ($\rho$-SQNE), if $\mathrm{Fix}\,(T) \neq \emptyset$ and, for all $x \in \mathcal{H}$ and $p \in \mathrm{Fix}(T)$, $\|Tx-p\|^2 \leq \|x-p\|^2-\rho\|Tx-x\|^2$ where the real number $\rho \geq 0$.
- $\tau$-demicontractive, if $\mathrm{Fix}\,(T) \neq \emptyset$ and it satisfies the condition

  $$\|Tx - p\|^2 \leq \|x - p\|^2 + \tau\|Tx - x\|^2 \ \forall x \in \mathcal{H}, p \in \mathrm{Fix}(T)$$

  with $\tau \in (0, 1)$.
- approximately shrinking (boundedly regular [17]) on a subset $D \subseteq \mathcal{H}$, if for any sequence $\{x^k\} \subseteq D$ the following implication holds

  $$\lim_{k\to\infty} \|Tx^k - x^k\| = 0 \Longrightarrow \lim_{k\to\infty} d(x^k, \mathrm{Fix}\,(T)) = 0, \tag{6}$$

  where $d(x, C) = \inf_{y\in C} \|x - y\|$ for any $x \in \mathcal{H}$ and $C \subset \mathcal{H}$.
- demiclosed, i.e., it satisfies the demiclosedness principle: if for any sequence $\{x^k\} \subset \mathcal{H}$ it holds

  $$w - \lim_{k\to\infty} x^k = x, \ \lim_{k\to\infty} \|(I - T)x^k\| = 0) \Longrightarrow Tx = x. \tag{7}$$

Clearly, a $\rho$-SQNE operator $T$ with $\rho = 0$ is quasi-nonexpansive and a nonexpansive operator $T$ with $\mathrm{Fix}\,(T) \neq \emptyset$ is quasi-nonexpansive. If an operator $T$ is quasi-nonexpansive or $\tau$-demicontractive on $H$, then the relaxation operator, defined by $\overline{T} = I + \beta(T - I)$, is $\rho$-SQNE with $\rho = \beta(1-\beta)$ for any fixed $\beta \in (0, 1]$ or $\rho = \beta(1 - \tau - \beta)$ for $\beta \in (0, 1 - \tau)$, respectively, and $\mathrm{Fix}\,(T) = \mathrm{Fix}\,(\overline{T})$ (see, [25, 24]). In this case $\mathrm{Fix}\,(T)$ is closed and convex. It is well known that (6) is only a necessary condition for implication (7) (see, [17]).

For each $i \in \mathcal{I} := \{1, ..., m\}$, let $C_i$ be a closed convex subset in $\mathcal{H}$ with $C := \cap_{i \in K} C_i \neq \emptyset$. We say that the family $\mathcal{C} = \{C_i : i \in \mathcal{I}\}$ is boundedly regular [17] if for any bounded sequence $\{x^k\} \subset \mathcal{H}$, there holds the following implication:

$$\lim_{k\to\infty} \max_{i \in K} d(u^k, C_i) = 0 \Longrightarrow \lim_{k\to\infty} d(u^k, C) = 0.$$

A mapping $F : \mathcal{H} \to \mathcal{H}$ is said to be $\eta$-strongly monotone and $l$-Lipschitz continuous, if it satisfies, respectively,

$$\langle Fx - Fy, x - y\rangle \geq \eta\|x - y\|^2,\ \|Fx - Fy\| \leq l\|x - y\|\ \forall x, y \in \mathcal{H}$$

with $l \geq \eta > 0$. It is well known that

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y\rangle\ \forall x, y \in \mathcal{H}.$$

**Lemma 2.1** ([31]) *Let $\mathcal{H}$ be a real Hilbert space and let $F$ be an $\eta$-strongly monotone and $l$-Lipschitz continuous operator on $\mathcal{H}$ with some positive constants $l \geq \eta > 0$. Let $T^{t,\mu} = I - t\mu F$. Then, for a fixed number $\mu \in (0, 2\eta/l^2)$ and any $t \in (0, 1)$, $T^{t,\mu}$ is a contractions with coefficient $1 - t\beta$, where $\beta = (1/2)\mu(2\eta - \mu l^2)$.*

**Lemma 2.2** ([30]) *Let $\{a_k\}$, $\{b_k\}$ and $\{c_k\}$ be sequences of real numbers such that, for all $k \geq 0$, $a_{k+1} \leq (1 - b_k)a_k + b_k c_k$; $a_k \geq 0$; $b_k$ satisfies a condition of type (t); and either $\sum_{k=1}^{\infty} b_k|c_k| < \infty$ or $\limsup_{k\to\infty} c_k \leq 0$. Then, $\lim_{k\to\infty} a_k = 0$.*

**Lemma 2.3** ([20]) *Let $\{c_k\}$ be a sequence of real numbers with a subsequence $\{l_k\}$ of $\{k\}$ such that $c_{l_k} < c_{l_k+1}$. Then, there exists a non-decreasing sequence $\{m_k\} \subseteq \{k\}$ such that $m_k \to \infty, c_{m_k} \leq c_{m_k+1}$ and $c_k \leq c_{m_k+1}$ for all (sufficiently large) numbers $k \geq 0$. In fact, $m_k = \max\{l \leq k : c_l \leq c_{l+1}\}$.*

**Lemma 2.4** ([15]) *Let $T_i : \mathcal{H} \to \mathcal{H}$ be a $\rho_i$-SQNE operator with $\rho_i > 0$ for all $i \in \mathcal{L} = \{1, ..., m\}$ for some integer $m \geq 1$ such that $\cap_{i\in\mathcal{L}}\mathrm{Fix}\,(T_i) \neq \emptyset$, $T := \sum_{i\in\mathcal{L}} \omega_i T_i$, where $\omega_i \geq 0$ for all $i \in \mathcal{L}$ and $\sum_{i\in\mathcal{L}} \omega_i = 1$, and let $\tilde{T} := T_m ... T_1$. Then, for any $x \in \mathcal{H}$ and $p \in \cap_{i\in\mathcal{L}}\mathrm{Fix}\,(T_i)$, we have*

$$\frac{1}{2R}\sum_{i\in\mathcal{L}} \omega_i\rho_i\|(T_i - I)x\|^2 \leq \|(T - I)x\|^2 \tag{8}$$

*and*

$$\frac{1}{2R}\sum_{i\in\mathcal{L}}\rho_i\|(T^i-T^{i-1})x\|^2\leq\|(\tilde{T}-I)x\|^2, \tag{9}$$

*where* $T^i=T_i...T_1, i\in L, T^0=I$ *and* $R\geq\|x-p\|$.

# 3 Main results

First of all, we formulate our block-iterative schemes.
*Block-iterative schemes:*

1. Choose any two points $x^{-1}$ and $x^0$ in $\mathcal{H}_0$ such that $x^{-1}\neq x^0$ and an integer $s\geq 1$. Set $k:=0$.

2. Set $x^{k,0}=x^k$, the current iterate. For $t=1,2,\cdots,s$, let $\mathcal{N}_t^k$ and $\mathcal{M}_t^k$ be two ordered non-empty subsets of $\mathcal{N}$ and $\mathcal{M}$, respectively, such that $\mathcal{N}=\mathcal{N}_1^k\cup\cdots\cup\mathcal{N}_s^k$ and $\mathcal{M}=\mathcal{M}_1^k\cup\cdots\cup\mathcal{M}_s^k$ and let define $x^{k,t}$ by the rule:
$$x^{k,t}=U_t^k y^{k,t},\ y^{k,t}=V_t^k x^{k,t-1}, \tag{10}$$
where either
$$U_t^k=\sum_{i^k(t)\in\mathcal{N}_t^k}\lambda_{i^k(t)}^k U_{i^k(t)}$$
or
$$U_t^k=U_{i^k_{|\mathcal{N}_t^k|}(t)}...U_{i_1^k(t)}$$
and either
$$V_t^k=\sum_{j^k(t)\in\mathcal{M}_t^k}\theta_{j^k(t)}^k\mathcal{L}_{\sigma_{j^k(t)}}\{V_{j^k(t)}\}$$
or
$$V_t^k=\mathcal{L}_{\sigma_{j^k_{|\mathcal{M}_t^k|}(t)}}\{V_{j^k_{|\mathcal{M}_t^k|}(t)}\}...\mathcal{L}_{\sigma_{j_1^k(t)}}\{V_{j_1^k(t)}\},$$
$\mathcal{N}_t^k=\{i_1^k(t),...,i_{|\mathcal{N}_t^k|}^k(t)\}$, $\mathcal{M}_t^k=\{j_1^k(t),...,j_{|\mathcal{M}_t^k|}^k(t)\}$, the parameters $\lambda_{i^k(t)}^k$ and $\theta_{j^k(t)}^k\geq\theta$ satisfy
$$\lambda_{i^k(t)}^k,\theta_{j^k(t)}^k\geq\theta>0,\ \sum_{i^k(t)\in\mathcal{N}_t^k}\lambda_{i^k(t)}^k=\sum_{j^k(t)\in\mathcal{M}_t^k}\theta_{j^k(t)}^k=1,$$
$\mathcal{L}_{\sigma_{j^k(t)}}\{V_{j^k(t)}\}$ is defined by (2)-(4) with $j=j^k(t)$ and $A=A_{j^k(t)}$, a function $\sigma_{j^k(t)}:\mathcal{H}_0\to[1,\infty)$ is an extrapolation functional bounded

from above by $\tau_{j^k(t)}$, defined by (3) with $V_{j^k(t)}$ and $A_{j^k(t)}$ instead of $V_j$ and $A$, respectively, i.e.,

$$\mathcal{L}_{\sigma_{j^k(t)}}\{V_{j^k(t)}\}x = x + \sigma_{j^k(t)}(x)(\mathcal{L}\{V_{j^k(t)}\}x - x),\ x \in \mathcal{H}_0,$$
$$\mathcal{L}\{V_{j^k(t)}\} = I + \frac{1}{\|A_{j^k(t)}\|^2}A^*_{j^k(t)}(V_{j^k(t)} - I)A_{j^k(t)},$$
$$\tau_{j^k(t)}(x) = \left(\frac{\|A_{j^k(t)}\|\|(V_{j^k(t)} - I)A_{j^k(t)}x\|}{\|A^*_{j^k(t)}(V_{j^k(t)} - I)A_{j^k(t)}x)\|}\right)^2$$

if $A_{j^k(t)}x \notin \mathrm{Fix}(V_{j^k(t)})$ and 1, otherwise.

If $x^{k,s} = x^k$, then stop. $x^k$ is a solution of (1). Otherwise, go to the next step.

3. Compute
$$x^{k+1} = (1 - \xi_k)(I - t_k\mu_k F)x^k + \xi_k x^{k,s}, \tag{11}$$
where $\xi_k \in [\xi, 1 - \xi]$ with a sufficiently small real number $\xi > 0$, $F$ is an $\eta$-strongly monotone and $l$-Lipschitz continuous operator on $\mathcal{H}_0$ with $l \geq \eta > 0$, $t_k$ satisfies $(t)$, and
$$\mu_k = \frac{\langle Fx^k - Fx^{k-1}, x^k - x^{k-1}\rangle}{\|Fx^k - Fx^{k-1}\|^2},$$
if $x^k \neq x^{k-1}$ and $\mu_{k-1}$, otherwise.
Set $x^{k-1} := x^k$, $x^k := x^{k+1}$ and $k := k + 1$. Return to step 2.

Further, we need the following lemmas.

**Lemma 3.1** *There holds the following inequality*
$$\|(I - V_j)A_jx\|^2 \leq \frac{2}{\rho_2 + 1}\langle A^*_j(I - V_j)A_jx, x - p\rangle,\ \forall x \in \mathcal{H}_0, \tag{12}$$
*for any* $p \in \Gamma$.

**Proof** This is followed from Lemmas 2.5, 2.6 in [14] and Fact 2.7, (i) in [13].

**Lemma 3.2** *For any sequence* $\{x^k\}$*, generated by one of our block-iterative schemes, we have:*
*(i)*
$$\|x^{k,s} - p\|^2 \leq \|x^k - p\|^2 - \tilde{\rho}\sum_{t=1}^{s}\left[\|(V^k_t - I)x^{k,t-1}\|^2 + \|(U^k_t - I)y^{k,t}\|^2\right], \tag{13}$$
*where* $p$ *is any point in* $\Gamma$, $\tilde{\rho} = \min\{\rho_1/r, \rho_2/q, \rho_2/(qa)\}$, *and* $a = \max_{1\leq j\leq q}\|A_j\|^2$.
*(ii)* $x^k$ *is a solution of (1) if and only if* $x^{k,s} = x^k$.

**Proof** (i) First, we consider the following case, when

$$U_t^k = \sum_{i^k(t)\in\mathcal{N}_t^k} \lambda_{i^k(t)}^k U_{i^k(t)},\ V_t^k = \sum_{j^k(t)\in\mathcal{M}_t^k} \theta_{j^k(t)}^k \mathcal{L}_{\sigma_{j^k(t)}}. \tag{14}$$

Using the definition of $y^{k,t}$ in (10), (12), the properties of $V_j$, and the definition of $\sigma_{k,t}, \theta_{j^k(t)}^k$, we get

$$\begin{aligned}
\|y^{k,t}-p\|^2 &= \|V_t^k x^{k,t-1}-p\|^2 \le \sum_{j^k(t)\in\mathcal{M}_t^k} \theta_{j^k(t)}^k \|\mathcal{L}_{\sigma_{j^k(t)}} x^{k,t-1}-p\|^2 \\
&= \sum_{j^k(t)\in\mathcal{M}_t^k} \theta_{j^k(t)}^k \|x^{k,t-1}-p \\
&\quad +\sigma_{j^k(t)}(x^{k,t-1})(\mathcal{L}\{V_{j^k(t)}\}-I)x^{k,t-1}\|^2 \\
&= \sum_{j^k(t)\in\mathcal{M}_t^k} \theta_{j^k(t)}^k \Bigg[\|x^{k,t-1}-p\|^2 \\
&\quad -2\frac{\sigma_{k,t}(x^{k,t-1})}{\|A_{j^k(t)}\|^2}\langle A_{j^k(t)}^*(I-V_{j^k(t)})A_{j^k(t)}x^{k,t-1}, x^{k,t-1}-p\rangle \\
&\quad +\frac{\sigma_{k,t}^2(x^{k,t-1})}{\|A_{j^k(t)}\|^4}\|A_{j^k(t)}^*(V_{j^k(t)}-I)A_{j^k(t)}x^{k,t-1}\|^2\Bigg] \\
&\le \sum_{j^k(t)\in\mathcal{M}_t^k} \theta_{j^k(t)}^k \Bigg[\|x^{k,t-1}-p\|^2 \\
&\quad -((\rho_2/q)+1)\frac{\sigma_{j^k(t)}(x^{k,t-1})}{\|A_{j^k(t)}\|^2}\|(V_{j^k(t)}-I)A_{j^k(t)}x^{k,t-1}\|^2 \\
&\quad +\frac{\sigma_{k,t}(x^{k,t-1})}{\|A_{j^k(t)}\|^2}\frac{\tau_{k,t}(x^{k,t-1})}{\|A_{j^k(t)}\|^2}\|A_{j^k(t)}^*(V_{j^k(t)}-I)A_{j^k(t)}x^{k,t-1}\|^2\Bigg] \\
&= \sum_{j^k(t)\in\mathcal{M}_t^k} \theta_{j^k(t)}^k \Bigg[\|x^{k,t-1}-p\|^2 \\
&\quad -((\rho_2/q)+1)\frac{\sigma_{j^k(t)}(x^{k,t-1})}{\|A_{j^k(t)}\|^2}\|(V_{j^k(t)}-I)A_{j^k(t)}x^{k,t-1}\|^2 \\
&\quad +\frac{\sigma_{k,t}(x^{k,t-1})}{\|A_{j^k(t)}\|^2}\|(V_{j^k(t)}-I)A_{j^k(t)}x^{k,t-1}\|^2\Bigg] \\
&= \sum_{j^k(t)\in\mathcal{M}_t^k} \theta_{j^k(t)}^k \Bigg[\|x^{k,t-1}-p\|^2 \\
&\quad -(\rho_2/q)\frac{\sigma_{j^k(t)}(x^{k,t-1})}{\|A_{j^k(t)}\|^2}\|(V_{j^k(t)}-I)A_{j^k(t)}x^{k,t-1}\|^2\Bigg].
\end{aligned}$$

From this, $\sigma_{k,t} \geq 1$, $a \geq \|A_{j^k(t)}\|^2$, $\|A_{j^k(t)}\| = \|A^*_{j^k(t)}\|$, and

$$\|A^*_{j^k(t)}\|\|(V_{j^k(t)} - I)A_{j^k(t)}x^{k,t-1}\| \geq \|A^*_{j^k(t)}(V_{j^k(t)} - I)A_{j^k(t)}x^{k,t-1}\|$$

we deduce that

$$\begin{aligned}
\|y^{k,t} - p\|^2 &\leq \|x^{k,t-1} - p\|^2 \\
&\quad -\frac{\rho_2}{qa}\sum_{j^k(t)\in\mathcal{M}^k_t} \theta^k_{j^k(t)} \frac{\sigma_{j^k(t)}(x^{k,t-1})}{\|A_{j^k(t)}\|^2}\|A^*_{j^k(t)}(V_{j^k(t)} - I)A_{j^k(t)}x^{k,t-1}\|^2 \\
&\leq \|x^{k,t-1} - p\|^2 \\
&\quad -\frac{\rho_2}{qa}\left\|\sum_{j^k(t)\in\mathcal{M}^k_t} \theta^k_{j^k(t)} \frac{\sigma_{j^k(t)}(x^{k,t-1})}{\|A_{j^k(t)}\|^2}A^*_{j^k(t)}(V_{j^k(t)} - I)A_{j^k(t)}x^{k,t-1}\right\|^2 \\
&= \|x^{k,t-1} - p\|^2 - \frac{\rho_2}{qa}\|(V^k_t - I)x^{k,t-1}\|^2.
\end{aligned}$$

Then,

$$\begin{aligned}
\|x^{k,t} - p\|^2 &\leq \sum_{i^k(t)\in\mathcal{N}^k_t} \lambda^k_{i^k(t)}\|U_{i^k(t)}y^{k,t} - p\|^2 \\
&\leq \|y^{k,t} - p\|^2 - \sum_{i^k(t)\in\mathcal{N}^k_t} \lambda^k_{i^k(t)}\rho_{i^k(t)}\|(U_{i^k(t)} - I)y^{k,t}\|^2 \\
&\leq \|y^{k,t} - p\|^2 - \rho_1\sum_{i^k(t)\in\mathcal{N}^k_t} \lambda^k_{i^k(t)}\|(U_{i^k(t)} - I)y^{k,t}\|^2 \\
&\leq \|y^{k,t} - p\|^2 - \rho_1\left\|\sum_{i^k(t)\in\mathcal{N}^k_t} \lambda^k_{i^k(t)}(U_{i^k(t)} - I)y^{k,t}\right\|^2 \\
&= \|y^{k,t} - p\|^2 - \rho_1\|(U^k_t - I)y^{k,t}\|^2.
\end{aligned}$$

This together with (15) implies that

$$\begin{aligned}
\|x^{k,t} - p\|^2 &\leq \|x^{k,t-1} - p\|^2 - \rho_1\|(U^k_t - I)y^{k,t}\|^2 \\
&\quad -\tfrac{\rho_2}{qa}\|(V^k_t - I)x^{k,t-1}\|^2.
\end{aligned} \tag{15}$$

Now, we consider the case when $U^k_t$ and $V^k_t$ are defined by

$$\begin{aligned}
U^k_t &= U_{i^k_{|\mathcal{N}^k_t|}(t)}...U_{i^k_1(t)}, \\
V^k_t &= \mathcal{L}_{\sigma_{j^k_{|\mathcal{M}^k_t|}(t)}}\{V_{j^k_{|\mathcal{M}^k_t|}(t)}\}...\mathcal{L}_{\sigma_{j^k_1(t)}}\{V_{j^k_1(t)}\}.
\end{aligned} \tag{16}$$

Then, the $\rho_1$-SQNE property of $U_i$ yields

$$\begin{aligned}\|x^{k,t}-p\|^2 &= \|U_t^k y^{k,t}-p\|^2 = \|U_{i^k_{|L^k_t|}(t)}U_{i^k_{|L^k_t|-1}(t)}...U_{i^k_1(t)}y^{k,t}-p\|^2\\ &\le \|U_{i^k_{|L^k_t|-1}(t)}...U_{i^k_1(t)}y^{k,t}-p\|^2\\ &\quad -\rho_1\|U_{i^k_{|L^k_t|}(t)}U_{i^k_{|L^k_t|-1}(t)}...U_{i^k_1(t)}y^{k,t}-U_{i^k_{|L^k_t|-1}(t)}...U_{i^k_1(t)}y^{k,t}\|^2\\ &\le \|y^{k,t}-p\|^2-\rho_1\sum_{h=1}^{|\mathcal{N}_t^k|}\|(U^{i^k_h(t)}-U^{i^k_{h-1}(t)})y^{k,t}\|^2\\ &\le \|y^{k,t}-p\|^2-\frac{\rho_1}{r}\|(U_t^k-I)y^{k,t}\|^2\end{aligned}$$

where $U^{i^k_h(t)}$ being a product of the first $h$ operators along the string $\mathcal{N}_t^k$ and $U^{i^k_0(t)}=I$, because $|\mathcal{N}_t^k|\le r$ and

$$\|(U_t^k-I)y^{k,t}\|^2\le|\mathcal{N}_t^k|\sum_{h=1}^{|\mathcal{N}_t^k|}\|(U^{i^k_h(t)}-U^{i^k_{h-1}(t)})y^{k,t}\|^2.$$

Similarly, we get

$$\begin{aligned}\|y^{k,t}-p\|^2 &= \|\mathcal{L}_{\sigma_{j^k_{|\mathcal{M}_t^k|}(t)}}\{V_{j^k_{|\mathcal{M}_t^k|}(t)}\}...\mathcal{L}_{\sigma_{j^k_1(t)}}\{V_{j^k_1(t)}\}x^{k,t-1}-p\|^2\\ &\le \|\mathcal{L}_{\sigma_{j^k_{|\mathcal{M}_t^k|-1}(t)}}\{V_{j^k_{|\mathcal{M}_t^k|}(t)}\}...\mathcal{L}_{\sigma_{j^k_1(t)}}\{V_{j^k_1(t)}\}x^{k,t-1}-p\|^2\\ &\quad -\rho_2\big\|\big(\mathcal{L}_{\sigma_{j^k_{|\mathcal{M}_t^k|}(t)}}\{V_{j^k_{|\mathcal{M}_t^k|}(t)}\}...\mathcal{L}_{\sigma_{j^k_1(t)}}\{V_{j^k_1(t)}\}\\ &\quad -\mathcal{L}_{\sigma_{j^k_{|\mathcal{M}_t^k|-1}(t)}}\{V_{j^k_{|\mathcal{M}_t^k|}(t)}\}...\mathcal{L}_{\sigma_{j^k_1(t)}}\{V_{j^k_1(t)}\}x^{k,t-1}\big)x^{k,t-1}\big\|^2\\ &\le \|x^{k,t-1}-p\|^2-\rho_2\sum_{h=1}^{|\mathcal{M}_t^k|}\|(\mathcal{L}^{i^k_h(t)}-\mathcal{L}^{i^k_{h-1}(t)})x^{k,t-1}\|^2, \qquad (17)\end{aligned}$$

where $\mathcal{L}^{i^k_h(t)}$ is defined similarly as $U^{i^k_h(t)}$. Since

$$\|(V_i^k-I)x^{k,t-1}\|^2\le q\sum_{h=1}^{|\mathcal{M}_t^k|}\|(\mathcal{L}^{i^k_h(t)}-\mathcal{L}^{i^k_{h-1}(t)})x^{k,t-1}\|^2,$$

from (17) we obtain (15) with $a=1$. Therefore,

$$\begin{aligned}\|x^{k,t}-p\|^2 &\le \|x^{k,t-1}-p\|^2-\rho_1\sum_{h=1}^{|L_t^k|}\|(U^{i^k_h(t)}-U^{i^k_{h-1}(t)})y^{k,t}\|^2\\ &\quad -\frac{\rho_2}{q}\|(V_t^k-I)x^{k,t-1}\|^2. \qquad (18)\end{aligned}$$

Noting $x^{k,0} = x^k$ and summing each (16) and (18) with $t = 1, ..., s$, we obtain (13). It is easy to see that in the rest cases, when $U_t^k, V_t^k$ are given by

$$U_t^k = \sum_{i^k(t)\in\mathcal{N}_t^k} \lambda_{i^k(t)}^k U_{i^k(t)},\ V_t^k = \mathcal{L}_{\sigma_{j^k_{|\mathcal{M}_t^k|}(t)}}\{V_{j^k_{|\mathcal{M}_t^k|}(t)}\}...\mathcal{L}_{\sigma_{j_1^k(t)}}\{V_{j_1^k(t)}\}, \quad (19)$$

or

$$U_t^k = U_{i^k_{|\mathcal{N}_t^k|}(t)}...U_{i_1^k(t)},\ V_t^k = \sum_{j^k(t)\in\mathcal{M}_t^k} \theta_{j^k(t)}^k \mathcal{L}_{\sigma_{j^k(t)}}, \quad (20)$$

we also obtain the same results.
(ii) It is obvious that if $x^k \in \Gamma$ then $x^{k,s} = x^k$. Inversely, we have to prove that if $x^{k,s} = x^k$ then $x^k \in \Gamma$. From (13) and $x^{k,s} = x^k$ we deduce that

$$\|y^{k,t} - x^{k,t-1}\| = \|(V_t^k - I)x^{k,t-1}\| = 0,\ \|(U_t^k - I)y^{k,t}\| = 0, \quad (21)$$

for $t = 1, ..., s$. On the other hand, by (10),

$$\|x^{k,t} - x^{k,t-1}\| = \|U_t^k y^{k,t} - x^{k,t-1}\| \le \|(U_t^k - I)y^{k,t}\| + \|(V_t^k - I)x^{k,t-1}\|,$$

for $t = 1, ..., s$. Then, by (21) $\|x^{k,t-1} - x^k\| = 0$. This and (21) yield

$$(U_t^k - I)x^k = 0,\ (V_t^k - I)x^k = 0, \quad (22)$$

for $t = 1, ..., s$.

Now, for arbitrary fixed $i \in \mathcal{N}$, $j \in \mathcal{M}$, and $k \ge 0$, from the definition of the proposed iterative schemes there exist integers $t_1^k, t_2^k \in \{1, ..., s\}$ such that $i \in \mathcal{N}_{t_1^k}^k$ and $j \in \mathcal{M}_{t_2^k}^k$. When $U_t^k$ and $V_t^k$ are defined by (14), from (8) and (22) we obtain

$$0 \le \frac{\rho_{1,i}\theta}{2R}\|(U_i - I)x^k\|^2 \le \|(U_{t_1^k}^k - I)x^k\|^2 = 0,$$
$$0 \le \frac{\rho_{2,j}\theta}{2R}\|(\mathcal{L}_{\sigma_j}\{V_j\} - I)x^k\|^2 \le \|(V_{t_2^k}^k - I)x^k\|^2 = 0.$$

It means that $x^k \in \text{Fix}(U_i)$ and $\|(\mathcal{L}_{\sigma_j}\{V_j\} - I)x^k\|^2 = 0$, which and (4) yield $A_j^*(V_j - I)A_j x^k = 0$. The last equality and (12) yield $(V_j - I)A_j x^k = 0$. It means that $A_j x^k \in \text{Fix}(V_j)$. Since $i$ and $j$ are arbitrary in $\mathcal{N}$ and $\mathcal{M}$, respectively, $x^k \in \Gamma$. When $U_{t_1^k}^k$ and $V_t^k$ are given by (16) we have

$$\|(U_{t_1^k}^k - I)x^k\|^2 = \|(U_{i^k_{|\mathcal{N}_{t_1^k}^k|}(t_1^k)}...U_i...U_{i_1^k(t_1^k)} - I)x^k\|^2 = 0,$$
$$\|(V_{t_2^k}^k - I)x^k\|^2 = \|(\mathcal{L}_{\sigma_j}\{V_j\} - I)x^k\|^2 = \|(\mathcal{L}_{\sigma_{j^k_{|\mathcal{M}_{t_2^k}^k|}(t_2^k)}}\{V_{j^k_{|\mathcal{M}_{t_2^k}^k|}(t_2^k)}\}$$
$$...\mathcal{L}_{\sigma_j}\{V_j\}...\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\} - I)x^k\|^2 = 0, \quad (23)$$

where $i$ coincides at least with one of $\{i_1^k(t_1^k), ..., i^k_{|\mathcal{N}^k_{t_1^k}|}(t_1^k)\}$ and $j$ is one of $\{j_1^k(t_2^k), ..., j^k_{|\mathcal{M}^k_{t_2^k}|}(t_2^k)\}$. Without loss of generality, we can write

$$\mathcal{N}^k_{t_1^k} = \{i_1^k(t_1^k), ..., i_l^k(t_1^k), i, ..., i^k_{|\mathcal{N}^k_t|}(t_1^k)\},$$
$$\mathcal{M}^k_{t_2^k} = \{j_1^k(t_2^k), ..., j_{l'}^k(t_2^k), j, ..., j^k_{|\mathcal{M}^k_t|}(t_2^k)\}.$$

By using (9) with $\tilde{T} = U^k_{t_1^k}$ and $\tilde{T} = V^k_{t_2^k}$, that satisfy (23), we get, respectively,

$$\begin{aligned} 0 \leq \quad & e\big[\|(U_i U_{i_l^k(t_1^k)}...U_{i_1^k(t_1^k)} - U_{i_l^k(t_1^k)}...U_{i_1^k(t_1^k)})x^k\|^2 + \\ & \|(U_{i_l^k(t_1^k)}...U_{i_1^k(t_1^k)} - U_{i_{l-1}^k(t_1^k)}...U_{i_1^k(t_1^k)})x^k\|^2 + ... + \\ & \|(U_{i_2^k(t_1^k)}U_{i_1^k(t_1^k)} - U_{i_1^k(t_1^k)})x^k\|^2 + \|(U_{i_1^k(t_1^k)} - I)x^k\|^2\big] \leq 0, \end{aligned}$$

where $e = \theta\rho_1/(2\tilde{R})$ and

$$\begin{aligned} 0 \leq \quad & e'\big[\|\big(\mathcal{L}_{\sigma_j}\{V_j\}\mathcal{L}_{\sigma_{j_{l'}^k(t_2^k)}}\{V_{j_{l'}^k(t_2^k)}\}...\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\} \\ & \quad -\mathcal{L}_{\sigma_{j_{l'}^k(t_2^k)}}\{V_{j_{l'}^k(t_2^k)}\}...\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\}\big)x^k\|^2 + \\ & \|\big(\mathcal{L}_{\sigma_{j_{l'}^k(t_2^k)}}\{V_{j_{l'}^k(t_2^k)}\}...\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\} \\ & \quad -\mathcal{L}_{\sigma_{j_{l'-1}^k(t_2^k)}}\{V_{j_{l'-1}^k(t_2^k)}\}...\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\}\big)x^k\|^2 + ... + \\ & \|\big(\mathcal{L}_{\sigma_{j_2^k(t_2^k)}}\{V_{j_2^k(t_2^k)}\}\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\} - \mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\}\big)x^k\|^2 \\ & + \|\big(\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\} - I\big)x^k\|^2\big] \leq 0, \end{aligned}$$

where $e' = \theta\rho_2/(2\tilde{R})$ with some constant $\tilde{R} > 0$. Hence,

$$\begin{aligned} & \|(U_i U_{i_l^k(t_1^k)}...U_{i_1^k(t_1^k)} - U_{i_l^k(t_1^k)}...U_{i_1^k(t_1^k)})x^k\|^2 = \\ & \|(U_{i_l^k(t_1^k)}...U_{i_1^k(t_1^k)} - U_{i_{l-1}^k(t_1^k)}...U_{i_1^k(t_1^k)})x^k\|^2 = ... = \\ & \|(U_{i_2^k(t_1^k)}U_{i_1^k(t_1^k)} - U_{i_1^k(t_1^k)})x^k\|^2 = \|(U_{i_1^k(t_1^k)} - I)x^k\|^2 = 0 \end{aligned}$$

and

$$\begin{aligned} & \|\big(\mathcal{L}_{\sigma_j}\{V_j\}\mathcal{L}_{\sigma_{j_{l'}^k(t_2^k)}}\{V_{j_{l'}^k(t_2^k)}\}...\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\} \\ & \quad -\mathcal{L}_{\sigma_{j_{l'}^k(t_2^k)}}\{V_{j_{l'}^k(t_2^k)}\}...\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\}\big)x^k\|^2 = \\ & \|\big(\mathcal{L}_{\sigma_{j_{l'}^k(t_2^k)}}\{V_{j_{l'}^k(t_2^k)}\}...\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\} \\ & \quad -\mathcal{L}_{\sigma_{j_{l'-1}^k(t_2^k)}}\{V_{j_{l'-1}^k(t_2^k)}\}...\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\}\big)x^k\|^2 = ... = 0, \\ & \|\big(\mathcal{L}_{\sigma_{j_2^k(t_2^k)}}\{V_{j_2^k(t_2^k)}\}\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\} - \mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\}\big)x^k\|^2 = 0, \\ & \|\big(\mathcal{L}_{\sigma_{j_1^k(t_2^k)}}\{V_{j_1^k(t_2^k)}\} - I\big)x^k\|^2\big] = 0. \end{aligned}$$

Consequently, $\|(U_i - I)x^k\|^2 = 0$ and $\|(\mathcal{L}_{\sigma_j}\{V_j\} - I)x^k\|^2 = 0$. Therefore, $x^k \in \text{Fix}(U_i)$ and $A_j x^k \in \text{Fix}(T_j)$.

Evidently, in the rest cases, when $U_t^k$ and $V_t^k$ are defined by (19) and (20), we also get the same results. This completes the proof.

**Lemma 3.3** *For any $p \in \Gamma$, the following inequality holds*

$$\begin{aligned}\|x^{k+1} \quad -p\|^2 \leq [1-(1-\theta_k)\alpha_k\beta_k]\|x^k-p\|^2 + 2(1-\theta_k)\alpha_k\big[\langle Fp, p-x^k\rangle \\ +\alpha_k\|Fp\|M_1\big] - \tilde{\rho}\textstyle\sum_{t=1}^{s}\big[\|(V_t^k - I)x^{k,t-1}\|^2 + \|(U_t^k - I)y^{k,t}\|^2\big] \quad (24)\end{aligned}$$

*for all $k \geq \tilde{k}$, a positive integer, where $\alpha_k = t_k\mu_k$ and $\beta_k = (1/2)(2\eta - \alpha_k l^2)$ such that $\alpha_k \in (0, \eta/l^2)$, $\eta \geq \beta_k \geq \eta/2$, and $1 - \alpha_k\beta_k \geq \tilde{c}$ for all $k \geq \tilde{k}$, and $M_1$, $\tilde{c}$ are positive real numbers.*

**Proof** Clearly, from (13) it follows that $\|x^{k,s} - p\| \leq \|x^k - p\|$ for all $k \geq 0$. We put

$$\eta_k = \frac{\langle Fx^k - Fx^{k-1}, x^k - x^{k-1}\rangle}{\|x^k - x^{k-1}\|^2},\ l_k = \frac{\|Fx^k - Fx^{k-1}\|}{\|x^k - x^{k-1}\|}.$$

Then,

$$\eta \leq \eta_k \leq \frac{\|Fx^k - Fx^{k-1}\|}{\|x^k - x^{k-1}\|} = l_k \leq l,$$

and hence

$$\frac{\eta}{l^2} \leq \mu_k = \frac{\eta_k}{l_k^2} \leq \frac{1}{\eta_k} \leq \frac{1}{\eta}. \quad (25)$$

Since $t_k \to 0$ as $k \to \infty$, by (25), there exists a positive integer $\tilde{k}$ such that $\alpha_k \in (0, \eta/l^2)$ and $\eta \geq \beta_k \geq \eta/2$ for all $k \geq \tilde{k}$. Thus, by using Lemma 2.1 and (25), we obtain

$$\begin{aligned}\|x^{k+1} - p\| \quad &= \|(1-\theta_k)(I - t_k\mu_k F)x^k + \theta_k x^{k,s} - p\| \\ &\leq (1-\theta_k)\|(I - t_k\mu_k F)x^k - p\| + \theta_k\|x^{k,s} - p\| \\ &= (1-\theta_k)\|(I - \alpha_k F)x^k - (I - \alpha_k F)p - \alpha_k Fp\| \\ &\quad +\theta_k\|x^{k,s} - p\| \\ &\leq (1-\theta_k)\big[(1-\alpha_k\beta_k)\|x^k - p\| + \alpha_k\|Fp\|\big] + \theta_k\|x^k - p\| \\ &= [1-(1-\theta_k)\alpha_k\beta_k]\|x^k - p\| + (1-\theta_k)\alpha_k\beta_k\|Fp\|/\beta_k \\ &\leq [1-(1-\theta_k)\gamma_k\beta_k]\|x^k - p\| + (1-\theta_k)\gamma_k\beta_k 2\|Fp\|/\eta \\ &\leq r' := \max\{\|x^{\tilde{k}} - p\|, 2\|Fp\|/\eta\}\ \forall k \geq \tilde{k}.\end{aligned}$$

It means that $\{x^k\}$ is bounded. Hence, $\{Fx^k\}$ is bounded. Thus, $\|Fx^k\| \leq M_1$

for all $k \geq 0$ where $M_1$ is some positive constant. Next, again by Lemma 2.1,

$$\begin{aligned}\|x^{k+1}-p\|^2 &\leq (1-\theta_k)\|(I-t_k\mu_k F)x^k - p\|^2 + \theta_k\|x^{k,s}-p\|^2\\ &= (1-\theta_k)\|(I-\alpha_k F)x^k - (I-\alpha_k F)p - \alpha_k Fp\|^2\\ &\qquad +\theta_k\|x^{k,s}-p\|^2\\ &\leq (1-\theta_k)\big[(1-\alpha_k\beta_k)\|x^k-p\|^2 + 2\alpha_k\big(\langle Fp, p-x^k\rangle\\ &\qquad +\alpha_k\|Fp\|M_1\big)\big]+\theta_k\|x^{k,s}-p\|^2. \qquad (26)\end{aligned}$$

Replacing $\|x^{k,s}-p\|^2$ in (26) by its upper bound in (13) we obtain (24). The proof is completed.

Now, we are in the position to prove strong convergence of any sequence, generated by one of our iterative schemes.

**Theorem 3.1** *Any sequence $\{x^k\}$, generated by one of our block-iterative schemes, as $k \to \infty$, converges strongly to a point $p_* \in \Gamma$, satisfying (5).*

**Proof** We need only to consider two cases.

*Case 1.* $\|x^{k+1}-p_*\| \leq \|x^k - p_*\|$ for all $k \geq \tilde{k}$, the integer in Lemma 3.3

Then, $\{x^k\}$ is bounded and $\lim_{k\to\infty}\|x^k-p_*\|$ exists. From (24) and the properties of $\theta_k, \alpha_k, \beta_k$ it follows that

$$d^k \leq \|x^k-p_*\|^2 - \|x^{k+1}-p_*\|^2 + d_k\alpha_k,$$

for all $k \geq \overline{k}$, a sufficiently large integer ($\overline{k} \geq \tilde{k}$), where $d_k = 2\|Fp_*\|(r'+\alpha_k M_1)$ and

$$d^k = \tilde{\rho}\sum_{t=1}^{s}\big[\|(V_t^k - I)x^{k,t-1}\|^2 + \|(U_t^k - I)y^{k,t}\|^2\big].$$

We prove that $\lim_{k\to\infty} d^k = 0$. Indeed, if $d^k \leq d_k\alpha_k$ for all $k \geq \overline{k}$ then the limit exists and equal to zero. Otherwise, i.e. $d^k > d_k\alpha_k$, we have

$$0 \leq \sum_{k=\tilde{k}}^{N}(d^k - d_k\alpha_k) \leq \|x^{\tilde{k}} - p_*\|^2 - \|x^{N+1}-p_*\|^2,$$

for any positive integer $N$. Thus, $\sum_{k=0}^{\infty}(d^k - d_k\alpha_k) \leq \|x^{\tilde{k}}-p_*\|^2$. Therefore, $\lim_{k\to\infty}(d^k - d_k\alpha_k) = 0$, that yields $\lim_{k\to\infty} d^k = 0$. This is equivalent to

$$\lim_{k\to\infty}\|(U_t^k - I)y^{k,t-1}\| = 0,\ \lim_{k\to\infty}\|(V_t^k - I)x^{k,t-1}\|^2 = 0,$$

for $t = 1, ..., s$. Next, by the same argument, as in the proof of (21) and (22) in Lemma 3.2, we have

$$\lim_{k\to\infty}\|y^{k,t} - x^{k,t-1}\| = 0,\ \lim_{k\to\infty}\|x^{k,t-1} - x^k\| = 0.$$

Hence,

$$\lim_{k\to\infty}\|(U_t^k - I)x^k\| = 0,\ \lim_{k\to\infty}\|(V_t^k - I)x^k\| = 0, \tag{27}$$

for $t = 1, ..., s$.

Since $\{x^k\}$ is a bounded sequence in the Hilbert space $\mathcal{H}_0$, there exists a subsequence $\{x^{n_k}\}$ of $\{x^k\}$ such that $\{x^{n_k}\}$ converges weakly to a point $\tilde{p} \in \mathcal{H}_0$ as $k \to \infty$. First, we prove that $\tilde{p} \in \Gamma$. Let $t_1^k, t_2^k, \mathcal{N}_{t_1^k}^k$, and $\mathcal{M}_{t_2^k}^k$ be as the above. Replacing $k$ and $t$, respectively, by $n_k$ and $t_1^k, t_2^k$ in (27), we obtain

$$\lim_{k\to\infty}\|(U_{t_1^k}^{n_k} - I)x^{n_k}\| = 0,\ \lim_{k\to\infty}\|(V_{t_2^k}^{n_k} - I)x^{n_k}\| = 0. \tag{28}$$

In the case that $U_t^k$ and $V_t^k$ are given by (14), using (8) with $T = U_{t_1^k}^{n_k}$ and $T = V_{t_2^k}^{n_k}$, we get

$$\begin{aligned}
&\frac{\theta\rho_1}{2R}\|(U_i - I)x^{n_k}\|^2 \le \|(U_{t_1^k}^{n_k} - I)x^{n_k}\|^2,\\
&\frac{\theta\rho_2}{2R}\|(\mathcal{L}_{\sigma_j}(\{V_j\} - I)x^{n_k}\|^2 \le \|(V_{t_1^k}^{n_k} - I)x^{n_k}\|^2,
\end{aligned} \tag{29}$$

where $R$ is some positive constant, because $\{x^{n_k}\}$ is bounded. Then, from (28) and (29) it follows that

$$\lim_{k\to\infty}\|(U_i - I)x^{n_k}\| = 0,\ \lim_{k\to\infty}\|(\mathcal{L}_{\sigma_j}\{V_j\} - I)x^{n_k}\| = 0. \tag{30}$$

The latter limit and (12) yield $\lim_{k\to\infty}\|(V_j - I)A_j x^{n_k}\| = 0$. Since $A_j$ is a bounded linear operator, the property of $\{x^{n_k}\}$ and the demiclosedness property of $U_i$, $V_j$ we have $\tilde{p} \in \mathrm{Fix}\,(U_i)$ and $A_j\tilde{p} \in \mathrm{Fix}\,(V_j)$ for each $i \in \mathcal{N}$ and $j \in \mathcal{M}$. It means that $\tilde{p} \in \Gamma$. Similarly, any weak cluster point of $\{x^k\}$ belongs to $\Gamma$. Therefore,

$$\limsup_{k\to\infty}\langle Fp_*, p_* - x^k\rangle \le 0.$$

By using this limit, the following one,

$$\|x^{k+1} - p_*\|^2 \le (1 - \theta_k\alpha_k\beta_k)\|x^k - p_*\|^2 + 2\alpha_k\big(\langle Fp_*, p_* - x^k\rangle + \alpha_k\|Fp_*\|M_1\big)$$

obtained by Lemma 3.3 with $p = p_*$, and Lemma **??** with properties of $\theta_k\alpha_k\beta_k$, we get $\lim_{k\to\infty}\|x^k - p_*\| = 0$.

Now, we consider the case when $U_t^k$ and $V_t^k$ are given by (16). By the similar argument as in the proof of Lemma 3.2 we have

$$\begin{aligned}
\lim_{k\to\infty}\big[&\|(U_iU_{i_{l-1}^k(t_1^k)}...U_{i_1^k(t_1^k)} - U_{i_{l-1}^k(t_1^k)}...U_{i_{1-2}^k(t_1^k)})x^k\|^2 +\\
&\|(U_{i_{l-2}^k(t_1^k)}...U_{i_1^k(t_1^k)} - U_{i_{l-3}^k(t_1^k)}...U_{i_1^k(t_1^k)})x^k\|^2 + ... +\\
&\|(U_{i_2^k(t_1^k)}U_{i_1^k(t_1^k)} - U_{i_1^k(t_1^k)})x^k\|^2 + \|(U_{i_1^k(t_1^k)} - I)x^k\|^2\big] = 0
\end{aligned}$$

and

$$\lim_{k\to\infty}\big[\|(\mathcal{L}_{\sigma_j}-\mathcal{L}^{\sigma_{j^k_{l'-1}(t^k_2)}})x^k\|^2+\|(\mathcal{L}^{\sigma_{j^k_{l'-1}(t^k_2)}}-\mathcal{L}^{\sigma_{j^k_{l'-2}(t^k_2)}})x^k\|^2+...$$
$$+\|(\mathcal{L}^{\sigma_{j^k_2(t^k_2)}}-\mathcal{L}^{\sigma_{j^k_1(t^k_2)}})x^k\|^2+\|(\mathcal{L}^{\sigma_{j^k_1(t^k_2)}}-I)x^k\|^2\big]=0.$$

Therefore, (30) is satisfied. Next, by the same argument as above we obtain the same result as for $U^k_t, V^k_t$ defined by (14). Clearly, when $U^k_t$ and $V^k_t$ are given by (19) or (20), we also obtain the same result.
*Case 2.* There exists a subsequence $\{l_k\}\subset\{k\}$ such that

$$\|x^{l_k}-p_*\|\le\|x^{l_k+1}-p_*\|\ \forall k\ge\tilde{k}.$$

Then, by Lemma **??**, there exists a non-decreasing sequence $\{n_k\}\subseteq\{k\}$ such that $n_k\to\infty$,

$$\|x^{n_k}-p_*\|\le\|x^{n_k+1}-p_*\|,\quad \|x^k-p_*\|\le\|x^{n_k+1}-p_*\| \tag{31}$$

for each $k\ge\tilde{k}$. So, from (24) with $k$ replaced by $n_k$ and the first inequality in (31), it follows that

$$\|x^{n_k}-p_*\|^2\le\frac{4}{\theta\eta}\big[\langle Fp_*,p_*-x^{n_k}\rangle+\alpha_{n_k}\|Fp\|M_1\big] \tag{32}$$

and $d^{n_k}\le d_{n_k}\alpha_{n_k}$ for sufficiently $n_k\ge\tilde{k}$. Hence, $\lim_{k\to\infty}d^{n_k}=0$ by the properties of $d_{n_k}$ and $\alpha_{n_k}$. By the same argument as in the proof of the *Case 1*, we obtain (30). Evidently, there exists a subsequence $\{x^{m_k}\}$ of $\{x^{n_k}\}$ such that $\{x^{m_k}\}$ converges weakly to a point $\underline{p}\in\mathcal{H}_0$ as $k\to\infty$. With each $i\in\mathcal{N}$ and $j\in\mathcal{M}$, for each $k\ge0$ let $t^k_1,t^k_2,\mathcal{N}^k_{t^k_1}$, and $\mathcal{M}^k_{t^k_2}$ be defined as the above. Next, again, by using the same argument as in the proof for the *Case 1* above, $\underline{p}\in\Gamma$ and any weak cluster point of $\{x^{m_k}\}$ belongs to $\Gamma$. Thus,

$$\limsup_{k\to\infty}\langle Fp_*,p_*-x^{n_k}\rangle\le0.$$

This, (32), and the property of $\alpha_k$ deduce $\lim_{k\to\infty}\|x^{n_k}-p_*\|=0$. Further, from Lemma **??** with $k$ and $p$ replaced, respectively, by $n_k$ and $p_*$, we get

$$\|x^{n_k+1}-p_*\|^2\le\|x^{n_k}-p_*\|^2+2\alpha_{n_k}\|Fp_*\|\big(r'+\alpha_{n_k}M_1\big).$$

This, the last limit, and the property of $\alpha_{n_k}$ yield $\lim_{k\to\infty}\|x^{n_k+1}-p_*\|^2=0$, from which and the second inequality in (31) we have $\lim_{k\to\infty}\|x^k-p_*\|=0$. The proof is completed.

**Remarks**

1. If we take $F=I$, then $F$ is $\eta$-strongly monotone and $l$-Lipschitz continuous mapping on $\mathcal{H}_0$ with $\eta\le1$ and $l\ge1$. Then, we obtain

$$x^{k+1}=(1-\xi_k)(1-t_k)x^k+\xi_kx^{k,s}, \tag{33}$$

since $\mu_k = 1$ for all $k \geq 0$. Method (33) is an improved modification of Krasnoselski-Mann [19, 21] iterative method for the considered problem.
2. Let $\alpha$ be any positive real number in $(0, 1)$. Take $F = (1-\alpha)I - (1-\alpha)u$, where $u$ is a fixed point in $\mathcal{H}_0$. Clearly, $F$ is $\eta$-strongly monotone and $l$-Lipschitz continuous operator on $\mathcal{H}_0$ with $l \geq \eta > 0$. Then, from (11) we deduce that

$$x^{k+1} = (1-\theta_k)[(1-t'_k)x^k + t'_k u] + \theta_k x^{k,s},$$

where $t'_k = (1-\alpha)t_k$. This method is a combination of the Krasnoselski-Mann iterative method with the Halpern one [18].
3. The results above have still value, if (12) is replaced by either

$$x^{k+1} = (1-\theta_k)x^k + \theta_k(I - t_k\mu_k F)x^{k,s} \tag{34}$$

with $\theta_k \in [\theta, 1]$ or

$$\begin{aligned} x^{k+1} &= (1-\theta_k)(I - t_k\mu_k F)x^{k,s} + \theta_k x^{k,s} \\ &= (I - \tilde{\theta}_k t_k \mu_k F)x^{k,s},\ \tilde{\theta}_k = 1-\theta_k, \end{aligned}$$

with $\theta_k \in [\theta, 1-\theta]$, where

$$\mu_k = \frac{\langle Fx^{k,s} - Fx^{k-1,s}, x^{k,s} - x^{k-1,s}\rangle}{\|Fx^{k.s} - Fx^{k-1,s}\|^2},$$

if $x^{k,s} \neq x^{k-1,s}$, $\mu_{k-1}$, otherwise, and $x^{-1,s} = x^{-1}$ and $x^{0,s} = x^0$. Taking $\theta_k = 1$ and $F = I$, (34) becomes a simple iterative method $x^{k+1} = (1-t_k)x^{k,s}$.
4. The approach in this paper can be extended to a generation of (1), that is to find a point

$$p \in C := \cap_{i\in\mathcal{N}}\text{Fix}(U_i),\ A_j p \in \cap_{l\in\mathcal{M}_{l,j}}\text{Fix}(V_{l,j}), \forall j \in \mathcal{M},$$

where $V_{l,j}$ is $\rho_{l,j}$-strongly quasi-nonexpansive operator on $\mathcal{H}_j$ with $\rho_{l,j} > 0$ for every $l \in \mathcal{M}_{l,j} = \{1, ..., q_{l,j}\}$ with integers $q_{l,j} \geq 1$.
5. Some similar split problems with iterative methods have been recently considered in [7, 6, 5].

# 4 An application to the MSSFP given by level sets and computational experiments

Let $\varphi_i$ and $\psi_j$ be boundedly Lipschitz continuous and convex functions on real Hilbert spaces $\mathcal{H}_0$ and $\mathcal{H}_j$, respectively, for $i \in \mathcal{N}$ and $j \in \mathcal{M}$. We consider the MSSFP: Find a point

$$\begin{aligned} &p \in C = \{p \in \mathcal{H}_0 : \varphi_i(p) \leq 0\}\ \forall i \in \mathcal{N}, \\ &A_j p \in \{v \in \mathcal{H}_j : \psi_j(v) \leq 0\}\ \forall j \in \mathcal{M} \end{aligned} \tag{35}$$

where $A_j$ is a bounded linear operator from $\mathcal{H}_0$ into $\mathcal{H}_j$, for all $j \in \mathcal{M}$. The solution set of (35) is denoted by $\Gamma$.

Let $\partial f(x) = \{g \in \mathcal{H} : f(y) - f(x) \geq \langle g, y - x\rangle$, for all $y \in \mathcal{H}\}$, the subdifferential of a continuous convex function $f$ on a Hilbert space $\mathcal{H}$. Since $f$ is continuous, the set $\partial f(x) \neq \emptyset$ (see, [3]). For an $\eta(x) \in \partial f(x)$, the operator $P_f$, so-called subgradient projection relative to $f$, is defined by

$$P_f x := x - \frac{[f(x)]_+}{\|\eta(x)\|^2}\eta(x)$$

if $\eta(x) \neq 0$; $x$, otherwise. where $[t]_+ = \max\{0, t\}$, has the properties: $\text{Fix}\,(P_f) = \{p \in \mathcal{H} : f(p) \leq 0\}$ and $P_f$ is 1-strongly quasi-nonexpansive, which satisfies the demiclosedness principle (see, [10] Theorem 4.2.7 and Corollary 4.2.6). Thus, $p_*$ is a solution of (35) if and only if it belongs to $\Gamma$ and solves (1) with $U_i = P_{\varphi_i}$ and $V_j = P_{\psi_j}$.

For computations, we take $\mathcal{H}_0 = \mathbf{E}^4, \mathcal{H}_j = \mathbf{E}^{j+1}$, Euclidean spaces, with $j \in \mathcal{M} = \{1, 2, 3, 4\}$.

$$\begin{aligned}
&\varphi_i(x) = \|(I - P_{\tilde{C}_i})x\|^2/2, \tilde{C}_i = \{x \in \mathbf{E}^4 : \|x - a^i\|^2 \leq 1\}, a^i \in \mathbf{E}^4,\\
&\psi_j(y) = \|(I - P_{\tilde{Q}_j})y\|^2/2, \tilde{Q}_j = \{y \in \mathbf{E}^j : \|y - b^j\|^2 \leq 1\}, b^j \in \mathbf{E}^j,\\
&a^i = (1 - i/10, 0, 0, 0),\ b^j = (1, 0, ..., 0),
\end{aligned}$$

for all $i \in \mathcal{N} = \{1, ..., 6\}$ and for all $j \in \mathcal{M}$ with a matrix $A_1$, having two rows $r_1 = (1, 2, 1, 2)$ and $r_2 = (2, 4, 2, 4)$, a matrix $A_2$, having three rows $r_1 = (1, 2, 0, 0)$, $r_2 = (2, 4, 0, 0)$ and $r_3 = (0, 0, 1, 0)$, a matrix $A_3$, having four rows $r_1 = (2, 1, 0, 0), r_2 = (0, 2, 1, 0)$, $r_3 = (0, 0, 2, 1), r_4 = (0, 0, 4, 2)$, and a matrix $A_4$, having fine rows $r_1 = (2, 1, 0, 0), r_2 = (0, 2, 1, 0)$, $r_3 = (0, 0, 2, 1), r_4 = (0, 0, 4, 2)$, $r_5 = (0, 0, 0, 1)$. It is easy to see that $\cap_{i\in L}\tilde{C}_i \neq \emptyset$ and the considered example has the unique minimum norm solution $p_* = (0, 0, 0, 0)$.

Clearly, $\partial\varphi_i(x) = \varphi_i'(x) = x - P_{\tilde{C}_i}x$, that has the form (see, [10]): $P_{\tilde{C}_i}x = a^i + (1/\|x - a^i\|)(x - a^i)$ if $\|x - a^i\| > 1$ and $x$ otherwise. Then, $P_{\varphi_i}x = (x + P_{\tilde{C}_i}x)/2$ if $x \notin \tilde{C}_i$; otherwise, $P_{\varphi_i}x = x$ and $P_{\psi_j}$ is similarly defined in $\mathbf{E}^{j+1}$. For computations, by taking $s = 2$, $\mathcal{N} = \mathcal{N}_1^k \cup \mathcal{N}_2^k$ and $\mathcal{M} = \mathcal{M}_1^k \cup \mathcal{M}_2^k$ with $\mathcal{N}_1^k = \{1, 2, 3\}$, $\mathcal{N}_2^k = \{4, 5, 6\}$, $\mathcal{M}_1^k = \{1, 2\}$, $\mathcal{M}_2^k = \{3, 4\}$, $\sigma_{j^k(t)}(x) = \tau_{j^k(t)}(x)$, and the starting point $x^0 = (3, 2, 5, 4)$, and using method (33), we have the following scheme,

$$\begin{aligned}
&x^{k,t} = U_t^k y^{k,t} y^{k,t} = V_t^k x^{k,t-1},\ t = 1, 2,\\
&x^{k+1} = (1 - \xi_k)(1 - t_k)x^k + \xi_k x^{k,2}.
\end{aligned} \tag{36}$$

where $\tau_{j^k(t)}(x)$ is defined in the block-iterative scheme. Computational results by (36) with (14), (15), (19), and (20), $t_k = 1/(k + 1)$, $\lambda_{i^k(t)}^k = 1/3$, and $\xi_k = \theta_{j^k(t)}^k = 1/2$ are given in the following tables.

Table 1: Result by (36) and (14).

| $k$ | $x_1^{k+1}$ | $x_2^{k+1}$ | $x_3^{k+1}$ | $x_4^{k+1}$ |
|---|---|---|---|---|
| 10 | 0.0088308449 | 0.0152433682 | 0.0454595158 | 0.0246655582 |
| 20 | 0.0004772087 | 0.0008681698 | 0.0027052801 | 0.0014691992 |
| 30 | 0.0000327203 | 0.0000595935 | 0.0001860741 | 0.0001010072 |
| 40 | 0.0000024101 | 0.0000043897 | 0.0000137075 | 0.0000074410 |
| 50 | 0.0000001846 | 0.0000003361 | 0.0000010497 | 0.0000005698 |

Table 2: Result by (36) and (16).

| $k$ | $x_1^{k+1}$ | $x_2^{k+1}$ | $x_3^{k+1}$ | $x_4^{k+1}$ |
|---|---|---|---|---|
| 10 | 0.0100743460 | 0.0151863428 | 0.0230477441 | 0.0110144905 |
| 20 | 0.0004430946 | 0.0006780287 | 0.0010239829 | 0.0004835695 |
| 30 | 0.0000231565 | 0.0000354934 | 0.0000535205 | 0.0000252499 |
| 40 | 0.0000012387 | 0.0000019907 | 0.0000030017 | 0.0000014155 |
| 50 | 0.0000000758 | 0.0000001161 | 0.0000001751 | 0.0000000826 |

Table 3: Result by (36) and (19).

| $k$ | $x_1^{k+1}$ | $x_2^{k+1}$ | $x_3^{k+1}$ | $x_4^{k+1}$ |
|---|---|---|---|---|
| 10 | 0.0044369738 | 0.0087323594 | 0.0387153882 | 0.0248808418 |
| 20 | 0.0002209196 | 0.0004826718 | 0.0022353282 | 0.0014344862 |
| 30 | 0.0000147412 | 0.0000328535 | 0.0001495178 | 0.0009595348 |
| 40 | 0.0000010566 | 0.0000023118 | 0.0000107169 | 0.0000068776 |
| 50 | 0.0000000787 | 0.0000001722 | 0.0000007986 | 0.0000005125 |

Table 4: Result by (36) and (20).

| $k$ | $x_1^{k+1}$ | $x_2^{k+1}$ | $x_3^{k+1}$ | $x_4^{k+1}$ |
|---|---|---|---|---|
| 10 | 0.0091396752 | 0.0147419320 | 0.0304096230 | 0.0155142279 |
| 20 | 0.0003689121 | 0.0006096060 | 0.0012723778 | 0.0064546818 |
| 30 | 0.0000180284 | 0.0000298010 | 0.0000622267 | 0.0000315668 |
| 40 | 0.0000009468 | 0.0000015352 | 0.0000032682 | 0.0000016579 |
| 50 | 0.0000000518 | 0.0000000856 | 0.0000001787 | 0.0000000906 |

The tables of numerical results show the effectiveness of the introduced iterative schemes.

## 5 Conclusion

In this paper, using the extrapolated Landweber-type operators, we proposed new strongly convergent iterative schemes for solving the the split feasibility problem with multiple output sets, governed by the fixed point sets of demiclosed strongly quasi-nonexpansive operators on infinite-dimensional Hilbert spaces. The strong convergence has been proved without the closedness property the range of $A_j$ as well as the bounded regular one of the family of fixed point sets of the operators, assumed recently in the literature for the similar problems. An application to the MSSFP with multiple output level sets was given with numerical experiments.

## Disclosure statement

No potential conflict of interest was supported by the authors.

## References

[1] Abass H.A. and Aphane M., *An algorithmic approach to solving split common fixed point problems for families of demicontractive operators in Hilbert spaces*, Boll. Unione Mat. Ital., doi:10.1007/s40574-023-00391-7.

[2] Aharoni R. and Censor Y., *Block-iterative projection methods for parallel computation of solutions to convex feasibility problems*, Linear Algebra Appl., 120 (1989), 165–175.

[3] Bauschke H.H. and Combettes P.L., *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, Springer, New York, Dordrecht, Heidelberg, London, 2011.

[4] Brooke M., Censor Y. and Gibali A., *Dynamic string-averaging CQ-methods for the split feasibility problem with percentage violation constraints arising in radiation therapy treatment planning*, Int. Trans. Oper. Res., 30(1) (2023), 181–215.

[5] Buong Ng. and Anh Ng.Th.Q., *Extrapolated simultaneous block-iterative cutter methods and applications*, Math. Methods Appl. Sci., 46(13) (2023), 14229–14242.

[6] Buong Ng. and Nguyen Ng.D., *Convergence of extrapolated dynamic string-averaging cutter methods and applications*, East Asian J. Appl. Math., 13(2) (2023), 257–275.

[7] Buong Ng., *Steepest-descent block-iterative methods for a finite family of quasi-nonexpansive mappings*, J. Ind. Manag. Optim., 18(5) (2022), 3735–3748.

[8] Cegielski A. and Al-Musallam F., *Strong convergence of a hybrid steepest descent method for the split common fixed point problem*, Optimization, 65(7) (2016), 1463–1476.

[9] Cegielski A., Gibali A., Reich S. and Zalas R., *Outer approximation methods for solving variational inequalities defined over the solution set of a split convex feasibility problem*, Numer. Funct. Anal. Optim., 41(9) (2020), 1089–1108.

[10] Cegielski A., *Iterative Methods for Fixed Point Problems in Hilbert Spaces*, Springer, Heidelberg, 2012. (Lecture Notes in Math., Vol. 2057)

[11] Cegielski A., *General method for solving the split common fixed point problem*, J. Optim. Theory Appl., 165(2) (2015), 385–404.

[12] Cegielski A., *Landweber-type operator and its properties*, Contemp. Math., 658 (2016), 139–148.

[13] Cegielski A., Reich S. and Zalas R., *Regular sequences of quasi-nonexpansive operators and their applications*, SIAM J. Optim., 28(2) (2018), 1508–1532.

[14] Cegielski A., Reich S. and Zalas R., *Weak, strong and linear convergence of the CQ-method via the regularity of Landweber operators*, Optimization, 69(3) (2020), 605–636.

[15] Cegielski A. and Zalas R., *Properties of a class of approximately shrinking operators and their applications*, Fixed Point Theory, 15(2) (2014), 399–426.

[16] Censor Y. and Segal A., *The split common fixed point problem for directed operators*, J. Convex Anal., 16(2) (2009), 587–600.

[17] Gibali A., Reich S. and Zalas R., *Outer approximation method for solving variational inequalities in Hilbert spaces*, Optimization, 66(3) (2017), 417–437.

[18] Halpern B., *Fixed points of nonexpanding maps*, Bull. Amer. Math. Soc., 73 (1967), 957–961.

[19] Krasnosel'skii M.A., *Two remarks on the method of successive approximations*, Uspekhi Mat. Nauk, 10 (1955), 123–127.

[20] Maingé F.E., *Strong convergence of projected subgradient methods for nonsmooth and nonstrictly convex minimization*, Set-Valued Var. Anal., 16(7) (2008), 899–912.

[21] Mann W.R., *Mean value methods in iteration*, Proc. Amer. Math. Soc., 4 (1953), 506–510.

[22] Mirzapour M., Cegielski A. and Elfving T., *Convergence and semiconvergence of a class of constrained block-iterative methods*, Numer. Funct. Anal. Optim., 42(14) (2021), 1718–1746.

[23] Moudafi A., *Viscosity approximation methods for fixed-point problems*, J. Math. Anal. Appl., 241(1) (2000), 46–55.

[24] Moudafi A., *The split fixed point problem for demicontractive mappings*, Inverse Problems, 26(5) (2010), 055007.

[25] Moudafi A., *A note on the split common fixed-point problem for quasi-nonexpansive operators*, Nonlinear Anal., 74(12) (2011), 4083–4087.

[26] Reich S. and Zalas A., *A modular string-averaging procedure for solving the common fixed point problems for quasi-nonexpansive mappings in Hilbert spaces*, Numer. Algorithms, 72(2) (2016), 297–323.

[27] Tang Y.C. and Liu L.W., *Several iterative algorithms for solving the split common fixed point problem of directed operators with applications*, Optimization, 65(1) (2016), 53–65.

[28] Wang F. and Xu H.K., *Cyclic algorithms for split feasibility problems in Hilbert spaces*, Nonlinear Anal., 74(12) (2011), 4105–4111.

[29] Wang F., *The split feasibility problem with multiple output sets for demicontractive mappings*, J. Optim. Theory Appl., 195(3) (2022), 837–853.

[30] Xu H.K., *An iterative approach to quadratic optimization*, J. Optim. Theory Appl., 116(3) (2003), 659–678.

[31] Yamada I., *The hybrid steepest descent method for the variational inequality problem over the intersection of fixed point sets of nonexpansive mappings*, in *Inherently Parallel Algorithms Feasibility Optim. Appl.* (Butnariu D., Censor Y. and Reich S., Eds.), North-Holland, Amsterdam, (2001), 473–504.