

ON THE PETERSON HIT PROBLEM OF FIVE VARIABLES AND ITS APPLICATIONS TO THE FIFTH SINGER TRANSFER

Nguyen Sum

*Department of Mathematics, Quy Nhon University
170 An Duong Vuong, Quy Nhon, Bình Định, Viet Nam.
e-mail: nguyensum@qnu.edu.vn*

Abstract

We study the Peterson hit problem of finding a minimal set of generators for the polynomial algebra $P_k := \mathbb{F}_2[x_1, x_2, \dots, x_k]$ as a module over the mod-2 Steenrod algebra, \mathcal{A} . In this paper, we explicitly determine a minimal set of \mathcal{A} -generators with $k = 5$ in degree 15. Using this results we show that the fifth Singer transfer is an isomorphism in this degree.

1 Introduction and statement of results

Let V_k be an elementary abelian 2-group of rank k . Denote by BV_k the classifying space of V_k . It may be thought of as the product of k copies of the real projective space $\mathbb{R}P^\infty$. Then

$$P_k := H^*(BV_k) \cong \mathbb{F}_2[x_1, x_2, \dots, x_k],$$

a polynomial algebra on k generators x_1, x_2, \dots, x_k , each of degree 1. Here the cohomology is taken with coefficients in the prime field \mathbb{F}_2 of two elements.

Being the cohomology of a space, P_k is a module over the mod 2 Steenrod algebra \mathcal{A} . The action of \mathcal{A} on P_k can explicitly be given by the formula

$$Sq^i(x_j) = \begin{cases} x_j, & i = 0, \\ x_j^2, & i = 1, \\ 0, & \text{otherwise,} \end{cases}$$

Key words: Steenrod squares, Hit problem, Singer transfer.

AMS Mathematics Subject Classification: Primary 55S10; 55S05, 55T15.

and subject to the Cartan formula $Sq^n(fg) = \sum_{i=0}^n Sq^i(f)Sq^{n-i}(g)$, for $f, g \in P_k$ (see Steenrod-Epstein [22]).

A polynomial f in P_k is called *hit* if it can be written as a finite sum $f = \sum_{i \geq 0} Sq^i(f_i)$ for some polynomials f_i . That means f belongs to \mathcal{A}^+P_k , where \mathcal{A}^+ denotes the augmentation ideal in \mathcal{A} . We are interested in the *hit problem*, set up by F. Peterson, of finding a minimal set of generators for the polynomial algebra P_k as a module over the Steenrod algebra. In other words, we want to find a basis of the \mathbb{F}_2 -vector space $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k := QP_k$.

Let $GL_k = GL_k(\mathbb{F}_2)$ be the general linear group over the field \mathbb{F}_2 . This group acts naturally on P_k by matrix substitution. Since the two actions of \mathcal{A} and GL_k upon P_k commute with each other, there is an action of GL_k on QP_k . The subspace of degree n homogeneous polynomials $(P_k)_n$ and its quotient $(QP_k)_n$ are GL_k -subspaces of the spaces P_k and QP_k respectively.

The hit problem was first studied by Peterson [15], Wood [26], Singer [20], and Priddy [16], who showed its relationship to several classical problems respectively in cobordism theory, modular representation theory, Adams spectral sequence for the stable homotopy of spheres, and stable homotopy type of classifying spaces of finite groups. The tensor product QP_k was explicitly calculated by Peterson [15] for $k = 1, 2$, by Kameko [10] for $k = 3$, and recently by us [23] for $k = 4$.

Many authors was then investigated the hit problem. (See Boardman [1], Bruner-Hà-Hung [2], Crabb-Hubbuck [5], Hà [6], Hung [7, 8], Kameko [10, 11], Nam [13, 14], Repka-Selick [18], Singer [21], Silverman [19], Wood [26, 27] and others.)

One of our main tools for studying the hit problem is the so-called Kameko squaring operation

$$Sq^0 : \mathbb{F}_2 \otimes_{GL_k} PH_*(BV_k) \rightarrow \mathbb{F}_2 \otimes_{GL_k} PH_*(BV_k).$$

Here $H_*(BV_k)$ is homology with \mathbb{F}_2 coefficients, and $PH_*(BV_k)$ denotes the primitive subspace consisting of all elements in the space $H_*(BV_k)$, which are annihilated by every positive-degree operation in the mod 2 Steenrod algebra; therefore, $\mathbb{F}_2 \otimes_{GL_k} PH_*(BV_k)$ is dual to $QP_k^{GL_k}$. The dual of the Kameko squaring is the homomorphism $Sq_*^0 : QP_k^{GL_k} \rightarrow QP_k^{GL_k}$. This homomorphism is given by the following GL_k -homomorphism $\widetilde{Sq}_*^0 : QP_k \rightarrow QP_k$. The latter is given by the \mathbb{F}_2 -linear map, also denoted by $\widetilde{Sq}_*^0 : P_k \rightarrow P_k$, given by

$$\widetilde{Sq}_*^0(x) = \begin{cases} y, & \text{if } x = x_1x_2 \dots x_ky^2, \\ 0, & \text{otherwise,} \end{cases}$$

for any monomial $x \in P_k$. Note that \widetilde{Sq}_*^0 is not an \mathcal{A} -homomorphism. However, $\widetilde{Sq}_*^0 Sq^{2t} = Sq^t \widetilde{Sq}_*^0$ and $\widetilde{Sq}_*^0 Sq^{2t+1} = 0$, for any nonnegative integer t .

The Kameko squaring operation commutes with the classical squaring operation on the cohomology of the Steenrod algebra through the Singer transfer

$$\mathrm{Tr}_k : \mathbb{F}_2 \otimes_{GL_k} PH_d(BV_k) \rightarrow \mathrm{Ext}_{\mathcal{A}}^{k,k+d}(\mathbb{F}_2, \mathbb{F}_2).$$

Boardman [1] used this fact to show that Tr_3 is an isomorphism. Bruner-Hà-Hung [2] applied it to prove that Tr_4 does not detect any element in the usual family $\{g_i\}_{i>0}$ of $\mathrm{Ext}_{\mathcal{A}}^4(\mathbb{F}_2, \mathbb{F}_2)$. Recently, Hung and his collaborators have completely determined the image of the fourth Singer transfer Tr_4 (in [2], [8], [6], [14], [9]). Singer showed in [20] that Tr_5 is not an epimorphism in degree 9. In [17], Quỳnh proved that Tr_5 is also not an epimorphism in degree 11. The Singer transfer was also investigated by Chon-Hà [3, 4].

In this paper, we explicitly determine all the admissible monomials (see Section 2) of P_5 in degree 15. Using this results, we prove that the fifth Singer transfer is an isomorphism in this degree. We have

Theorem 1.1. *There exist exactly 432 admissible monomials of degree 15 in P_5 . Consequently $\dim(QP_5)_{15} = 432$.*

By using Theorem 1.1, we compute $(QP_5)_{15}^{GL_5}$.

Theorem 1.2. *$(QP_5)_{15}^{GL_5}$ is an \mathbb{F}_2 -vector space of dimension 2 with a basis consisting of the 2 classes represented by the following polynomials:*

$$\begin{aligned} p &= x_1^{15} + x_2^{15} + x_3^{15} + x_4^{15} + x_5^{15} + x_1x_2^{14} + x_1x_3^{14} + x_1x_4^{14} + x_1x_5^{14} + x_2x_3^{14} \\ &\quad + x_2x_4^{14} + x_2x_5^{14} + x_3x_4^{14} + x_3x_5^{14} + x_4x_5^{14} + x_1x_2^2x_3^{12} + x_1x_2^2x_4^{12} + x_1x_2^2x_5^{12} \\ &\quad + x_1x_3^2x_4^{12} + x_1x_3^2x_5^{12} + x_1x_4^2x_5^{12} + x_2x_3^2x_4^{12} + x_2x_3^2x_5^{12} + x_2x_4^2x_5^{12} + x_3x_4^2x_5^{12} \\ &\quad + x_1x_2^2x_3^4x_4^8 + x_1x_2^2x_3^4x_5^8 + x_1x_2^2x_4^4x_5^8 + x_1x_3^2x_4^4x_5^8 + x_2x_3^2x_4^4x_5^8 + x_1x_2^2x_3^4x_4^4x_5^4, \\ q &= x_1x_2x_3x_4^6x_5^6 + x_1x_2x_3^6x_4x_5^6 + x_1x_2x_3^6x_4^6x_5 + x_1x_2^6x_3x_4x_5^6 + x_1x_2^6x_3x_4^6x_5 \\ &\quad + x_1x_2^3x_3^6x_4x_5^4 + x_1x_2^3x_3^6x_4^4x_5 + x_1x_2^6x_3^3x_4x_5^4 + x_1x_2^6x_3^3x_4^4x_5 + x_1^3x_2x_3x_4^6x_5^6 \\ &\quad + x_1^3x_2x_3x_4^6x_5^4 + x_1^3x_2x_3^4x_4x_5^6 + x_1^3x_2x_3^4x_4^6x_5 + x_1^3x_2^4x_3x_4x_5^6 + x_1^3x_2^4x_3x_4^6x_5 \\ &\quad + x_1x_2^3x_3^3x_4^4x_5^4 + x_1x_2^3x_3^3x_4^4x_5^4 + x_1^3x_2^3x_3x_4^4x_5^4 + x_1^3x_2^3x_3^4x_4x_5^4 + x_1^3x_2^3x_3^4x_4^4x_5 \\ &\quad + x_1^3x_2^4x_3^3x_4x_5^4 + x_1^3x_2^4x_3^3x_4^4x_5. \end{aligned}$$

Using Theorem 1.2, we prove the following which was proved in Hung [8] by using computer computation.

Theorem 1.3 (Hung [8]). *The fifth Singer transfer*

$$\mathrm{Tr}_5 : \mathbb{F}_2 \otimes_{GL_5} PH_{15}(BV_5) \rightarrow \mathrm{Ext}_{\mathcal{A}}^{5,20}(\mathbb{F}_2, \mathbb{F}_2)$$

is an isomorphism.

This paper is organized as follows. In Section 2, we recall some needed information on the admissible monomials in P_k and Singer criterion on the hit monomials. We prove Theorem 1.1 in Section 3 by explicitly determine all the admissible monomials of degree 15. Theorems 1.2 and 1.3 will be proved in Sections 4.

2 Preliminaries

In this section, we recall some results in Kameko [10] and Singer [21] which will be used in the next sections.

Notation 2.1. Let $\alpha_i(a)$ denote the i -th coefficient in dyadic expansion of a nonnegative integer a . That means $a = \alpha_0(a)2^0 + \alpha_1(a)2^1 + \alpha_2(a)2^2 + \dots$, for $\alpha_i(a) = 0, 1$ and $i \geq 0$.

Let $x = x_1^{a_1} x_2^{a_2} \dots x_k^{a_k} \in P_k$. Set $I_i(x) = \{j \in \mathbb{N}_k : \alpha_i(a_j) = 0\}$, for $i \geq 0$. Then we have

$$x = \prod_{i \geq 0} X_{I_i(x)}^{2^i}.$$

For a polynomial f in P_k , we denote by $[f]$ the class in $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k$ represented by f . For a subset $S \subset P_k$, we denote

$$[S] = \{[f] : f \in S\} \subset QP_k.$$

Definition 2.2. For a monomial $x = x_1^{a_1} x_2^{a_2} \dots x_k^{a_k} \in P_k$, we define two sequences associated with x by

$$\begin{aligned} \omega(x) &= (\omega_1(x), \omega_2(x), \dots, \omega_i(x), \dots), \\ \sigma(x) &= (a_1, a_2, \dots, a_k), \end{aligned}$$

where $\omega_i(x) = \sum_{1 \leq j \leq k} \alpha_{i-1}(a_j) = \deg X_{I_{i-1}(x)}$, $i \geq 1$.

The sequence $\omega(x)$ is called the weight vector of x (see Wood [27]). The weight vectors and the sigma vectors can be ordered by the left lexicographical order.

Let $\omega = (\omega_1, \omega_2, \dots, \omega_i, \dots)$ be a sequence of nonnegative integers such that $\omega_i = 0$ for $i \gg 0$. Define $\deg \omega = \sum_{i \geq 0} 2^{i-1} \omega_i$. Denote by $P_k(\omega)$ the subspace of P_k spanned by all monomials y such that $\deg y = \deg \omega$, $\omega(y) \leq \omega$ and $P_k^-(\omega)$ the subspace of P_k spanned by all monomials $y \in P_k(\omega)$ such that $\omega(y) < \omega$. Denote by \mathcal{A}_s^+ the subspace of \mathcal{A} spanned by all Sq^j with $1 \leq j < 2^s$. Define

$$QP_k(\omega) = P_k(\omega) / ((\mathcal{A}^+ P_k \cap P_k(\omega)) + P_k^-(\omega)).$$

Then we have

$$(QP_k)_n = \oplus_{\deg \omega = n} QP_k(\omega).$$

Definition 2.3. Let x be a monomial and f, g two homogeneous polynomials of the same degree in P_k . We define $f \equiv g$ if and only if $f - g \in \mathcal{A}^+ P_k$. If $f \equiv 0$ then f is called hit.

We recall some relations on the action of the Steenrod squares on P_k .

Proposition 2.4. Let f be a homogeneous polynomial in P_k .

- i) If $i > \deg f$ then $Sq^i(f) = 0$. If $i = \deg f$ then $Sq^i(f) = f^2$.
- ii) If i is not divisible by 2^s then $Sq^i(f^{2^s}) = 0$ while $Sq^{r2^s}(f^{2^s}) = (Sq^r(f))^{2^s}$.

Definition 2.5. Let x, y be monomials of the same degree in P_k . We say that $x < y$ if and only if one of the following holds

- i) $\omega(x) < \omega(y)$;
- ii) $\omega(x) = \omega(y)$ and $\sigma(x) < \sigma(y)$.

Definition 2.6. A monomial x is said to be inadmissible if there exist monomials y_1, y_2, \dots, y_t such that $y_j < x$ for $j = 1, 2, \dots, t$ and $x \equiv y_1 + y_2 + \dots + y_t$.

A monomial x is said to be admissible if it is not inadmissible.

Obviously, the set of all the admissible monomials of degree n in P_k is a minimal set of \mathcal{A} -generators for P_k in degree n .

The following theorem is a modification of a result in [10].

Theorem 2.7 (Kameko [10], Sum [24]). Let x, w be monomials in P_k such that $\omega_i(x) = 0$ for $i > r > 0$. If w is inadmissible, then xw^{2^r} is also inadmissible.

Proposition 2.8 ([24]). Let x be an admissible monomial in P_k . Then we have

- i) If there is an index i_0 such that $\omega_{i_0}(x) = 0$, then $\omega_i(x) = 0$ for all $i > i_0$.
- ii) If there is an index i_0 such that $\omega_{i_0}(x) < k$, then $\omega_i(x) < k$ for all $i > i_0$.

Now, we recall a result of Singer [21] on the hit monomials in P_k .

Definition 2.9. A monomial $z = x_1^{b_1} x_2^{b_2} \dots x_k^{b_k}$ is called a spike if $b_j = 2^{s_j} - 1$ for s_j a nonnegative integer and $j = 1, 2, \dots, k$. If z is a spike with $s_1 > s_2 > \dots > s_{r-1} \geq s_r > 0$ and $s_j = 0$ for $j > r$, then it is called a minimal spike.

The following is a criterion for the hit monomials in P_k .

Theorem 2.10 (Singer [21]). Suppose $x \in P_k$ is a monomial of degree n , where $\mu(n) \leq k$. Let z be the minimal spike of degree n . If $\omega(x) < \omega(z)$ then x is hit.

For latter use, we set

$$P_k^0 = \langle \{x = x_1^{a_1} x_2^{a_2} \dots x_k^{a_k} ; a_1 a_2 \dots a_k = 0\} \rangle,$$

$$P_k^+ = \langle \{x = x_1^{a_1} x_2^{a_2} \dots x_k^{a_k} ; a_1 a_2 \dots a_k > 0\} \rangle.$$

It is easy to see that P_k^0 and P_k^+ are the \mathcal{A} -submodules of P_k . Furthermore, we have the following.

Proposition 2.11. *We have a direct summand decomposition of the \mathbb{F}_2 -vector spaces*

$$QP_k = QP_k^0 \oplus QP_k^+.$$

Here $QP_k^0 = P_k^0/\mathcal{A}^+.P_k^0$ and $QP_k^+ = P_k^+/\mathcal{A}^+.P_k^+$.

For $1 \leq i \leq k$, define the homomorphism $f_i = f_{k;i} : P_{k-1} \rightarrow P_k$ of algebras by substituting

$$f_i(x_j) = \begin{cases} x_j, & \text{if } 1 \leq j < i, \\ x_{j+1}, & \text{if } i \leq j < k. \end{cases}$$

It is easy to see that

Proposition 2.12. *If $B_{k-1}(n)$ is the set of all admissible monomials of degree n in P_{k-1} , then $f(B_{k-1}(n)) := \cup_{1 \leq i \leq k} f_i(B_{k-1}(n))$ is the set of all admissible monomials of degree n in P_k^0 .*

For $1 \leq i \leq k$, define $\varphi_i : QP_k \rightarrow QP_k$, the homomorphism induced by the \mathcal{A} -homomorphism $\overline{\varphi}_i : P_k \rightarrow P_k$, which is determined by $\overline{\varphi}_1(x_1) = x_1 + x_2$, $\overline{\varphi}_1(x_j) = x_j$ for $j > 1$, and $\overline{\varphi}_i(x_i) = x_{i-1}$, $\overline{\varphi}_i(x_{i-1}) = x_i$, $\overline{\varphi}_i(x_j) = x_j$ for $j \neq i, i-1$, $1 < i \leq k$. Note that the general linear group GL_k is generated by $\overline{\varphi}_i$, $0 \leq i \leq k$ and the symmetric group Σ_k is generated by $\overline{\varphi}_i$, $1 < i \leq k$.

For any $I = (i_0, i_1, \dots, i_r)$, $0 < i_0 < i_1 < \dots < i_r \leq k$, $0 \leq r < k$, we define the homomorphism $p_I : P_k \rightarrow P_{k-1}$ of algebras by substituting

$$p_I(x_j) = \begin{cases} x_j, & \text{if } 1 \leq j < i_0, \\ \sum_{1 \leq s \leq r} x_{i_s-1}, & \text{if } j = i_0, \\ x_{j-1}, & \text{if } i_0 < j \leq k. \end{cases}$$

Then p_I is a homomorphism of \mathcal{A} -modules. In particular, for $I = (i)$, we have $p_{(i)}(x_i) = 0$.

3 Proof of Theorem 1.1

In this section, we explicitly determine all the admissible monomials of degree 15.

Consider the Kameko homomorphism $(\widetilde{Sq}_*^0)_5^5 : (QP_5)_{15} \rightarrow (QP_5)_5$. Since this homomorphism is an epimorphism, we have

$$(QP_5)_{15} \cong \text{Ker}(\widetilde{Sq}_*^0)_5^5 \oplus (QP_5)_5 = ((QP_5^0)_{15} \oplus ((QP_5^+)_{15} \cap \text{Ker}(\widetilde{Sq}_*^0)_5^5) \oplus (QP_5)_5.$$

By Proposition 2.12, to compute $(QP_5^0)_{15}$ we need to compute

$$(QP_4)_{15} = (QP_4)_{15}^0 \oplus (QP_4)_{15}^+.$$

Using Kameko's results in [10], we have

$$B_3(15) = \{x_1^{15}, x_2^{15}, x_3^{15}, x_1x_2^{14}, x_1x_3^{14}, x_2x_3^{14}, x_1x_2^2x_3^{12}, \\ x_1x_2^7x_3^7, x_1^7x_2x_3^7, x_1^7x_2^7x_3, x_1^3x_2^5x_3^7, x_1^3x_2^7x_3^5, x_1^7x_2^3x_3^5\}.$$

By a direct computation using Proposition 2.12, we see that $f(B_3(15))$ is the set consisting of 38 admissible monomials in $(P_5^0)_{15}$.

Lemma 3.1. *If x is an admissible monomial of degree 15 in P_4 then either $\omega(x) = (1, 1, 1, 1)$ or $\omega(x) = (3, 2, 2)$.*

Proof. Since $\deg x$ is odd, we have $\omega_1(x) = 1$ or $\omega_1(x) = 3$.

Suppose $\omega_1(x) = 1$, then $x = x_i y^2$ with y a monomial of degree 7. Since x is admissible, by Theorem 2.7, y is admissible. If $y \notin P_4^+$ then from Kameko [10], $\omega(y) = (1, 1, 1)$ or $\omega(y) = (3, 2)$. A direct computation shows that $x = x_i y^2$ is inadmissible for all monomials y in P_4 with $\omega(y) = (3, 2)$. Hence $\omega(x) = (1, 1, 1, 1)$. If $y \in P_4^+$, then y is a permutation of one of the following monomial $x_1x_2x_3x_4^4, x_1x_2x_3^2x_4^3, x_1x_2^2x_3^2x_4^2$. By a direct computation we see that $x = x_i y^2$ is inadmissible.

If $\omega_1(x) = 3$, then $x = x_i x_j y^2, i < j$ with y a monomial of degree 6 in P_4 . By Theorem 2.7, y is admissible. So $\omega_1(y) = 2$ or $\omega_1(y) = 4$. If $\omega_1(y) = 4$, then by Proposition 2.8, x is inadmissible. Hence $\omega_1(y) = 2$ and $\omega(x) = (3, 2, 2)$. The lemma is proved. \square

Proposition 3.2. $(QP_4^+)_{15}$ is an \mathbb{F}_2 -vector space of dimension 37 with a basis consisting of all the classes represented by the admissible monomials $d_i, 1 \leq i \leq 37$, which are determined as follows:

- | | | | | |
|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| 1. $x_1x_2x_3^6x_4^7$ | 2. $x_1x_2x_3^7x_4^6$ | 3. $x_1x_2^2x_3^5x_4^7$ | 4. $x_1x_2^2x_3^7x_4^5$ | 5. $x_1x_2^3x_3^4x_4^7$ |
| 6. $x_1x_2^3x_3^5x_4^6$ | 7. $x_1x_2^3x_3^6x_4^5$ | 8. $x_1x_2^3x_3^7x_4^4$ | 9. $x_1x_2^6x_3x_4^7$ | 10. $x_1x_2^6x_3^3x_4^5$ |
| 11. $x_1x_2^6x_3^7x_4^4$ | 12. $x_1x_2^7x_3x_4^6$ | 13. $x_1x_2^7x_3^2x_4^5$ | 14. $x_1x_2^7x_3^3x_4^4$ | 15. $x_1x_2^7x_3^6x_4^3$ |
| 16. $x_1^3x_2x_3^4x_4^7$ | 17. $x_1^3x_2x_3^5x_4^6$ | 18. $x_1^3x_2x_3^6x_4^5$ | 19. $x_1^3x_2x_3^7x_4^4$ | 20. $x_1^3x_2^3x_3^4x_4^5$ |
| 21. $x_1^3x_2^3x_3^5x_4^4$ | 22. $x_1^3x_2^4x_3x_4^7$ | 23. $x_1^3x_2^4x_3^3x_4^5$ | 24. $x_1^3x_2^4x_3^7x_4^4$ | 25. $x_1^3x_2^5x_3x_4^6$ |
| 26. $x_1^3x_2^5x_3^2x_4^5$ | 27. $x_1^3x_2^5x_3^3x_4^4$ | 28. $x_1^3x_2^5x_3^6x_4^3$ | 29. $x_1^3x_2^7x_3x_4^4$ | 30. $x_1^3x_2^7x_3^4x_4^3$ |
| 31. $x_1^7x_2x_3x_4^6$ | 32. $x_1^7x_2x_3^2x_4^5$ | 33. $x_1^7x_2x_3^3x_4^4$ | 34. $x_1^7x_2x_3^6x_4^3$ | 35. $x_1^7x_2^3x_3x_4^4$ |
| 36. $x_1^7x_2^3x_3^4x_4^3$ | 37. $x_1x_2^2x_3^4x_4^8$ | | | |

Proof. From the proof of Lemma 3.1, if x is an admissible monomial of degree 15 in P_4 , then x is a permutation of one of the following monomials:

$$x_1x_2x_3^6x_4^7, x_1x_2^2x_3^5x_4^7, x_1x_2^3x_3^4x_4^7, x_1x_2^3x_3^5x_4^6, x_1^2x_2^3x_3^5x_4^5, x_1^3x_2^3x_3^4x_4^5.$$

By a direct computation we see that if $x \neq d_t, 1 \leq t \leq 37$, then x is inadmissible.

Now we prove that the set $\{[d_t] : 1 \leq t \leq 37\}$ is linearly independent in QP_4^+ . Suppose there is a linear relation

$$\mathcal{S} = \sum_{1 \leq t \leq 37} \gamma_t d_t \equiv 0, \quad (3.1)$$

with $\gamma_t \in \mathbb{F}_2$.

By Kameko [10], $B_3(15)$ is the set consisting of 7 monomials:

$$\begin{aligned} v_1 &= x_1 x_2^7 x_3^7, \quad v_2 = x_1^3 x_2^5 x_3^7, \quad v_3 = x_1^3 x_2^7 x_3^5, \\ v_4 &= x_1^7 x_2 x_3^7, \quad v_5 = x_1^7 x_2^3 x_3^5, \quad v_6 = x_1^7 x_2^7 x_3, \quad v_7 = x_1 x_2^2 x_3^{12}. \end{aligned}$$

By a direct computation, we explicitly compute $p_I(\mathcal{S})$ in terms of v_1, v_2, \dots, v_7 . From the relations $p_I(\mathcal{S}) \equiv 0$ for $I = (i, j)$ with $1 \leq i < j \leq 4$ and for $I = (1, i, j)$ with $2 \leq i < j \leq 4$, one gets $\gamma_t = 0$ for $t \neq 1, 2, 9, 11, 12, 15, 16, 19, 22, 24, 29, 30, 31$ and $\gamma_1 = \gamma_9 = \gamma_{16} = \gamma_{22}, \gamma_2 = \gamma_{11} = \gamma_{19} = \gamma_{24}, \gamma_{12} = \gamma_{15} = \gamma_{29} = \gamma_{30}, \gamma_{31} = \gamma_{34} = \gamma_{35} = \gamma_{36}$. Hence the relation (3.1) becomes

$$\gamma_1 \theta_1 + \gamma_2 \theta_2 + \gamma_{12} \theta_3 + \gamma_{31} \theta_4 + \gamma_{37} d_{37} \equiv 0, \quad (3.2)$$

where

$$\begin{aligned} \theta_1 &= d_1 + d_9 + d_{16} + d_{22}, \quad \theta_2 = d_2 + d_{11} + d_{19} + d_{24}, \\ \theta_3 &= d_{12} + d_{15} + d_{29} + d_{30}, \quad \theta_4 = d_{31} + d_{34} + d_{35} + d_{36}. \end{aligned}$$

Now, we prove that $\gamma_1 = \gamma_2 = \gamma_{12} = \gamma_{31} = 0$.

The proof is divided into 4 steps.

Step 1. Under the homomorphism φ_1 , the image of (3.2) is

$$\gamma_1 \theta_1 + \gamma_2 \theta_2 + \gamma_{12} \theta_3 + \gamma_{31} (\theta_4 + \theta_3) + \gamma_{37} (d_{37} + v_7) \equiv 0. \quad (3.3)$$

Since $v_7 \in P_4^0$, $\gamma_{37} = 0$. Combining (3.2) and (3.3), we get

$$\gamma_{31} \theta_3 \equiv 0. \quad (3.4)$$

If the polynomial θ_3 is hit, then we have

$$\theta_3 = Sq^1(A) + Sq^2(B) + Sq^4(C),$$

for some polynomials $A \in (P_4^+)_{14}, B \in (P_4^+)_{13}, C \in (P_4^+)_{11}$. Let $(Sq^2)^3$ act on the both sides of this equality. We get

$$(Sq^2)^3(\theta_3) = (Sq^2)^3 Sq^4(C),$$

By a direct calculation, we see that the monomial $x = x_1^8 x_2^7 x_3^4 x_4^2$ is a term of $(Sq^2)^3(\theta_3)$. If this monomial is a term of $(Sq^2)^3 Sq^4(y)$ for a monomial

$y \in (P_4^+)_{11}$, then $y = x_2^7 f_2(z)$ with $z \in P_3$ and $\deg z = 4$. Using the Cartan formula, we see that x is a term of $x_2^7 (Sq^2)^3 Sq^4(z) = x_2^7 (Sq^2)^3 (z^2) = 0$. Hence

$$(Sq^2)^3(\theta_3) \neq (Sq^2)^3 Sq^4(C),$$

for all $C \in (P_4^+)_{11}$ and we have a contradiction. So $[\theta_3] \neq 0$ and $\gamma_{31} = 0$.

Step 2. Since $\gamma_{31} = 0$, the homomorphism φ_2 sends (3.2) to

$$\gamma_1 \theta_1 + \gamma_2 \theta_2 + \gamma_{12} \theta_4 \equiv 0. \quad (3.5)$$

Using the relation (3.5) and by the same argument as given in Step 1, we get $\gamma_{12} = 0$.

Step 3. Since $\gamma_{31} = \gamma_{12} = 0$, the homomorphism φ_3 sends (3.2) to

$$\gamma_1 [\theta_1] + \gamma_2 [\theta_3] = 0. \quad (3.6)$$

Using the relation (3.6) and by the same argument as given in Step 2, we obtain $\gamma_3 = 0$.

Step 4. Since $\gamma_{31} = \gamma_{12} = \gamma_2 = 0$, the homomorphism φ_4 sends (3.2) to

$$\gamma_1 \theta_2 = 0.$$

Using this relation and by the same argument as given in Step 3, we obtain $\gamma_1 = 0$. The proposition is proved. \square

Corollary 3.3. *The set $[f(B_4(15))]$ is a basis of the \mathbb{F}_2 -vector space $(QP_5^0)_{15}$. Consequently $\dim(QP_5^0)_{15} = 270$.*

Now we compute $(QP_5)_5 = (QP_5^0)_5 \oplus (QP_5^+)_5$. Using Kameko's results in [10], we have $B_3(5) = \{x_1 x_2 x_3^3, x_1 x_2^3 x_3, x_1^3 x_2 x_3\}$. A direct computation, we easily obtain

$$B_4(5) = f(B_3(5)) \cup \{x_1 x_2^2 x_3 x_4, x_1 x_2 x_3^2 x_4, x_1 x_2 x_3 x_4^2\}.$$

This implies $\dim(QP_4)_5 = 15$. It is easy to see that $(QP_5^+)_5 = \langle [x_1 x_2 x_3 x_4 x_5] \rangle$. So we get

$$B_5(5) = f(B_4(5)) \cup \{x_1 x_2 x_3 x_4 x_5\}.$$

Combining this with Proposition 2.12 we obtain

Proposition 3.4. *The set $[B_5(5)]$ is a basis of the \mathbb{F}_2 -vector space $(QP_5)_5$. Consequently $\dim(QP_5)_5 = 46$.*

Now we compute $(QP_5^+)_5 \cap \text{Ker}(\widetilde{Sq}_*^0)^5$.

Lemma 3.5. *If x is an admissible monomial of degree 15 in P_5^+ and $[x] \in \text{Ker}(\widetilde{Sq}_*^0)$, then $\omega(x)$ is one of the sequences: $(1, 1, 3)$, $(3, 2, 2)$, $(3, 4, 1)$.*

Proof. Since $x \in P_5^+$ and $[x] \in \text{Ker}(\widetilde{Sq}_*^0)$, using Proposition 2.8, we see that x is a permutation of one of the following monomials:

$$\begin{array}{cccccc} x_1x_2x_3^2x_4^4x_5^7, & x_1x_2^2x_3^2x_4^3x_5^7, & x_1x_2x_3x_4^6x_5^6, & x_1x_2x_3^2x_4^5x_5^6, & x_1x_2x_3^3x_4^4x_5^6, \\ x_1x_2^2x_3^2x_4^4x_5^6, & x_1x_2^2x_3^3x_4^3x_5^6, & x_1^2x_2^2x_3^2x_4^3x_5^6, & x_1x_2^2x_3^2x_4^5x_5^5, & x_1x_2^2x_3^3x_4^4x_5^5, \\ x_1^2x_2^2x_3^3x_4^3x_5^5, & x_1x_2^2x_3^4x_4^4x_5^4, & x_1^2x_2^2x_3^3x_4^4x_5^4, & x_1x_2^3x_3^3x_4^4x_5^4. \end{array}$$

We have

$$\begin{aligned} x_1x_2^2x_3^2x_4^4x_5^6 &= x_1x_2^2x_3^2x_4^2x_5^8 + Sq^1(x_1^2x_2x_3x_4^4x_5^6 + x_2^2x_3x_4^2x_5^8) \\ &\quad + Sq^2(x_1x_2x_3x_4^4x_5^6 + x_1x_2x_3x_4^2x_5^8) \\ x_1^2x_2^2x_3^3x_4^4x_5^4 &= x_1x_2^2x_3^4x_4^4x_5^4 + Sq^1(x_1x_2^2x_3^3x_4^4x_5^4) \\ x_1^2x_2^2x_3^2x_4^3x_5^6 &= x_1x_2^2x_3^2x_4^4x_5^6 + Sq^1(x_1x_2^2x_3^3x_4^3x_5^6). \end{aligned}$$

Since $\omega(x_1x_2^2x_3^2x_4^2x_5^8) = (1, 3, 0, 1) < (1, 3, 2, 0) = \omega(x_1x_2^2x_3^2x_4^4x_5^6)$, $\omega(x_1x_2^2x_3^4x_4^4x_5^4) = (1, 1, 3) < (1, 3, 2) = \omega(x_1^2x_2^2x_3^3x_4^4x_5^4)$, $\omega(x_1x_2^2x_3^2x_4^4x_5^6) = (1, 3, 2) < (1, 5, 1) = \omega(x_1^2x_2^2x_3^2x_4^3x_5^6)$, if the monomial x is a permutation of one of the monomials $x_1x_2^2x_3^2x_4^4x_5^6$, $x_1^2x_2^2x_3^3x_4^4x_5^4$, $x_1^2x_2^2x_3^2x_4^3x_5^6$, then x is inadmissible. The lemma follows. \square

From Lemma 3.5, we have

$$\begin{aligned} (QP_5^+)_{15} \cap \text{Ker}(\widetilde{Sq}_*^0)_5^5 &= ((QP_5^+) \cap QP_5(1, 1, 3)) \oplus \\ &\quad \oplus ((QP_5^+) \cap QP_5(3, 4, 1)) \oplus ((QP_5^+) \cap QP_5(3, 2, 2)). \end{aligned}$$

Proposition 3.6. $QP_5^+ \cap QP_5(1, 1, 3) = \langle [x_1x_2^2x_3^4x_4^4x_5^4] \rangle$.

Proof. From the proof of Lemma 3.5, if x is a monomial of degree 15 in P_5 and $\omega(x) = (1, 1, 3)$ then x is a permutation of the monomial $x_1x_2^2x_3^4x_4^4x_5^4$. By a direct computation, we have $x \equiv x_1x_2^2x_3^4x_4^4x_5^4$, completing the proof. \square

Proposition 3.7. $QP_5^+ \cap QP_5(3, 4, 1)$ is an \mathbb{F}_2 -vector space of dimension 40 with a basis consisting of all the classes represented by the admissible monomials a_i , $1 \leq i \leq 40$, which are determined as follows:

1. $x_1x_2^2x_3^2x_4^3x_5^7$
2. $x_1x_2^2x_3^2x_4^7x_5^3$
3. $x_1x_2^2x_3^3x_4^2x_5^7$
4. $x_1x_2^2x_3^3x_4^3x_5^6$
5. $x_1x_2^2x_3^3x_4^6x_5^3$
6. $x_1x_2^2x_3^3x_4^7x_5^2$
7. $x_1x_2^2x_3^7x_4^2x_5^3$
8. $x_1x_2^2x_3^7x_4^3x_5^2$
9. $x_1x_2^3x_3^2x_4^2x_5^7$
10. $x_1x_2^3x_3^2x_4^3x_5^6$
11. $x_1x_2^3x_3^2x_4^6x_5^3$
12. $x_1x_2^3x_3^2x_4^7x_5^2$
13. $x_1x_2^3x_3^3x_4^2x_5^6$
14. $x_1x_2^3x_3^3x_4^6x_5^2$
15. $x_1x_2^3x_3^6x_4^2x_5^3$
16. $x_1x_2^3x_3^6x_4^3x_5^2$
17. $x_1x_2^3x_3^7x_4^2x_5^2$
18. $x_1x_2^3x_3^7x_4^3x_5^2$
19. $x_1x_2^7x_3^2x_4^3x_5^2$
20. $x_1x_2^7x_3^3x_4^2x_5^2$
21. $x_1^3x_2x_3^2x_4^2x_5^7$
22. $x_1^3x_2x_3^2x_4^3x_5^6$
23. $x_1^3x_2x_3^2x_4^6x_5^3$
24. $x_1^3x_2x_3^2x_4^7x_5^2$
25. $x_1^3x_2x_3^3x_4^2x_5^6$
26. $x_1^3x_2x_3^3x_4^6x_5^2$
27. $x_1^3x_2x_3^6x_4^2x_5^3$
28. $x_1^3x_2x_3^6x_4^3x_5^2$
29. $x_1^3x_2x_3^7x_4^2x_5^2$
30. $x_1^3x_2x_3^7x_4^3x_5^2$
31. $x_1^3x_2^3x_3x_4^2x_5^2$
32. $x_1^3x_2^3x_3x_4^3x_5^2$
33. $x_1^3x_2^5x_3^2x_4^2x_5^3$
34. $x_1^3x_2^5x_3^2x_4^3x_5^2$
35. $x_1^3x_2^5x_3^3x_4^2x_5^2$
36. $x_1^3x_2^5x_3^3x_4^3x_5^2$
37. $x_1^7x_2x_3^2x_4^2x_5^3$
38. $x_1^7x_2x_3^2x_4^3x_5^2$
39. $x_1^7x_2x_3^3x_4^2x_5^2$
40. $x_1^7x_2x_3^3x_4^3x_5^2$.

Proof. Let x be an admissible monomial of degree 15 in P_5 and $\omega(x) = (3, 4, 1)$. From the proof of Lemma 3.5, x is a permutation of one of the monomials $x_1x_2^2x_3^2x_4^7x_5^5$, $x_1x_2^2x_3^3x_4^6x_5^5$, $x_1^2x_2^2x_3^3x_4^5x_5^5$. A direct computation shows that if $x \neq a_t$, $1 \leq t \leq 40$, then x is inadmissible.

Now, we prove that the set $\{[a_t] : 1 \leq t \leq 40\}$ is linearly independent in QP_5 . Suppose there is a linear relation

$$\mathcal{S} = \sum_{1 \leq t \leq 40} \gamma_t a_t \equiv 0,$$

with $\gamma_t \in \mathbb{F}_2$. By a direct computation, we explicitly compute $p_{(1,j)}(\mathcal{S})$ in terms of d_i , $1 \leq j \leq 37$. From the relations $p_{(1,j)}(\mathcal{S}) \equiv 0$ for $1 \leq j \leq 5$, we obtain $\gamma_t = 0$ for $1 \leq t \leq 40$. The proposition is proved. \square

Proposition 3.8. $QP_5^+ \cap QP_5(3, 2, 2)$ is an \mathbb{F}_2 -vector space of dimension 75 with a basis consisting of all the classes represented by the admissible monomials b_t , $1 \leq t \leq 75$, which are determined as follows:

- | | | | |
|-------------------------------|-------------------------------|-------------------------------|-------------------------------|
| 1. $x_1x_2x_3x_4^6x_5^6$ | 2. $x_1x_2x_3^2x_4^7x_5^7$ | 3. $x_1x_2x_3^2x_4^5x_5^6$ | 4. $x_1x_2x_3^2x_4^6x_5^5$ |
| 5. $x_1x_2x_3^2x_4^7x_5^4$ | 6. $x_1x_2x_3^3x_4^6x_5^6$ | 7. $x_1x_2x_3^3x_4^6x_5^4$ | 8. $x_1x_2x_3^6x_4x_5^6$ |
| 9. $x_1x_2x_3^6x_4^2x_5^5$ | 10. $x_1x_2x_3^6x_4^3x_5^4$ | 11. $x_1x_2x_3^6x_4x_5$ | 12. $x_1x_2x_3^7x_4^2x_5^4$ |
| 13. $x_1x_2^2x_3x_4^4x_5^7$ | 14. $x_1x_2^2x_3x_4^5x_5^6$ | 15. $x_1x_2^2x_3x_4^6x_5^5$ | 16. $x_1x_2^2x_3x_4^7x_5^4$ |
| 17. $x_1x_2^2x_3^3x_4^4x_5^5$ | 18. $x_1x_2^2x_3^3x_4^5x_5^4$ | 19. $x_1x_2^2x_3^4x_4^7x_5^5$ | 20. $x_1x_2^2x_3^4x_4^3x_5^5$ |
| 21. $x_1x_2^2x_3^4x_4^7x_5$ | 22. $x_1x_2^2x_3^4x_4^6x_5^5$ | 23. $x_1x_2^2x_3^4x_4^2x_5^5$ | 24. $x_1x_2^2x_3^4x_4^3x_5^4$ |
| 25. $x_1x_2^2x_3^4x_4^6x_5$ | 26. $x_1x_2^2x_3^4x_4^4x_5^4$ | 27. $x_1x_2^2x_3^4x_4^4x_5$ | 28. $x_1x_2^2x_3^4x_4^6x_5^5$ |
| 29. $x_1x_2^2x_3^4x_4^6x_5^4$ | 30. $x_1x_2^2x_3^4x_4^2x_5^5$ | 31. $x_1x_2^2x_3^4x_4^2x_5^4$ | 32. $x_1x_2^2x_3^4x_4^4x_5^4$ |
| 33. $x_1x_2^2x_3^4x_4^6x_5$ | 34. $x_1x_2^2x_3^4x_4^2x_5^5$ | 35. $x_1x_2^2x_3^4x_4^3x_5^4$ | 36. $x_1x_2^2x_3^4x_4^6x_5$ |
| 37. $x_1x_2^2x_3^5x_4^2x_5^4$ | 38. $x_1x_2^2x_3^6x_4x_5^4$ | 39. $x_1x_2^2x_3^6x_4x_5$ | 40. $x_1x_2^2x_3^6x_4x_5^6$ |
| 41. $x_1x_2^2x_3^6x_4^2x_5^5$ | 42. $x_1x_2^2x_3^6x_4^3x_5^4$ | 43. $x_1x_2^2x_3^6x_4x_5$ | 44. $x_1x_2^2x_3^6x_4x_5^4$ |
| 45. $x_1x_2^2x_3^6x_4x_5$ | 46. $x_1x_2^2x_3^6x_4^2x_5^4$ | 47. $x_1x_2^2x_3^6x_4x_5^4$ | 48. $x_1x_2^2x_3^6x_4^2x_5$ |
| 49. $x_1^3x_2x_3^4x_4^6x_5$ | 50. $x_1^3x_2x_3^4x_4^6x_5^4$ | 51. $x_1^3x_2x_3^4x_4^5x_5^5$ | 52. $x_1^3x_2x_3^4x_4^5x_5^4$ |
| 53. $x_1^3x_2x_3^4x_4^4x_5^5$ | 54. $x_1^3x_2x_3^4x_4^6x_5$ | 55. $x_1^3x_2x_3^4x_4^2x_5^5$ | 56. $x_1^3x_2x_3^4x_4^3x_5^4$ |
| 57. $x_1^3x_2x_3^4x_4^6x_5$ | 58. $x_1^3x_2x_3^5x_4^2x_5^4$ | 59. $x_1^3x_2x_3^6x_4x_5^4$ | 60. $x_1^3x_2x_3^6x_4x_5$ |
| 61. $x_1^3x_2x_3^3x_4^4x_5^4$ | 62. $x_1^3x_2x_3^4x_4^4x_5^4$ | 63. $x_1^3x_2x_3^4x_4^4x_5$ | 64. $x_1^3x_2x_3^4x_4^6x_5^5$ |
| 65. $x_1^3x_2x_3^4x_4^2x_5^5$ | 66. $x_1^3x_2x_3^4x_4^3x_5^4$ | 67. $x_1^3x_2x_3^4x_4^6x_5$ | 68. $x_1^3x_2x_3^4x_4^4x_5^4$ |
| 69. $x_1^3x_2x_3^4x_4^4x_5$ | 70. $x_1^3x_2x_3^5x_4^2x_5^4$ | 71. $x_1^3x_2x_3^5x_4x_5^4$ | 72. $x_1^3x_2x_3^5x_4^2x_5$ |
| 73. $x_1^7x_2x_3x_4^2x_5^4$ | 74. $x_1^7x_2x_3^2x_4x_5^4$ | 75. $x_1^7x_2x_3^2x_4x_5$ | |

Proof. Let x be an admissible monomial of degree 15 in P_5 and $\omega(x) = (3, 2, 2)$. From the proof of Lemma 3.5, x is a permutation of one of the monomials:

$$\begin{array}{llll} x_1x_2x_3^2x_4^4x_5^7, & x_1x_2x_3x_4^6x_5^6, & x_1x_2x_3^2x_4^5x_5^6, & x_1x_2x_3^3x_4^4x_5^6, \\ x_1x_2^2x_3^2x_4^5x_5^5, & x_1x_2^2x_3^3x_4^4x_5^5, & x_1x_2^2x_3^3x_4^4x_5^4. \end{array}$$

By a direct computation, we see that if $x \neq b_t$, $1 \leq t \leq 75$, then x is inadmissible.

Now, we prove that the set $\{[b_t] : 1 \leq t \leq 75\}$ is linearly independent in QP_5 . Suppose there is a linear relation

$$\mathcal{S} = \sum_{1 \leq t \leq 75} \gamma_t b_t \equiv 0, \quad (3.7)$$

with $\gamma_t \in \mathbb{F}_2$. By a direct computation, we explicitly compute $p_{(i,j)}(\mathcal{S})$ in terms of d_t , $1 \leq t \leq 37$. From the relations $p_{(i,j)}(\mathcal{S}) \equiv 0$ for $1 \leq i < j \leq 5$, one gets $\gamma_t = 0$ for $t \notin J$ with

$$J = \{1, 8, 11, 32, 38, 39, 40, 43, 44, 45, 49, 50, 53, 54, 57, 61, 62, 63, 64, 67, 68, 69\}$$

and $\gamma_t = \gamma_1$ for $t \in J$. Hence the relation (3.7) becomes

$$\gamma_1 q \equiv 0,$$

where $q = b_1 + b_8 + b_{11} + b_{32} + b_{38} + b_{39} + b_{40} + b_{43} + b_{44} + b_{45} + b_{49} + b_{50} + b_{53} + b_{54} + b_{57} + b_{61} + b_{62} + b_{63} + b_{64} + b_{67} + b_{68} + b_{69}$.

If the polynomial q is hit, then we have

$$q = Sq^1(A) + Sq^2(B) + Sq^4(C),$$

for some polynomials $A \in (P_5^+)_{14}$, $B \in (P_5^+)_{13}$, $C \in (P_5^+)_{11}$. Let $(Sq^2)^3$ act on the both sides of this equality. Since $(Sq^2)^3 Sq^1 = 0$ and $(Sq^2)^3 Sq^2 = 0$ we get

$$(Sq^2)^3(q) = (Sq^2)^3 Sq^4(C).$$

By a direct calculation, we have

$$(Sq^2)^3(q) = D + \text{other terms},$$

where $D = x_1^3(x_2^2 x_3^8 x_4^4 x_5^4 + x_2^8 x_3^2 x_4^4 x_5^4 + x_2^8 x_3^4 x_4^2 x_5^4 + x_2^8 x_3^4 x_4^4 x_5^2 + x_2^4 x_3^8 x_4^2 x_5^4 + x_2^8 x_3^4 x_4^4 x_5^2 + x_2^6 x_3^4 x_4^4 x_5^4 + x_2^4 x_3^6 x_4^4 x_5^4)$. Hence there is a polynomial $C' \in (P_4)_8$ such that D is a term of $(Sq^2)^3 Sq^4(x_1^3 f_1(C'))$. Using the Cartan formula we see that D is a term of $x_1^3 f_1((Sq^2)^3 Sq^4(C'))$. A direct computation shows that D is not a term of $x_1^3 f_1((Sq^2)^3 Sq^4(C'))$ for any $C' \in (P_4)_8$. Hence

$$(Sq^2)^3(q) \neq (Sq^2)^3 Sq^4(C),$$

for all $C \in (P_5^+)_{11}$ and we have a contradiction. So $[q] \neq 0$ and $\gamma_1 = 0$. The proposition is proved. \square

4 Proof of Theorems 1.2 and 1.3

Proof of Theorem 1.2. Since $\widetilde{Sq}_*^0 = (\widetilde{Sq}_*^0)_{15}^5 : (QP_5)_{15} \rightarrow (QP_5)_5$ is a homomorphism of GL_5 -modules, we have a direct summand decomposition of the GL_5 -modules: $(QP_5)_{15} = \text{Ker}(\widetilde{Sq}_*^0)_{15}^5 \oplus (QP_5)_5$. Hence

$$(QP_5)_{15}^{GL_5} = (\text{Ker}(\widetilde{Sq}_*^0)_{15}^5)^{GL_5} \oplus (QP_5)_5^{GL_5}.$$

By a direct computation using Proposition 3.4 we easily obtain $(QP_5)_5^{GL_5} = 0$.

It is easy to see that

$$\text{Ker}(\widetilde{Sq}_*^0)_5^5 = QP_5(1, 1, 1, 1) \oplus QP_5(1, 1, 3) \oplus QP_5(3, 2, 2) \oplus QP_5(3, 4, 1),$$

where $QP_5(1, 1, 1, 1) \oplus QP_5(1, 1, 3)$, $QP_5(3, 2, 2)$ and $QP_5(3, 4, 1)$ are the GL_5 -submodules of $\text{Ker}(\widetilde{Sq}_*^0)_5^5$. By a direct computation using Theorem 1.1 and the homomorphisms $\varphi_i : QP_5 \rightarrow QP_5$, $1 \leq i \leq 5$, one gets

$$\begin{aligned} (QP_5(1, 1, 1, 1) \oplus QP_5(1, 1, 3))^{GL_5} &= \langle [p] \rangle, \\ QP_5(3, 2, 2)^{GL_5} &= \langle [q] \rangle, \quad QP_5(3, 4, 1)^{GL_5} = 0. \end{aligned}$$

The theorem is proved. \square

Proof of Theorem 1.3. First of all, we briefly recall the definition of the Singer transfer. Let \widehat{P}_1 be the submodule of $\mathbb{F}_2[x_1, x_1^{-1}]$ spanned by all powers x_1^i with $i \geq -1$. The usual \mathcal{A} -action on $P_1 = \mathbb{F}_2[x_1]$ is canonically extended to an \mathcal{A} -action on $\mathbb{F}_2[x_1, x_1^{-1}]$ (see Singer [20]). \widehat{P}_1 is an \mathcal{A} -submodule of $\mathbb{F}_2[x_1, x_1^{-1}]$. The inclusion $P_1 \subset \widehat{P}_1$ gives rise to a short exact sequence of \mathcal{A} -modules:

$$0 \longrightarrow P_1 \longrightarrow \widehat{P}_1 \longrightarrow \Sigma^{-1}\mathbb{F}_2 \longrightarrow 0.$$

Let e_1 be the corresponding element in $\text{Ext}_{\mathcal{A}}^1(\Sigma^{-1}\mathbb{F}_2, P_1)$. Singer set $e_k = e_1 \otimes \dots \otimes e_1 \in \text{Ext}_{\mathcal{A}}^k(\Sigma^{-k}\mathbb{F}_2, P_k)$. Then, he defined $\text{Tr}_k^* : \text{Tor}_k^{\mathcal{A}}(\mathbb{F}_2, \Sigma^{-k}\mathbb{F}_2) \rightarrow \text{Tor}_0^{\mathcal{A}}(\mathbb{F}_2, P_k) = QP_k$ by $\text{Tr}_k^*(z) = e_k \cap z$. Its image is a submodule of $(QP_k)^{GL_k}$. The k -th Singer transfer is defined to be the dual of Tr_k^* .

The algebra $\text{Ext}_{\mathcal{A}}^{*,*}(\mathbb{F}_2, \mathbb{F}_2)$ is described in terms of the mod-2 lambda algebra Λ (see Lin [12]). Recall that Λ is a bigraded differential algebra over \mathbb{F}_2 generated by $\lambda_j \in \Lambda^{1,j}$, $j \geq 0$, with the relations

$$\lambda_j \lambda_{2j+1+m} = \sum_{\nu \geq 0} \binom{m - \nu - 1}{\nu} \lambda_{j+m-\nu} \lambda_{2j+1+\nu},$$

for $m \geq 0$ and the differential

$$\delta(\lambda_k) = \sum_{\nu \geq 0} \binom{k - \nu - 1}{\nu + 1} \lambda_{k-\nu-1} \lambda_{\nu},$$

for $k > 0$ and that $H^{s,t}(\Lambda, \delta) = \text{Ext}_{\mathcal{A}}^{s,t+s}(\mathbb{F}_2, \mathbb{F}_2)$. It is easy to see that $\lambda_{2^i-1} \in \Lambda^{1,2^i-1}$, $i \geq 0$, and $\bar{d}_0 = \lambda_6 \lambda_2 \lambda_3^2 + \lambda_4^2 \lambda_3^2 + \lambda_2 \lambda_4 \lambda_5 \lambda_3 + \lambda_1 \lambda_5 \lambda_1 \lambda_7 \in \Lambda^{4,14}$ are the cycles in the lambda algebra Λ .

Proposition 4.1 (See Lin [12]). $\text{Ext}_{\mathcal{A}}^{5,20}(\mathbb{F}_2, \mathbb{F}_2) = \text{Span}\{h_0^4 h_4, h_1 d_0\}$, with $h_i = [\lambda_{2^i-1}] \in \text{Ext}_{\mathcal{A}}^{1,2^i}(\mathbb{F}_2, \mathbb{F}_2)$ and $d_0 = [\bar{d}_0] \in \text{Ext}_{\mathcal{A}}^{4,18}(\mathbb{F}_2, \mathbb{F}_2)$.

It is well known that $H_*(B\mathbb{V}_k)$ is the dual of $H^*(B\mathbb{V}_k) = P_k$. So

$$H_*(B\mathbb{V}_k) = \Gamma(a_1, a_2, \dots, a_k)$$

is the divided power algebra generated by a_1, a_2, \dots, a_k , where a_i is dual to $x_i \in P_k$ with respect to the basis of P_k consisting of all monomials in x_1, x_2, \dots, x_k . In [3], Chon and Hà defined a homomorphism of algebras

$$\phi = \bigoplus_{k \geq 1} \phi_k : \bigoplus_{k \geq 1} H_*(B\mathbb{V}_k) \rightarrow \bigoplus_{k \geq 1} \Lambda_k = \Lambda,$$

which induces the Singer transfer. Here the homomorphism $\phi_k : H_*(B\mathbb{V}_k) \rightarrow \Lambda_k$ is defined by the following inductive formula:

$$\phi_k(a^{(I,t)}) = \begin{cases} \lambda_t, & \text{if } k-1 = \ell(I) = 0, \\ \sum_{i \geq t} \phi_{k-1}(Sq^{i-t}a^I)\lambda_i, & \text{if } k-1 = \ell(I) > 0, \end{cases}$$

for any $a^{(I,t)} = a_1^{(i_1)} a_1^{(i_2)} \dots a_{k-1}^{(i_{k-1})} a_k^{(t)} \in H_*(B\mathbb{V}_k)$ and $I = (i_1, i_2, \dots, i_{k-1})$.

Proposition 4.2 (See Chon and Hà [3]). *If $b \in PH_*(BV_k)$, then $\phi_k(b)$ is a cycle in the lambda algebra Λ and $\text{Tr}_k([b]) = [\phi_k(b)]$.*

Now we are ready to prove Theorem 1.3.

According to Theorem 1.2, $\{[p], [q]\}$ is a basis of $(QP_5)_{15}^{GL_5}$. Let $\{p^*, q^*\}$ be the basis of $\mathbb{F}_2 \otimes_{GL_5} PH_{15}(B\mathbb{V}_5)$ which is dual to $\{[p], [q]\}$. It is easy to see that $a_5^{(15)} \in PH_{15}(B\mathbb{V}_5)$ and $\langle a_5^{(15)}, p \rangle = 1, \langle a_5^{(15)}, q \rangle = 0$. Consider the element $b = \sum_{I \in \mathcal{J}} a^I \in H_{15}(B\mathbb{V}_5)$, where \mathcal{J} is the set of all the following sequences:

(1, 1, 1, 6, 6), (1, 2, 2, 5, 5), (1, 2, 1, 6, 5), (1, 1, 2, 5, 6), (1, 4, 2, 5, 3), (1, 4, 1, 6, 3),
 (1, 3, 2, 6, 3), (1, 2, 4, 3, 5), (1, 1, 4, 3, 6), (1, 4, 4, 3, 3), (1, 6, 1, 1, 6), (1, 5, 2, 2, 5),
 (1, 6, 1, 2, 5), (1, 5, 2, 1, 6), (1, 5, 2, 4, 3), (1, 6, 1, 4, 3), (1, 6, 2, 3, 3), (1, 3, 4, 2, 5),
 (1, 3, 4, 1, 6), (1, 3, 3, 2, 6), (1, 3, 4, 4, 3), (1, 1, 6, 1, 6), (1, 2, 5, 2, 5), (1, 2, 6, 1, 5),
 (1, 1, 5, 2, 6), (1, 4, 5, 2, 3), (1, 4, 6, 1, 3), (1, 3, 6, 2, 3), (1, 2, 3, 4, 5), (1, 1, 3, 4, 6),
 (1, 4, 3, 4, 3), (1, 3, 1, 5, 5), (1, 5, 5, 1, 3), (1, 5, 1, 3, 5), (1, 5, 3, 1, 5), (1, 5, 3, 3, 3).

By a direct computation we see that $b \in PH_{15}(B\mathbb{V}_5)$ and $\langle b, p \rangle = 0, \langle b, q \rangle = 1$. Hence we obtain $[a_5^{(15)}] = p^*$ and $[b] = q^*$. A direct computation shows

$$\begin{aligned} \phi_5(a_5^{(15)}) &= \lambda_0^4 \lambda_{15}, \\ \phi_5(b) &= \lambda_1 \bar{d}_0 + \delta(\lambda_1 \lambda_9 \lambda_3^2 + \lambda_1 \lambda_3 \lambda_9 \lambda_3). \end{aligned}$$

Using Proposition 4.2, one gets $\text{Tr}_5(p^*) = \text{Tr}_5([a_5^{(15)}]) = h_0^4 h_4$ and $\text{Tr}_5(q^*) = \text{Tr}_5([b]) = h_1 d_0$. The theorem follows. \square

Acknowledgment. The paper was completed when the author was visiting to Vietnam Institute for Advanced Study in Mathematics (VIASM). The author would like to thank the VIASM for support and kind hospitality. His thanks go to Dr. Phan Hoàng Chon for helpful discussions. This work was also partially supported by the Reseach Project No. B2013.28.129.

References

- [1] J. M. Boardman, *Modular representations on the homology of power of real projective space*, in: M. C. Tangora (Ed.), *Algebraic Topology*, Oaxtepec, 1991, in: *Contemp. Math.*, vol. 146, 1993, pp. 49-70, MR1224907.
- [2] R. R. Bruner, L. M. Hà and N. H. V. Hung, *On behavior of the algebraic transfer*, *Trans. Amer. Math. Soc.* 357 (2005), 473-487, MR2095619.
- [3] P. H. Chon and L. M. Hà, *Lambda algebra and the Singer transfer*, *C. R. Math. Acad. Sci. Paris* 349 (2011), 21-23, MR2755689.
- [4] P. H. Chon and L. M. Hà, *On May spectral sequence and the algebraic transfer*, *Manuscripta Math.* 138 (2012), 141-160, MR2898751.
- [5] M. C. Crabb and J. R. Hubbuck, *Representations of the homology of BV and the Steenrod algebra II*, *Algebraic Topology: new trend in localization and periodicity*, *Progr. Math.* 136 (1996), 143-154, MR1397726.
- [6] L. M. Hà, *Sub-Hopf algebras of the Steenrod algebra and the Singer transfer*, "Proceedings of the International School and Conference in Algebraic Topology, Hà Nội 2004", *Geom. Topol. Monogr., Geom. Topol. Publ.*, Coventry, vol. 11 (2007), 81-105, MR2402802.
- [7] N. H. V. Hung, *Spherical classes and the algebraic transfer*, *Trans. Amer. Math. Soc.* 349 (1997), 3893-3910, MR1433119.
- [8] N. H. V. Hung, *The cohomology of the Steenrod algebra and representations of the general linear groups*, *Trans. Amer. Math. Soc.* 357 (2005), 4065-4089, MR2159700.
- [9] N. H. V. Hung and V. T. N. Quỳnh, *The image of Singer's fourth transfer*, *C. R. Math. Acad. Sci. Paris* 347 (2009), 1415-1418, MR2588792.
- [10] M. Kameko, *Products of projective spaces as Steenrod modules*, PhD. Thesis, Johns Hopkins University, 1990.
- [11] M. Kameko, *Generators of the cohomology of BV_3* , *J. Math. Kyoto Univ.* 38 (1998), 587-593, MR1661173.
- [12] W. H. Lin, $\text{Ext}_A^{4,*}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $\text{Ext}_A^{5,*}(\mathbb{Z}/2, \mathbb{Z}/2)$, *Topology Appl.*, 155 (2008), 459-496, MR2380930.
- [13] T. N. Nam, *\mathcal{A} -générateurs génériques pour l'algèbre polynomiale*, *Adv. Math.* 186 (2004), 334-362, MR2073910.
- [14] T. N. Nam, *Transfert algébrique et action du groupe linéaire sur les puissances divisées modulo 2*, *Ann. Inst. Fourier (Grenoble)* 58 (2008), 1785-1837, MR2445834.
- [15] F. P. Peterson, *Generators of $H^*(\mathbb{R}P^\infty \times \mathbb{R}P^\infty)$ as a module over the Steenrod algebra*, *Abstracts Amer. Math. Soc. No.* **833** April 1987.
- [16] S. Priddy, *On characterizing summands in the classifying space of a group, I*, *Amer. Jour. Math.* 112 (1990), 737-748, MR1073007.
- [17] V. T. N. Quỳnh, *On behavior of the fifth algebraic transfer*, "Proceedings of the International School and Conference in Algebraic Topology, Hà Nội 2004", *Geom. Topol. Monogr., Geom. Topol. Publ.*, Coventry, vol. 11 (2007), 309-326, MR2402811.
- [18] J. Repka and P. Selick, *On the subalgebra of $H_*((\mathbb{R}P^\infty)^n; \mathbb{F}_2)$ annihilated by Steenrod operations*, *J. Pure Appl. Algebra* 127 (1998), 273-288, MR1617199.

- [19] J. H. Silverman, *Hit polynomials and the canonical antiautomorphism of the Steenrod algebra*, Proc. Amer. Math. Soc. 123 (1995), 627-637, MR1254854.
- [20] W. M. Singer, *The transfer in homological algebra*, Math. Zeit. 202 (1989), 493-523, MR1022818.
- [21] W. M. Singer, *On the action of the Steenrod squares on polynomial algebras*, Proc. Amer. Math. Soc. 111 (1991), 577-583, MR1045150.
- [22] N. E. Steenrod and D. B. A. Epstein, *Cohomology operations*, Annals of Mathematics Studies 50, Princeton University Press, Princeton N.J (1962), MR0145525.
- [23] N. Sum, *The negative answer to Kameko's conjecture on the hit problem*, C. R. Math. Acad. Sci. Paris, Ser. I 348 (2010), 669-672.
- [24] N. Sum, *The negative answer to Kameko's conjecture on the hit problem*, Adv. Math. 225 (2010), 2365-2390.
- [25] N. Sum, *On the hit problem for the polynomial algebra*, C. R. Math. Acad. Sci. Paris, Ser. I, 351 (2013), 565-568.
- [26] R. M. W. Wood, *Steenrod squares of polynomials and the Peterson conjecture*, Math. Proc. Cambriges Phil. Soc. 105 (1989), 307-309, MR0974986.
- [27] R. M. W. Wood, *Problems in the Steenrod algebra*, Bull. London Math. Soc. 30 (1998) 449-517, MR1643834.